

Iowa Influenza Surveillance Network (IISN)

Influenza-like Illness (ILI) and Other Respiratory Viruses

Weekly Activity Report



All data presented in this report are provisional and may change as additional reports are received



Iowa Influenza Geographic Spread						
No Activity	Sporadic	Local	Regional	Widespread		

Note: See CDC activity estimates for definition www.cdc.gov/flu/weekly/overview.htm

Quick Stats	
Predominate influenza subtype	A(H3)
Percent of influenza rapid test positive	33% (1217/3672)
Percent of RSV rapid tests positive	11% (54/487)
Influenza-associated hospitalizations	68/2274 inpatients surveyed
Percent of outpatient visits for ILI	4.49% (baseline 1.6%)
Percent school absence due to illness	3.21%
Number of long-term care outbreaks	9
Number of schools with ≥10% absence due to illness	55
Influenza-associated mortality - all ages (Cumulative)	34
Influenza-associated pediatric mortality (Cumulative)	1
Predominate non-influenza virus	Rhinovirus/enterovirus

Note: Deaths are considered influenza-associated when influenza is listed on the death certificate. This is an underestimate of influenza-related deaths. Cumulative mortality totals are from 9/30/2018 to the current week.

Iowa statewide activity summary:


Influenza activity is still elevated on multiple measures, and the geographic spread of influenza in Iowa is widespread. Influenza A(H3) was the predominate subtype detected at the State Hygienic Laboratory this week with 62 influenza A(H3), 45 influenza A(H1N1)pdm09 and two influenza B (Victoria lineage) viruses detected from submitted samples as well as two influenza A positive specimens with no subtype reported. Sentinel hospitals reported 68 influenza-related hospitalizations. The proportion of outpatient visits due to influenza-like illness (ILI) decreased slightly to 4.49 percent, which is above the regional baseline of 1.6 percent. Nine long-term care influenza outbreaks were reported with onset in week 11. Fifty-five schools reported at least 10 percent absenteeism due to illness. Nine influenza deaths were reported. Surveillance sites most frequently detected the following non-influenza respiratory illnesses: 139 rhinovirus/enterovirus 129 coronavirus and 80 RSV.

International activity summary - (WHO):

In the temperate zone of the northern hemisphere influenza activity continued to be reported. In North America, influenza activity continued but in recent weeks influenza A(H3N2) was the dominant virus, followed by influenza A(H1N1)pdm09. In the temperate zones of the southern hemisphere, influenza activity remained at inter-seasonal levels, with the exception of some parts of Australia where influenza activity remained above inter-seasonal levels. Worldwide, seasonal influenza A viruses accounted for the majority of detections.

Visit <u>www.who.int/influenza/surveillance monitoring/updates/latest update GIP surveillance/en/</u> for more information. It was last updated 3/18/2019.

National activity summary - (CDC)-Last Updated for Week 11:



Synopsis: Influenza activity remains elevated in the United States. Influenza A(H1N1)pdm09, influenza A(H3N2), and influenza B viruses continue to co-circulate. Below is a summary of the key influenza indicators for the week ending March 16, 2019.

Viral Surveillance: The percentage of respiratory specimens testing positive for influenza viruses in clinical laboratories increased slightly. Nationally, during the most recent three weeks, influenza A(H3) viruses were reported more frequently than influenza A(H1N1)pdm09 viruses and in HHS Regions 2, 4, 5, 6, 7, 8, 9 and 10. The majority of influenza viruses characterized antigenically are similar to the cell-grown reference viruses representing the 2018–2019 Northern Hemisphere influenza vaccine viruses. However, an increasing proportion of influenza A(H3N2) viruses are antigenically distinguishable from A/Singapore/INFIMH-16-0019/2016 (3C.2a1), a cell-propagated reference virus representing the A(H3N2) component of 2018-19 Northern Hemisphere influenza vaccines. The vast majority of influenza viruses tested (>99%) show susceptibility to oseltamivir and peramivir. All influenza viruses tested showed susceptibility to zanamivir.

Outpatient Illness Surveillance: The proportion of outpatient visits for influenza-like illness (ILI) remained at 4.4%, which is above the national baseline of 2.2%. All 10 regions reported ILI at or above their region-specific baseline level.

Geographic Spread of Influenza: The geographic spread of influenza in 44 states was reported as widespread; Puerto Rico and four states reported regional activity; the District of Columbia and two states reported local activity; the U.S. Virgin Islands reported sporadic activity; and Guam did not report.

Influenza-associated Hospitalizations: A cumulative rate of 47.1 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported. The highest hospitalization rate is among adults 65 years and older (146.0 hospitalizations per 100,000 population).

Pneumonia and Influenza Mortality: The proportion of deaths attributed to pneumonia and influenza (P&I) was below the system-specific epidemic threshold in the National Center for Health Statistics (NCHS) Mortality Surveillance System.

Influenza-associated Pediatric Deaths: Eight influenza-associated pediatric deaths were reported to CDC during week 11.

Laboratory surveillance program:

The State Hygienic Laboratory (SHL) is the primary laboratory in Iowa characterizing specimens for influenza surveillance. SHL reports the number of tests performed and the type and subtype/lineage of positive tests to the influenza surveillance network daily. SHL also sends a portion of specimens to CDC for further characterization.


Table 1: Influenza A viruses detected by SHL by age group

		CURREN	IT WEEK		CUMI	JLATIVE (9/30/	18- CURRENT	Γ WEEK)
Age Group	A(H1N1) pdm09	A(H3)	Not subtyped	Total	A(H1N1) pdm09	A(H3)	Not subtyped	Total
0-4	12	4	0	16 (15%)	60	13	0	73 (10%)
5-17	5	11	0	16 (15%)	68	41	0	109 (15%)
18-24	2	14	1	17 (16%)	27	90	2	119 (17%)
25-49	13	3	0	16 (15%)	83	22	2	107 (15%)
50-64	7	5	0	12 (11%)	86	34	3	123 (17%)
>64	6	25	1	32 (29%)	63	111	10	184 (26%)
Total	45	62	2	109	387	311	17	715
Pct.	41%	57%	2%		54%	43%	2%	

Table 2: Influenza B viruses detected by SHL by age group

	CURRENT WEEK				CUMI	JLATIVE (9/30/	18– CURRENT	WEEK)
Age Group	Victoria Lineage	Yamagata Lineage	Not subtyped	Total	Victoria Lineage	Yamagata Lineage	Not subtyped	Total
0-4	0	0	0	0 (0%)	0	0	0	0 (0%)
5-17	0	0	0	0 (0%)	5	3	0	8 (53%)
18-24	1	0	0	1 (50%)	1	0	0	1 (7%)
25-49	1	0	0	1 (50%)	2	2	0	4 (27%)
50-64	0	0	0	0 (0%)	1	0	0	1 (7%)
>64	0	0	0	0 (0%)	0	0	1	1 (7%)
Total	2	0	0	2	9	5	1	15
Pct.	100%	0%	0%		60%	33%	7%	

Table 1 and 2 Notes: Cell counts of three or less are sometimes suppressed to protect confidentiality. Totals by age may not add up to totals by subtype/lineage due to missing age information. Only cases of lowa residents are included. Specimens listed as "not subtyped" may be pending or were not able to be subtyped due to weak positive lab results. This can be due to poor collection, timing of collection or stage of infection.




Rapid influenza and RSV test surveillance:


The State Hygienic Laboratory (SHL) runs a weekly web-based survey program where laboratorians report the number of influenza and respiratory syncytial virus (RSV) rapid tests performed and the number of tests positive. This table includes only the number of patients tested for influenza or RSV at laboratory surveillance sites throughout the state. This table does not provide case counts.

Table 3: Percent of influenza rapid tests positive and number of tests performed by region for the present week

REGION	RAF	RAPID ANTIGEN INFLUENZA TESTS			RAPID ANTIGEN RSV TESTS		
REGION	Tested	Flu A	Flu B	% Positive	Tested	Positive	% Positive
Region 1 (Central)	1414	517	18	38	83	10	12
Region 2 (NE)	149	48	0	32	27	1	4
Region 3 (NW)	363	64	3	18	127	12	9
Region 4 (SW)	141	29	1	21	26	6	23
Region 5 (SE)	182	43	0	24	37	8	22
Region 6 (Eastern)	1423	487	7	35	187	17	9
Total	3672	1188	29	33	487	54	11

Note: see map in the school section for the counties in each region.







Influenza-associated hospitalizations:

Sentinel hospitals that participate in IISN voluntarily track and report the number of influenza-associated hospitalizations and the total number of inpatients each week. Iowa hospitals interested in joining this surveillance program should contact Andy Weigel at 515-322-1937 or andy.weigel@idph.iowa.gov for more information.

Table 4: Number of influenza-associated hospitalization reported by age group

AGE	CURRENT WEEK	CUMULATIVE (9/30/18- CURRENT WEEK)
Age 0-4	6	54
Age 5-24	6	38
Age 25-49	13	74
Age 50-64	13	140
Age >64	30	314
Total	68	620






Outpatient health care provider surveillance program (ILINet):

Outpatient health care providers who participate in the ILINet (a national influenza surveillance program) report the number of patients seen with influenza-like illness and the total number of patient visits each week. This system is a key part of Iowa's influenza surveillance. Iowa health care providers interested in joining this surveillance program should contact Andy Weigel at 515-322-1937 or andy.weigel@idph.iowa.gov for more information.

Table 5: Outpatient visits for influenza-like illness (ILI)


Week	Age 0-4	Age 5-24	Age 25-49	Age 50-64	Age >64	Total ILI	ILI Percent
Week 9, ending Mar 2	48	128	15	9	13	213	3.07
Week 10, ending Mar 9	67	151	17	10	3	248	4.54
Week 11, ending Mar 16	79	99	27	12	14	231	4.49

Note: Influenza-like Illness is defined as a fever of ≥100° F as well as cough and/or sore throat.



Seasonal influenza vaccination:

Seasonal influenza vaccination data in Iowa is based on doses reported to the Iowa Immunization Registry Information System (IRIS). IRIS is a confidential, computerized, population-based system that tracks immunization for children, adolescents and adults who are seen in a variety of public and private healthcare provider sites throughout the state of Iowa. For more information on the immunization data, contact Kim Tichy, IRIS coordinator at 515-281-4288 or kimberly.tichy@idph.iowa.gov.




Note: The data for the 2018-2019 season is only up to the current week and there is a lag between the vaccine administration date and the date reported to the IRIS. The current season's data will be adjusted as additional data is received.

Long-term Care Outbreaks:

Table 6: Number of long-term care outbreaks investigated


REGION	CURRENT WEEK	CUMULATIVE (9/30/18- CURRENT WEEK)
Region 1 (Central)	6	18
Region 2 (NE)	1	2
Region 3 (NW)	0	5
Region 4 (SW)	0	7
Region 5 (SE)	0	7
Region 6 (Eastern)	2	7
Total	9	46

Note: see map in the school section for the counties in each region.



School surveillance program

IDPH monitors illnesses in schools from two different types of reporting: 10 percent school absence reports and weekly sentinel illness reporting. Iowa schools (K-12) track and report (including non-influenza illnesses) when the number of students absent with illness reaches or exceeds 10 percent of total student enrollment. Iowa sentinel schools that participate in IISN voluntarily track and report absence due to all illness and the total enrollment each week. This data provides excellent trends for influenza activity as well as age-specific information used to target vaccination efforts and messages.




^{*}School data not reported for week 52 due to holiday closings

Table 7: Number of schools reporting >10% absenteeism due to any illness

REGION	CURRENT WEEK	CUMULATIVE (9/30/18-CURRENT WEEK)
Region 1 (Central)	12	40
Region 2 (NE)	8	12
Region 3 (NW)	1	25
Region 4 (SW)	4	20
Region 5 (SE)	8	23
Region 6 (Eastern)	22	57
Total	55	177

Note: See map in the school section for the counties in each region. Each school that reports 10 percent illness is counted only once per week for weekly numbers and only once per season for the cumulative reports.

Iowa Influenza Region Map




Non-influenza respiratory viruses:

The State Hygienic Laboratory (SHL) runs a weekly web-based survey program where laboratorians report the number of positive tests for non-influenza respiratory viruses. This table also includes the positive non-influenza virus tests reported from the Dunes Medical Laboratories at Mercy Medical Center in Sioux City. The table includes only the number of positive tests at laboratory surveillance sites throughout the state. The table does not provide case counts.

Table 8: Number of positive results for non-influenza respiratory virus collected by SHL, Mercy Dunes in Sioux City and Iowa Methodist Medical Center

Viruses	CURRENT WEEK	CUMULATIVE (9/30/18–CURRENT WEEK)
Adenovirus	42	902
Parainfluenza Virus Type 1	1	14
Parainfluenza Virus Type 2	1	428
Parainfluenza Virus Type 3	20	201
Parainfluenza Virus Type 4	1	396
Rhinovirus/Enterovirus	139	3473
Respiratory syncytial virus (RSV)	80	1938
Human metapneumovirus (hMPV)	48	356
Coronavirus	129	1770
Total	461	9478



Other resources:

Vaccine:

Influenza vaccine recommendation: idph.iowa.gov/immtb/immunization/vaccine

CDC vaccine information: www.cdc.gov/flu/faq/flu-vaccine-types.htm

Vaccine finder: http://vaccinefinder.org/

Neighboring states' influenza information:

Illinois: www.dph.illinois.gov/topics-services/diseases-and-conditions/influenza/influenza-

surveillance#publications

Minnesota: www.health.state.mn.us/divs/idepc/diseases/flu/stats/index.html

Missouri: health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php

South Dakota: doh.sd.gov/diseases/infectious/flu/

Wisconsin: www.dhs.wisconsin.gov/influenza/index.htm