

# Immunization Schedule with Combination Vaccines

EVERY FALL: FLU VACCINE<sup>4</sup> for anyone 6 months and older

	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	4-6 YEARS
<b>PEDIARIX<sup>®</sup></b> <b>PROQUAD<sup>®</sup></b> <b>QUADRACEL<sup>™</sup></b> or <b>KINRIX<sup>®</sup></b>	<b>PEDIARIX<sup>®</sup></b> DTaP, IPV, HepB <b>+</b> PCV Rotavirus Hib	<b>PEDIARIX<sup>®</sup></b> DTaP, IPV, HepB <sup>1</sup> <b>+</b> PCV Rotavirus Hib	<b>PEDIARIX<sup>®</sup></b> DTaP, IPV, HepB <b>+</b> PCV Rotavirus <sup>2</sup> Hib <sup>3</sup>	PCV Hib HepA MMR <sup>6</sup> Varicella <sup>6</sup>	DTaP	HepA	<b>QUADRACEL<sup>™</sup></b> or <b>KINRIX<sup>®5</sup></b> DTaP, IPV <b>+</b> <b>PROQUAD<sup>®</sup></b> MMRV
<b>PENTACEL<sup>®5</sup></b> <b>PROQUAD<sup>®</sup></b> <b>QUADRACEL<sup>™</sup></b> or <b>KINRIX<sup>®</sup></b>	<b>PENTACEL<sup>®</sup></b> DTaP, IPV, Hib <b>+</b> PCV Rotavirus HepB	<b>PENTACEL<sup>®</sup></b> DTaP, IPV, Hib <b>+</b> PCV Rotavirus HepB <sup>1</sup>	<b>PENTACEL<sup>®</sup></b> DTaP, IPV, Hib <b>+</b> PCV Rotavirus <sup>2</sup> HepB	PCV HepA MMR <sup>6</sup> Varicella <sup>6</sup>	<b>PENTACEL<sup>®</sup></b> DTaP, IPV, Hib	HepA	<b>QUADRACEL<sup>™</sup></b> or <b>KINRIX<sup>®5</sup></b> DTaP, IPV <b>+</b> <b>PROQUAD<sup>®</sup></b> MMRV
<b>PROQUAD<sup>®</sup></b> <b>QUADRACEL<sup>™</sup></b> or <b>KINRIX<sup>®</sup></b>	DTaP IPV HepB Hib PCV Rotavirus	DTaP IPV HepB <sup>1</sup> Hib PCV Rotavirus	DTaP IPV HepB Hib <sup>3</sup> PCV Rotavirus <sup>2</sup>	Hib PCV HepA MMR <sup>6</sup> Varicella <sup>6</sup>	DTaP	HepA	<b>QUADRACEL<sup>™</sup></b> or <b>KINRIX<sup>®5</sup></b> DTaP, IPV <b>+</b> <b>PROQUAD<sup>®</sup></b> MMRV

The above is a suggested schedule using combination vaccines. For alternatives, details and catch-up, refer to the most current *Recommended Immunization Schedules* from CDC.

1. A dose of HepB vaccine is not necessary at 4 months if doses are given at birth and 2 months, but may be included as part of combination vaccine.
2. The 6 month dose is not needed if Rotarix is used exclusively for both dose 1 and 2 in rotavirus vaccine series.
3. The 6 month Hib dose is not needed if PedvaxHIB is used exclusively for the 2<sup>nd</sup> and 4<sup>th</sup> month doses.
4. A child may need 1 or 2 doses of influenza vaccine based on previous vaccination history. Refer to most current influenza vaccine recommendations.
5. ACIP recommends whenever feasible, use the same vaccine manufacturer's DTaP for each dose in the series, however vaccination should not be delayed based on the type of DTaP previously administered.
6. CDC recommends administration of separate MMR and Varicella vaccines for the 12-15 month dose; however MMRV is acceptable based on clinical discretion.

Iowa Department of Public Health Immunization Program - 1-800-831-6293