

**Methodology for Iowa's 2010 Water Quality Assessment,
Listing, and Reporting Pursuant to
Sections 305(b) and 303(d) of the Federal Clean Water Act.**

Prepared by:

Iowa Department of Natural Resources:
Environmental Services Division
Geological and Water Survey
Watershed Monitoring & Assessment Section

April 2011

Table of Contents	Page
Introduction	4
Overview of the assessment and listing process	4
The <i>Iowa Water Quality Standards</i>	5
The total maximum daily load (TMDL)	5
Deadlines	6
The “integrated report”	6
Changes in methodology since the 2006 reporting/listing cycle	13
The assessment and listing process	17
Sources of existing and readily available water quality data	17
Identifying impairments	30
Types of assessments (“evaluated” and “monitored”)	31
Magnitude of impairment	
Data quantity considerations (“data completeness” guidelines)	33
Data quality considerations (“credible data” requirements)	33
Rationale for any decision not to use existing and readily available data	34
List of waters in need of further investigation	40
Overwhelming evidence of impairment	41
How water quality data and other water-related information are summarized to determine whether waters are Section 303(d) “impaired”	42
<ul style="list-style-type: none"> • Physical-chemical data from fixed station monitoring 	42
<ul style="list-style-type: none"> • Data from biological monitoring in support of biocriteria development 	45
<ul style="list-style-type: none"> • Data from IDNR-sponsored lake monitoring by Iowa State University and by SHL 	45
<ul style="list-style-type: none"> • Data from monitoring of bacterial indicators in rivers, lakes, and beach areas 	46
<ul style="list-style-type: none"> • Data from programs to monitor fish tissue for toxic contaminants 	50
<ul style="list-style-type: none"> • Reports of pollutant-caused fish kills 	51
<ul style="list-style-type: none"> • Data from the statewide survey of freshwater mussels from 1998-1999 	54
<ul style="list-style-type: none"> • Data from public water supplies on the quality of raw and finished water 	56
<ul style="list-style-type: none"> • Data from special studies of water quality and aquatic communities 	57
<ul style="list-style-type: none"> • Results of volunteer monitoring that meet “credible data” requirements 	57
Exclusion of waters from the 2006 Section 303(d) list	58
Prioritization and scheduling of waters for TMDL development	60
Addressing interstate inconsistencies in Section 303(d) lists	61
Public participation	62
References	63

Table 1. Summary of changes in Iowa DNR's listing methodology between the 2006 and 2008 listing cycles	66
Table 2. Summary of U.S. EPA's "integrated reporting" format as used for Iowa's 2008 Section 305(b) and Section 303(d) cycle.	66
Table 3. Comparison of Iowa DNR's assessment reaches for the Upper Mississippi River to those agreed upon in 2004 by the Upper Mississippi River Basin Association (UMRBA)	68
Table 4. Monitoring stations on the Upper Mississippi River and its tributaries monitored as part of the USGS Long-Term Resource Monitoring Program (LTRMP)	69
Table 5. Iowa beaches monitored by DNR/SHL or by local cooperators for indicator bacteria during the period 2004-2006.	70
Table 6. Data completeness guidelines for using results of routine ambient water quality monitoring to make assessments of support of beneficial uses	72
Table 7. Summary of Iowa water quality criteria used to make assessments of support of beneficial designated uses of Iowa surface waters	73
Table 8. Summary of Iowa water quality criteria for indicator bacteria (<i>E. coli</i>) in surface waters designated in the <i>Iowa Water Quality Standards</i> (IAC 2003).	76
Table 9. General water quality criteria to protect beneficial general uses for all Iowa surface waters	76
Table 10. Methods for determining support of aquatic life uses	77
Table 11. Methods for determining support of classified, beneficial uses for fish consumption, primary contact recreation, and drinking water uses	78
Table 12. Sample size and number of exceedances to determine an impaired beneficial use.	80
Table 13. Summary of Iowa's revised (2006) protocol for issuing fish consumption advisories.	86
Table 14. Placement of fish kill-affected waters into IR categories for Iowa's 2010 Integrated Reporting cycle.	86
Figure 1. Use of water quality data and information for Iowa's 2008 integrated Section 305(b)/303(d) report/list	87
Attachment 1. Excerpt from Senate File 2371: Iowa's credible data legislation	88
Attachment 2. Guidelines for determining Section 305(b) aquatic life use support using stream biocriteria sampling data	93
Addendum to Attachment 2: Establishment of biological impairment criteria (BIC) for determining support of warmwater stream aquatic life designated uses	105
Attachment 3. The use of the trophic state index to identify water quality impairments in Iowa lakes	113
Attachment 4. Methodology for assessment the degree to which Iowa's shallow natural lakes support their designated aquatic life uses	129
Attachment 5. Iowa DNR interpretations of Section 305(b)/303(d) causes of impairment	139

Introduction

Iowa's 2010 assessment and listing methodology attempts to incorporate recommendations in U.S. EPA's guidance for the 2010 assessment, listing, and reporting requirements pursuant to Sections 303(d) and 305(b) of the federal Clean Water Act (U.S. EPA 2005, 2006, 2009). This EPA guidance establishes the formats for an "integrated report" (IR) that satisfies the listing requirements of Section 303(d) and the reporting requirements of Sections 305(b) and 314 of the Clean Water Act (CWA). This EPA guidance replaces all previous guidance pertaining to Sections 305(b) and 303(d) except EPA's Consolidated Assessment and Listing Methodology (CALM) (U.S. EPA 2002). Due to the lack of details regarding the mechanics of water quality assessment in more recent EPA guidance (e.g., U.S. EPA 2002), IDNR continues to use assessment methods recommended in previous EPA guidance for Section 305(b) reporting (U.S. EPA 1997). Iowa's 2010 methodology meets the requirements of CWA, Section 303(d)(1)(a) and 40 CFR Section 130.24 and incorporates requirements of Iowa's credible data law (Attachment 1). The changes in methodology between the 2008 and 2010 listing cycles are summarized in Table 1 and are explained throughout this document.

Overview of the assessment and listing process:

The process of assessing water quality and adding waterbodies to the state list of "impaired" waters involves three interrelated program areas of the federal Clean Water Act (CWA): (1) establishment of state water quality standards that identify beneficial uses for the state's waterbodies and that identify criteria to determine whether each use is being achieved, (2) development of water quality assessments by comparing water quality information to water quality standards to determine whether or not beneficial uses are being achieved, and (3) addition of the appropriate waters assessed as "not fully supporting" beneficial uses (i.e., "impaired") to the state's Section 303(d) list. The state's 303(d) list is thus a public accounting of all assessed waterbodies determined to be impaired where a TMDL needs to be developed. A waterbody that is placed on the 303(d) list has been assessed as "not fully meeting" water quality standards including designated uses (e.g., for primary contact recreation, aquatic life, and/or as a source of drinking water for a public water supply). The failure to fully meet state standards can result from the following: violations of numeric criteria, narrative criteria, anti-degradation requirements as defined in U.S. EPA's regulations regarding water quality standards (40 CFR 131), and/or a determination that a specific designated use cannot be achieved. The violations of water quality standards might be due to an individual pollutant, multiple pollutants, or an unknown cause of impairment. As provided for in U.S. EPA's (2005, 2006, 2009) guidance for integrated reporting, other waterbodies may be assessed as impaired but not included on the 303(d) list. These waters will be included in Category 4 of the Integrated Report (Table 1). IR Category 4 includes three types of impaired

waterbodies: (1) waters for which a TMDL has been completed but water quality standards have not yet been attained; (2) waters where other required control measures are expected to result in attainment of water quality standards in a reasonable period of time; and (3) the impairment or threat is not caused by a "pollutant" as defined by U.S. EPA. In addition, Iowa waters assessed as impaired by pollutant-caused fish kills are placed in Category 4 if an IDNR investigation identified the person responsible for the kill and monetary restitution for the fish killed has been sought.

The Iowa Water Quality Standards:

According to U.S. EPA, a water quality standard is composed of three components: (1) a description of beneficial use, (2) water quality criteria to protect this use, and (3) an anti-degradation policy that ensures protection of water quality where water quality exceeds levels necessary to protect fish and wildlife propagation and recreation in and on the water. Thus, the basis for a state's 305(b) assessments and Section 303(d) lists of impaired waters is ultimately the state's water quality standards. That is, the state water quality standards contain the benchmarks (criteria) to which water quality data are compared to determine the degree to which beneficial uses are supported. The versions of the *Iowa Water Quality Standards* and the accompanying *Surface Water Classification* with the effective date of June 11, 2008 were used as the basis for water quality assessments prepared for this (2010) assessment and listing cycle. This version of the *Standards* was the most recent EPA-approved version available during the period of time covered by the 2010 assessment and listing cycle (2006 through 2008). These versions of the standards and surface water classification are available upon request from Iowa DNR's Watershed Monitoring & Assessment Section.

The Total Maximum Daily Load (TMDL):

The Watershed Monitoring & Assessment Section of the Iowa DNR's Geological and Water Survey conducts water quality assessments as required by Clean Water Act Section 305(b). Based on these assessments, section staff identify waterbodies in the state of Iowa that may require a total maximum daily load (TMDL) allocation to address the causes and sources of pollutants contributing to impairment of a designated use or other applicable beneficial use (IR Category 5 waters). In general terms, a TMDL defines the level of water quality needed to support a water quality standard, including the designated uses, water quality criteria, and the anti-degradation policy that comprise the standard. Conceptually, a TMDL is the maximum pollutant load from point sources and nonpoint sources, plus a load allocated to a "margin of safety," that a waterbody can receive and continue to meet water quality standards. The margin of safety accounts for the lack of understanding of the relationship between pollutant loads and water quality.

Deadlines:

According to current EPA regulations, the Section 303(d) list of impaired waterbodies must be submitted to EPA by April 1 of every even numbered year. Thus, this methodology was designed to meet the deadline for submission of the list to be submitted to U.S. EPA in April 2010.

The “integrated report”:

Based on previous guidance from U.S. EPA (e.g., U.S. EPA 1997), most states, including Iowa, had historically produced separate Section 305(b) reports and Section 303(d) lists. Section 305(b) reports have attempted to characterize water quality statewide and thus identified not only designated use impairments but also water quality concerns that are worthy of note and further investigation but do not constitute Section 303(d)-type water quality impairments. The 303(d) lists, on the other hand, have represented the subset of waterbodies assessed for the Section 305(b) report with known and reasonably verifiable impairments of a designated use or general use, as defined in the *Iowa Water Quality Standards* that are appropriate for Section 303(d) listing. Based on development of revised guidance by U.S. EPA (2003), however, an “integrated report” was prepared for the 2004 cycle that incorporated elements of both the Section 305(b) report and Section 303(d) list. Based on updated guidance from U.S. EPA (2005, 2006, 2009), IDNR has continued to use the integrated report format for the 2006, 2008, and the current (2010) Section 305(b) reporting and Section 303(d) listing cycles.

In their guidance for the integrated assessment, reporting, and listing cycles, U.S. EPA (2003, 2005, 2006) recommended that reporting requirements of Sections 305(b) and 303(d) be “integrated” into a report that contains five assessment categories and associated subcategories:

- Category 1: All designated uses are met.
- Category 2: Some of the designated uses are met but there is insufficient data to determine if remaining designated uses are met.
- Category 3: Insufficient data exist to determine whether any designated uses are met.
- Category 4: Water is impaired or threatened but a TMDL is not needed because one of the following occur:
 - 4a. A TMDL has been completed;
 - 4b. Other required control measures are expected to result in attainment of water quality standards in a reasonable period of time;

4c: The impairment or threat is not caused by a “pollutant.”

- Category 5: Water is impaired or threatened and a TMDL is needed [IR Category 5 is the state's Section 303(d) list].

The five categories of EPA's integrated reporting and listing format used for Iowa's integrated reports since the 2004 reporting cycle are further explained below and are summarized in Table 2. In the descriptions below, the text in italics is taken directly from U.S. EPA's (2005) guidance for integrated reporting. The notes that follow these excerpts contain IDNR's interpretations and modifications of this guidance.

Category 1 waterbodies: *Waters belong in Category 1 if they are attaining all designated uses and no use is threatened. Segments should be listed in this category if there are data and information that are consistent with the State's methodology and this guidance, and support a determination that all WQSs [water quality standards] are attained and no designated use is threatened.*

Iowa DNR has made no modifications to the definition or intent of IR Category 1.

Category 2 waterbodies: *Waters should be placed in Category 2 if there are data and information that meet the requirements of the State's assessment and listing methodology that support a determination that some, but not all, designated uses are attained and none are threatened. Attainment status of the remaining designated uses is unknown because data are insufficient to categorize a water consistent with the State's listing methodology.*

Iowa DNR has made the following modifications to IR Category 2: the renaming of EPA's Category 2 as Category 2a and the addition of Category 2b.

Category 2a: Some uses supported; insufficient information to determine whether other uses are supported. This wording is consistent with U.S. EPA's definition of IR Category 2.

Category 2b: At least one use assessed as fully supported with at least one other use “evaluated” as impaired. An “evaluated” assessment of impairment lacks sufficient confidence to take forward to either Category 5 (Section 303(d) list) or

Category 4 (impaired but TMDL not required). This subcategory allows tracking of the “impaired / evaluated” waterbodies (e.g., a biological assessment of impairment based on data generated by a non-IDNR sampling protocol). Waters placed into subcategory 2b will be added to Iowa’s list of “waters in need of further investigation.”

As part of revisions to its biological assessment protocol for the 2010 Integrated Reporting cycle, IDNR has added the following subcategories to IR subcategory 2b to improve IDNR’s ability to better target follow-up monitoring on streams and rivers where potential biological impairments have been identified. That is, these subcategories have been added to allow IDNR to track potentially impaired streams and rivers that are (1) within the calibration watershed size of Iowa’s biological assessment protocol (watersheds from ~ 10 to 500 square miles) and (2) that are outside (i.e., watersheds too small or too large) this calibration range. The following subcategories have been added for the 2010 cycle:

2b-c [calibrated]: At least one use (contact recreation, drinking water, or fish consumption) is assessed as “fully supported,” but the aquatic life use of a stream segment with a watershed size within the calibrated range of the biological assessment protocol has been assessed as potentially impaired;

2b-u [un-calibrated]: At least one use (contact recreation, drinking water, or fish consumption) is assessed as “fully supported,” but the aquatic life use of a stream segment with a watershed size outside the calibrated range of the biological assessment protocol has been assessed as potentially impaired;

Category 3 waterbodies: *Waters belong in Category 3 if there are insufficient or no data and information to determine, consistent with the State's listing methodology, if any designated use is attained. To assess the attainment status of these waters, States should schedule monitoring on a priority basis to obtain data and should also make efforts obtain information necessary to move these waters into Categories 1, 2, 4, and 5.*

Iowa DNR has made the following modifications to IR Category 3: the renaming of EPA Category 3 to Category 3a and the addition of Category 3b.

Category 3a: Insufficient data exist to determine whether any uses are met; no uses are assessed [either “evaluated” or “monitored”]. This wording is consistent with U.S. EPA’s definition of IR Category 3.

Category 3b: Insufficient data exist to determine whether any designated uses are met, but at least one use is assessed as potentially impaired based on an “evaluated” assessment. This category is similar to IDNR’s Category 2b, but no other uses are assessed as “fully supported” (i.e., the only use assessed is the one assessed as “impaired/evaluated”). Similar to IDNR subcategory 2b, this subcategory allows tracking of the “impaired / evaluated” waterbodies. Waters placed into subcategory 3b will be added to Iowa’s list of “waters in need of further investigation.”

As part of revisions to its biological assessment protocol for the 2010 Integrated Reporting cycle, IDNR has added the following subcategories to IR subcategory 3b to improve IDNR’s ability to better target follow-up monitoring on streams and rivers where potential biological impairments have been identified. That is, these subcategories have been added to allow IDNR to track potentially impaired streams and rivers that are (1) within the calibration watershed size of Iowa’s biological assessment protocol (watersheds from ~ 10 to 500 square miles) and (2) that are outside (i.e., watersheds too small or too large) this calibration range. The following subcategories have been added for the 2010 cycle:

3b-c [calibrated]: the aquatic life use of a stream segment within the calibrated range of the biological assessment protocol has been assessed as potentially impaired; no other uses are assessed due to lack of water quality information;

3b-u [un-calibrated]: the aquatic life use of a stream segment with a watershed size outside the calibrated range of the biological assessment protocol has been assessed as potentially impaired; no other uses are assessed due to lack of water quality information;

Category 4 waterbodies: *Waters belong in Category 4 if one or more designated uses are impaired or threatened but establishment of a TMDL is not required. States may place an impaired or threatened water that does not require a TMDL in one of the following three subcategories:*

- Category 4a: a TMDL has been completed for the water-pollutant combination. Waters should only be placed in Category 4a when all TMDLs needed to result in attainment of all applicable WQ Standards have been approved or established by EPA. Current regulations do not require TMDLs for all waters.
- Category 4b: other required control measures are expected to result in the attainment of WQSs in a reasonable period of time. Some waters may be excluded from Category 5, and placed into Category 4b. In order to meet the requirements to place these waters into Category 4b, the State must demonstrate that "other pollution control requirements (e.g., best management practices) required by local, State or Federal authority" (see 40 CFR 130.7(b)(1)(iii)) are expected to address all water-pollutant combinations and attain all WQ Standards in a reasonable period of time. EPA expects that States will provide adequate documentation that the required control mechanisms will address all major pollutant sources and establish a clear link between the control mechanisms and WQ Standards.
- Category 4c: the impairment or threat is not caused by a pollutant. Waters should be listed in Category 4c when an impairment is not caused by a pollutant. "Pollution," as defined by the Clean Water Act, is the "man-made or man-induced alteration of the chemical, physical, biological and radiological integrity of water." In some cases, the pollution is caused by the presence of a pollutant and a TMDL is required. In other cases, pollution does not result from a pollutant and a TMDL is not required. An example of a pollutant stressor would be copper; an example of a non-pollutant stressor ("pollution") would be "low flow."

Iowa DNR has made no modifications to the definitions or intents of IR Categories 4a, 4b, or 4c. Iowa DNR has, however, made the following modification to IR Category 4: the addition of Category 4d.

Category 4d: Water is impaired due to a pollutant-caused fish kill and enforcement actions were taken against the party responsible for the kill: a TMDL is neither appropriate nor needed. For purposes of Section 305(b) assessments in Iowa, all waters affected by a fish kill caused by a known pollutant or a suspected pollutant are assessed as impaired. Those kills where a pollutant cause was identified are placed into either Category 4d (responsible party identified and enforcement action taken: TMDL not required) or Category 5 (no responsible party identified; enforcement action not taken: a pollutant problem may remain and a TMDL is potentially needed).

Category 5 waterbodies: *This category constitutes the Section 303(d) list that EPA will approve or disapprove under the CWA. Waters should be placed in Category 5 when it is determined, in accordance with the State's assessment and listing methodology, that a pollutant has caused, is suspected of causing, or is projected to cause an impairment or threat. If that impairment or threat is due to a pollutant, the water should be placed in Category 5 and the pollutant causing the impairment identified.*

Iowa DNR has made the following modifications to IR Category 5: the renaming of EPA's Category 5 to Category 5a and the addition of categories 5b and 5p.

Category 5a: Water is impaired or threatened by a pollutant stressor and a TMDL is needed. This wording is consistent with U.S. EPA's definition of IR Category 5.

Category 5b: Impairment is based on results of biological monitoring or a fish kill investigation where specific causes and/or sources of the impairment have not yet been identified. The biological assessment adequately demonstrates that an impairment exists, but either the cause or the source of the impairment is unknown. The primary use of this subcategory is for biologically-based (biomonitoring) impairments with the cause listed as "unknown" and for fish kill-based impairments where a pollutant cause was identified but no source was found. Additional monitoring / investigation, such as that conducted as part of IDNR's stressor identification procedure, is needed to determine causes or sources before the TMDL can be developed.

As part of revisions to its biological assessment protocol for the 2010 Integrated Reporting cycle, IDNR has added the following subcategories to IR Subcategory 5b to improve IDNR's ability to track the impairment status of streams and rivers and to better target follow-up monitoring where both biological impairments and potential de-listings have been identified.

5b-t [tentative]: The aquatic life uses of a stream segment with a watershed size within the calibration range of the IDNR biological assessment protocol (~10 to 500 square miles) are assessed as Section 303(d)-impaired based on an evaluated assessment. The reasons for

residency in this subcategory include: 1) data quantity (single sample assessments), 2) data age (data older than five years), 3) data quality (marginal sampling conditions), and 4) sampling frequency (multiple samples collected in same year, not multiple years).

5b-v [verified]: The aquatic life uses of a stream with a watershed size within the calibration range of IDNR biological assessment protocol (~10 to 500 square miles) are assessed as Section 303(d)-impaired based on results of the required two or more biological sampling events in multiple years within the previous five years needed to confirm the existence of a biological impairment.

Category 5p: Impairment occurs on a waterbody presumptively designated for Class A1 primary contact recreation use or Class B(WW1) aquatic life use. Due to changes in the *Iowa Water Quality Standards* that became effective in March 2006, all perennially-flowing streams and intermittent streams with perennial pools are presumed to be capable to supporting the highest level of primary contact recreation use (Class A1) and the highest level of aquatic life use [Class B(WW1)]. These changes to the *Iowa Water Quality Standards* were approved by U.S. EPA in February 2008. Under this approach to stream classification, the Class A1 (primary contact recreation) use is applied to all of Iowa's perennial rivers and streams and intermittent streams with perennial pools, and the Class B(WW1) aquatic life use is applied to all of Iowa's perennial rivers and streams and intermittent streams with perennial pools unless the water is already designated for Class B(WW2) or Class B(WW3) uses in Iowa's surface water classification. A "use attainability analysis" or UAA must be conducted, including field investigations, to determine whether a presumptively-applied use is, in fact, the appropriate designated use for the stream segment in question. Until the time when a UAA has been conducted and the appropriate designated uses have been applied and approved, any impairments on presumptively-designated Iowa streams will be placed in IR Category 5p.

According to U.S. EPA's (2005) guidance, the Section 303(d) list is composed of waters included in IR Category 5 of the Integrated Report which includes those waters for which a TMDL needs to be developed. This list includes waterbodies impaired by "pollutants" such as nitrate and

indicator bacteria. The source of impairment might be from point sources, nonpoint sources, groundwater or atmospheric deposition. Some sources of impairment of Iowa waterbodies originate outside of the state. Historically, Iowa has listed impaired waterbodies regardless of whether the source of pollutant is known and regardless of whether the pollutant source(s) can be legally controlled or acted upon by the state of Iowa. This methodology is consistent with that history.

As specified in Iowa's credible data law, waterbodies where the assessment indicates a potential impairment, but where sufficient and credible data are lacking, will not be included on the state's 303(d) list (IR Category 5). According to this methodology, these waters will be included in IR subcategories 2b or 3b and placed on the state list of "waters in need of further investigation" as provided for by Iowa's credible data legislation.

Changes in methodology since the 2008 reporting/listing cycle

The changes in IDNR's assessment and listing methodology between the 2008 and current (2010) cycles are summarized in Table 1. The following changes were made.

(1) Assigning magnitudes of bacterial impairment:

Despite U.S. EPA's tendency to have states move away from reporting magnitude of impairment as part of Integrated Reporting, DNR has continued to use the "partially supporting" assessment category to indicate an impairment of moderate magnitude and the "not supporting" category to indicate a relatively severe impairment. IDNR feels that ranking impairments according to magnitude is useful, especially regarding the need to prioritize impairments for TMDL development. Prior to the 2010 IR cycle, impairments due only to single-sample maximum criteria violations were considered to be of a "moderate" impairment magnitude while impairments due to violations of the geometric mean criteria were considered a "severe" magnitude. For the 2010 IR cycle, the identification of impairment magnitudes has been revised and will now also be based only on the distribution of geometric means from the 2008 IR cycle. That is, an impairment due to a geometric mean of less than or equal to 1,000 organisms/100 ml is considered of a "moderate" magnitude (use is "partially supported"), while a geometric mean of greater than 1,000 organisms/100 ml will be considered to be a "severe" impairment (use is "not supported"). This revised framework is summarized in Table 11.

(2) Revision of IDNR's biological assessment protocol:

IDNR has historically based biological impairment decisions on only one biological sampling event that incorporated IDNR's biological assessment protocol or an equivalent protocol. Problems with impairment flip-flopping, however, have occurred on streams where impairment magnitudes have been slight. That is, instances have occurred where a watershed improvement project has been initiated on a stream segment with a slight biological impairment based on only one sampling event. Results of additional biological monitoring conducted as part of the watershed project, however, have sometimes shown "full support" of aquatic life uses, thus leaving the just-initiated watershed improvement project with no impairment to correct.

In order to minimize the chance of impairment flip-flopping, IDNR's biological assessment protocol was revised for the 2010 Integrated Reporting cycle. Results from two independent biological sampling events from different years and within a five-year period that both suggest impairment of the aquatic life uses of a stream segment are now required before a biologically-based Section 303(d) impairment is identified.

(3) Addition of Integrated Report subcategories related to biological assessments of streams and rivers:

As part of revisions to its biological assessment protocol for the 2010 Integrated Reporting cycle, IDNR has added subcategories to IR Categories 2, 3, and 5 to improve IDNR's ability to track the impairment status of streams and rivers and to better target follow-up monitoring where potential biological impairments have been identified. Subcategories for biologically-assessed waters have been added to IR Categories 2 and 3 to allow IDNR to track potentially impaired streams and rivers that are (1) within the calibration watershed size of Iowa's biological assessment protocol (watersheds from 10 to 500 square miles) and (2) that are outside (i.e., watersheds too small or too large) this calibration range. The following subcategories have been added for the 2010 cycle:

2b-c [calibrated]: At least one other use (contact recreation, drinking water, or fish consumption) is assessed as "fully supported," but the aquatic life use of a stream segment with a watershed size within the calibrated range of the biological assessment protocol has been assessed as potentially impaired;

2b-u [un-calibrated]: At least one other use (contact recreation, drinking water, or fish consumption) is assessed as "fully supported," but the aquatic life use of a stream

segment with a watershed size outside the calibrated range of the biological assessment protocol has been assessed as potentially impaired;

3b-c [calibrated]: the aquatic life use of a stream segment within the calibrated range of the biological assessment protocol has been assessed as potentially impaired; no other uses are assessed due to lack of water quality information;

3b-u [un-calibrated]: the aquatic life use of a stream segment with a watershed size outside the calibrated range of the biological assessment protocol has been assessed as potentially impaired; no other uses are assessed due to lack of water quality information;

5b-t [tentative]: The aquatic life uses of a stream segment with a watershed size within the calibration range of the IDNR biological assessment protocol are assessed as Section 303(d)-impaired based on only one of the two biological sampling events needed to confirm the existence of a biological impairment.

5b-v [verified]: The aquatic life uses of a stream with a watershed size within the calibration range of IDNR biological assessment protocol are assessed as Section 303(d)-impaired based on results of the required two or more biological sampling events in multiple years within the previous five years needed to confirm the existence of a biological impairment.

These subcategories have been incorporated into IDNR's Section 305(b) assessment database, ADBNet (<http://programs.iowadnr.gov/adbnet/index.aspx>).

(5) Changes in the Iowa Water Quality Standards since the 2008 reporting cycle:

Revised water quality criteria in the *Iowa Water Quality Standards* were approved by U.S. EPA for the following toxic parameters in 2009:

- aluminum
- arsenic
- cadmium
- chromium
- copper
- cyanide
- DDT+DDD+DDE
- dieldrin (human health only)
- lead

- lindane (human health only)
- mercury
- zinc

These criteria changes have been incorporated into the assessment and listing process for the 2010 Integrated Reporting cycle.

The Assessment and Listing Process

Preparation of Iowa's integrated [305(b)/303(d)] report includes the following basic steps:

- Assemble all existing and readily available water quality-related data and information not previously used for 305(b) water quality assessments;
- Identify water quality-related data and information of sufficient quality and quantity for purposes of developing scientifically defensible water quality assessments;
- Compare these water quality-related data and information to state water quality standards to determine the degree to which assessed waters meet these standards;
- Identify Section 303(d) impairments that are based on water quality-related data and information that meet the state's requirements for data quantity and data quality (Table 6);
- Place all waters into one of the five categories specified in U.S. EPA's (2003, 2005) "integrated report" guidance for water quality assessment and listing;
- Prepare the state list of waters in need of further investigation as required by state law;
- Prioritize the waterbodies on the draft Section 303(d) list (Category 5) for TMDL development (high, medium, and low);
- Provide the draft integrated report, including the draft Section 303(d) list (Category 5), to the public for review and comment;
- Revise and finalize the integrated report based on new information and public input;
- Submit the finalized integrated report, including the Section 303(d) list, to U.S. EPA for approval/disapproval;
- Develop a schedule for development of TMDLs for Section 303(d)-listed (IR Category 5) waterbodies.

Sources of existing and readily available water quality-related data and information:

As specified in U.S. EPA's current (1992) TMDL rule (40 CFR 130.7), sources of existing and readily available water quality-related data and information to be considered as part of Section 303(d) listing include, but are not limited to, the following:

- the state's most recent CWA Section 305(b) assessments;
- CWA Section 319 nonpoint source assessments;
- dilution calculations, trend analyses, or predictive models for determining the physical, chemical, or biological integrity of streams, rivers, lakes, and estuaries;
- water quality-related data and water-related information from local, State, Territorial, or Federal agencies (especially the U.S. Geological Survey's National Water Quality Assessment Program (NAWQA) and National Stream Quality Accounting Network (NASQAN)), Tribal governments, members of the public, and academic institutions.

Historically, the majority of information used by IDNR to develop Iowa's Section 303(d) list of impaired waters has been taken from its Section 305(b) assessments. Data sources used to assess water quality conditions in Iowa for purposes of Section 305(b) assessment include, but are not limited to, the following:

- Physical, chemical, and biological data from ambient fixed station water quality monitoring networks conducted by IDNR and other agencies (e.g., U.S. Geological Survey; U.S. Army Corps of Engineers);
- Data from water quality monitoring conducted by adjacent states on border rivers and waters flowing into the state;
- Data from biological monitoring being conducted by IDNR in cooperation with the State Hygienic Laboratory at The University of Iowa (SHL) as part of a current effort to establish biological criteria for Iowa's ecoregions and subcoregions and as part of other projects (e.g., the 2000-2005 regional Environmental Monitoring and Assessment Program (REMAP) project);
- Data from the IDNR-sponsored statewide lake monitoring project conducted by Iowa State University and SHL;
- Data from monitoring of bacterial indicators in rivers and at beaches of publicly-owned lakes;
- Data from programs to monitor fish tissue for toxic contaminants;
- Reports of pollutant-caused fish kills;
- Where readily available, data from public water supplies on the quality of raw and finished water;
- Drinking water-related source water assessments under Section 1453 of the Safe Drinking Water Act;
- Data from special studies of water quality and aquatic communities;
- Best professional judgment of IDNR staff;

- Results of volunteer monitoring (e.g., by IOWATER-trained volunteers);
- Water-related information received from the public.

The cutoff date for the data collection period for Iowa's 2010 Integrated Report is the end of the calendar year 2008. This is a general guideline used by IDNR, and more recent information may be used for some types of water quality information that becomes available infrequently or at irregular intervals (e.g., fish consumption advisories and reports of pollution-caused fish kills). Large amounts of staff time are needed to summarize monitoring data, compare the summarized results to water quality standards, develop the waterbody-specific assessments of the degree to which designated uses are supported, and to solicit and respond to public comments on the draft Section 303(d) list. Also, water quality data generated by the various agencies are not available immediately following sample collection: a lag time of from a few months up to half a year is associated with obtaining results of water quality monitoring networks. Given these time requirements, and given the other work responsibilities of IDNR staff that prepare Iowa's Integrated Report, the allowance of a 15-month window for report preparation prior to the April deadline is not excessive.

For purposes of developing water quality assessments for the 2010 integrated report, three years of water quality data from streams and rivers were typically used for both conventional pollutant parameters (e.g., indicator bacteria) and the less frequently monitored toxic parameters (e.g., toxic metals). This is the fourth consecutive 305(b)/303(d) cycle for which IDNR has used a three-year data gathering period. Prior to the 2004 cycle, only two years of data were used for Iowa's Section 305(b) reports. For most assessments, the use of three years of data increases the number of samples upon which the decision on use support is based and will help address the problem of year-to-year fluctuations in water quality. More recent data and information are used where appropriate to supplement the current assessment. Older data, up to five years old (data from 2004 through 2008 for the 2010 Integrated Report cycle), are used to supplement data from the current assessment period for water quality parameters with low collection frequency (e.g., toxic metals). Due to the lower sampling frequency in Iowa's ambient lake monitoring programs, five years of data (2004-2008) are used for developing Section 305(b) assessments and for identifying Section 303(d) listings for Iowa lakes.

As specified in Iowa's credible data law, and based on the uncertainty inherent in using old data to characterize current water quality conditions, data between five and ten years old are used for Section 305(b) assessments but are not used for purposes of adding waters to Iowa's Section 303(d) list of impaired waters (i.e., Category Five of the Integrated Report). Chemical/physical data older than five years are generally believed to be less reflective of current ambient water quality than are more recent

data (U.S EPA 1997, pages 1-5 and 1-9). Of course, nearly all recent water quality data from Iowa waters have already been used for Section 305(b) assessments and thus have already been considered for Section 303(d) listings. Also, a listed waterbody will not be removed from the state's Section 303(d) list simply because the data upon which the impairment was based have aged beyond five years. Thus, the restrictions placed on use of old water quality data have little effect on impaired waters listings or de-listings in Iowa.

The sources of water quality data used for water quality assessments and impaired waters listings in Iowa are discussed in more detail below.

- **Physical, chemical, and biological data from ambient fixed station water quality monitoring networks conducted in Iowa by IDNR and other agencies**

IDNR, in cooperation with SHL, has conducted routine ambient monitoring of river water quality in Iowa since the early 1980s. Due to resource constraints, the majority of this monitoring prior to 1999 was limited to relatively few (16) locations. An appropriation from the Iowa Legislature, however, allowed a significant expansion of this monitoring program beginning in October 1999. Iowa rivers are now monitored monthly at approximately 85 sites for a variety of physical, chemical, and bacterial parameters through a contract with the SHL which provides both data collection and laboratory services. Sixty-two of these sites are classified as ambient (background) sites. These sites are distributed throughout every major river basin in an effort to provide good geographic coverage of the state. Twenty-three of the sites are associated with 10 major cities, with monitoring stations located both upstream and downstream from each city. In addition to the standard parameters, the upstream/downstream urban sites are being tested for a variety of industrial chemicals and insecticides. For more information on the IDNR's ambient and city monitoring programs see the following web site: <http://www.igsb.uiowa.edu/wqm/>.

Long-term ambient water-quality monitoring has also been conducted in Iowa by the following agencies: U.S. Army Corps of Engineers, U.S. Geological Survey (USGS), and water utilities such as the Des Moines Water Works, the Cedar Rapids Water Department, and the Rathbun Rural Water Association. The monitoring networks in Iowa conducted by agencies other than IDNR are typically designed to answer questions specific to drinking water sources or to the effects of in-stream structures or large facilities on water quality (e.g., flood control reservoirs or power generating facilities). For example, networks have been established by the U.S. Army Corps of Engineers on the Des Moines, Raccoon, and

Iowa rivers to evaluate changes in water quality caused by Saylorville, Red Rock, and Coralville reservoirs (see Lutz and Francois 2007, 2008, and Lutz and Steffen 2009). In general, stations in these networks have remained fixed for approximately four decades, and they have been monitored more frequently than stations in the IDNR/SHL network. Thus, these networks provide a relatively long-term database that can be used to characterize water quality conditions. For information on the monitoring networks on the Des Moines and Raccoon rivers, see the following web site:

<http://home.eng.iastate.edu/~dslutz/dmrwqn/dmrwqn.html>.

Currently, USGS conducts routine water quality monitoring at three fixed stations in Iowa: the Mississippi River at Clinton, the Missouri River at Omaha (Council Bluffs), and the Big Sioux River at Akron. All three of these sites are remnants of the USGS National Stream Quality Accounting Network (NASQAN). In late 1994, the USGS began routine monitoring at selected locations in the Skunk, Iowa, Cedar, and Wapsipinicon river basins as part of the National Water Quality Assessment Program (NAWQA) in the Eastern Iowa Basins study unit. This monitoring was conducted through September 1998. The NAWQA program was designed to generate comprehensive and nationally-consistent water quality information that could be used to describe the status and trends of the nation's water resources. During the 2006-2008 data gathering period for the current (2010) Integrated Reporting cycle, the following streams were routinely monitored by USGS such that 10 or more samples were collected over the three-year period:

	USGS Monitoring Station:	Site Number
1.	Boyer River at Logan, Harrison Co.	06609500
2.	Cedar River at Edgewood Road, Cedar Rapids, Linn Co.	05464480
3.	Des Moines River at Keosauqua, Van Buren Co.	05490500
4.	Iowa River at Iowa City, Johnson Co.	05454500
5.	Iowa River at Wapello, Louisa Co.	05465500
6.	Little Sioux River at Turin, Monona Co.	06607500
7.	Maquoketa River at Spragueville, Jackson Co.	05418600
8.	Nishnabotna River at Hamburg, Fremont Co.	06810000
9.	Skunk River at Augusta, Lee Co.	05474000
10.	South Fork Iowa River headwaters near Blairsburg, Hamilton Co.	04541070

	USGS Monitoring Station:	Site Number
11.	South Fork Iowa River NE of New Providence, Hardin Co.	05451210
12.	South Fork Iowa River near Blairsburg, Hamilton Co.	05451080
13.	Turkey River at Garber, Clayton Co.	05412500
14.	Wapsipinicon River near DeWitt, Clinton Co.	05422000
15.	Wapsipinicon River near Tripoli, Bremer Co.	05420680

Data from USGS monitoring in Iowa are available at the following web site:

<http://waterdata.usgs.gov/nwis/sw>.

- **Data for Iowa tributaries of the Upper Mississippi River generated by the Long Term Resource Monitoring Program**

Intensive water quality monitoring of Pool 13 of the Upper Mississippi River and several Iowa tributaries is conducted by Iowa DNR staff at Bellevue, Iowa, as part of the Long-Term Resource Monitoring Program (LTRMP). Staff from other LTRMP field stations (e.g., Onalaska, WI) also sample Iowa tributaries. The LTRMP was authorized under the Water Resources Development Act of 1986 as an element of the U.S. Army Corps of Engineers "Environmental Management Program" (EMP) and is currently being implemented by the U.S. Geological Survey in cooperation with the five Upper Mississippi River basin states (Illinois, Iowa, Minnesota, Missouri, and Wisconsin). State staff at six field stations in the Upper Mississippi River system conduct monitoring of fisheries, macroinvertebrates, vegetation, as well as water quality on specified reaches of the river. Water quality monitoring by the LTRMP began in 1988 and continues. LTRMP stations with chemical data used for Section 305(b) water quality assessments and Section 303(d) listings in Iowa are summarized in Table 4. Data from this network are available from the Upper Midwest Environmental Sciences Center (see

<http://www.umesc.usgs.gov/ltrmp.html>).

- **Data from water quality monitoring conducted by adjacent states on border rivers and waters flowing into the state**

States adjacent to Iowa (South Dakota, Minnesota, Wisconsin, Illinois, Missouri, and Nebraska) also have fixed station ambient water quality monitoring programs that generate data useful for purposes of water quality assessments in Iowa. Data from these monitoring networks are available either through the U.S. EPA's national water quality

database "STORET and WQX" [<http://www.epa.gov/storet/>] or through personal contacts with water quality monitoring staff of environmental agencies in these states. These data are used with the guidelines described in this document to assess the degree to which the relevant *Iowa Water Quality Standards* are being met. In addition, decisions on assessment and listing for interstate waters are coordinated to the extent possible with water quality staff from the adjacent states. For example, assessments and listings for the Iowa portion of the Upper Mississippi River are made in consultation with the states of Minnesota, Wisconsin, Illinois, and Missouri as part of ongoing interstate 305(b)/303(d) consultations through the Upper Mississippi River Basin Association's *Water Quality Task Force* (<http://www.umrba.org/wq.htm>). UMRBA consultations and coordination of assessments and listings are based on a uniform set of assessment reaches for the Upper Mississippi River that was adopted by all five UMR states in 2004 (Table 3).

- **Data from biological monitoring being conducted by IDNR in cooperation with SHL as part of a current effort to establish biological criteria for Iowa's ecoregions and subcoregions and as part of the 2002-2005 Regional EMAP project**

Biological criteria or "biocriteria" are narrative or numeric expressions that describe the best attainable biological integrity (reference condition) of aquatic communities inhabiting waters of a given designated aquatic life use. In order to develop biocriteria, knowledge of the variation in the ecological and biological conditions within a state is necessary. Ecoregions, generally defined as regions of relative homogeneity in ecological systems and relationships between organisms and their environments, have been used by several states when developing biocriteria for their water quality standards. Biological reference sites are located on the least impacted streams within an ecoregion. Monitoring results from regional reference sites can thus serve as benchmarks to which other streams in the region can be compared.

In Iowa, a list of candidate stream reference sites was generated in the early 1990s for the state's ten ecoregions and subcoregions (see Attachment 2). Sampling of reference sites began in 1994 and continues; the current rate of sampling is 20 sites per year. Stream biological sampling is conducted from July 15 to October 15. In addition to reference site sampling, sampling at "test" sites is conducted to determine how much a stream's biological health is impacted by disturbances such as channelization, livestock grazing, manure spills, wastewater discharges and urban runoff. Currently, approximately 40 test sites are sampled per year. At both reference sites and test sites, standard

sampling procedures are used so that data from all sites are comparable. The samples measure how many types of benthic macroinvertebrates and fish are present and the abundance of each type in relation to the whole sample. Benthic macroinvertebrates are collected from several types of habitat including aquatic vegetation, boulders, leaf packs, overhanging vegetation, rocks, root mats and woody debris. Fish are sampled in one pass through the sampling area using electrofishing gear. These bioassessment sampling protocols have also been used to examine the location and amount of biological impairment in TMDL-targeted watersheds (see IDNR/WRS 2001). More watershed sampling is planned to support development of stream restoration plans, including TMDLs. The data from the sampling of reference sites, test sites, and watershed sites are being used to develop indicators of stream biological integrity that will form the basis for establishment of numeric biocriteria that will be used for assessments of aquatic life use support as part of Integrated Reporting.

From 2002 through 2006, Iowa DNR, in cooperation with the State Hygienic Laboratory at the University of Iowa (SHL), conducted biological sampling as part of a Regional Environmental Monitoring and Assessment Program (REMAP) project designed to randomly select Iowa stream sites over four years to objectively measure biological integrity in flowing streams. This project was based on a random sampling design was used to obtain an unbiased sample population from which accurate statements about the status of Iowa's perennial streams can be extrapolated. Approximately 60 sites a year were sampled and included measures of several indicators of stream ecosystem health including: fish tissue, sediment, and water contaminant levels; physical habitat quality; and fish and benthic macroinvertebrate populations. This study was designed to determine the current biological health of Iowa streams and help provide a uniform assessment of stream conditions in the Central Plains of the United States.

- **Data from the IDNR-sponsored lake monitoring conducted by Iowa State University and the University of Iowa Hygienic Laboratory**

Data from statewide surveys of Iowa lakes completed in the early 1980s (110 lakes) and early 1990s (115 lakes) by Iowa State University served as the basis for past assessments of lake water quality in Iowa. Beginning in 2000, 131 lakes throughout Iowa were monitored annually as part of an IDNR-sponsored five-year project to assess their condition and measure the temporal variability in lake water quality. This monitoring was

being conducted by Iowa State University. All lakes assessed as part of the 1990/1992 statewide lake survey were sampled as well as 16 additional lakes.

Each lake is sampled three times during the summer season to assess seasonal variability. Samples are taken at the deepest point in each lake basin. Vertical probes are lowered through the water column to determine vertical profiles for temperature, dissolved oxygen, specific conductivity, pH, turbidity, and chlorophyll. An integrated column sampler is used to collect water from the upper mixed zone in thermally stratified lakes and from the entire water column in lakes that lack stratification.

In 2005, IDNR sponsored lake monitoring by SHL to supplement the ISU survey. Samples were collected at the 131 lakes sampled as part of the ISU study to expand the summer season monitored from three months (typically June, July, and August) to include samples from May, September and October. The SHL samples have been collected and analyzed with field methods and laboratory procedures comparable to those used in the ISU study. In 2006, five monthly samples were collected as part of IDNR/SHL lake monitoring at all 131 lakes. The data from these samples are used to develop water quality assessments for the lakes monitored.

- **Data from monitoring of bacterial indicators in rivers and at beaches of publicly-owned lakes**

Indicator bacteria, such as fecal coliform bacteria and *E. coli*, are commonly monitored by state environmental agencies to indicate the degree to which surface waters support their designated uses for primary contact recreation. High levels of these indicator bacteria suggest that using a river or lake for either primary contact recreation (e.g., swimming or water skiing) or secondary contact recreation (e.g., wading while fishing) presents a health risk due to the potential for waterborne diseases. As part of fixed station monitoring networks in Iowa, river and stream reaches designated for primary or secondary contact recreation uses are monitored for bacterial indicators on a monthly basis.

Historically, this type of monitoring had not been conducted at Iowa's lakes. In 1999, however, the IDNR Division of Parks, Recreation and Preserves monitored ten of Iowa's public beaches for indicator bacteria. In 2000, beach monitoring was expanded to thirty-one Iowa beaches and was placed under the direction of IDNR's Watershed Monitoring and Assessment Section. From May through September, these beaches were monitored

weekly. All beaches were monitored for three U.S. EPA-recommended bacterial indicators: fecal coliform, enterococci, and *E. coli*. Since 2001, annual monitoring at approximately thirty-five beaches at state-owned lakes as been conducted on a weekly basis during summer recreational seasons.

In addition, 28 beaches at city and county-owned lakes were monitored for indicator bacteria during the period 2006 through 2008. The data from this monitoring is available in the Iowa STORET/WQX water quality database (<http://programs.iowadnr.gov/iastoret/>). These data will also be evaluated to determine the degree to which primary contact recreation (Class A1) uses are supported. The Iowa beaches monitored for indicator bacteria during the 2006-2008 period, including state-owned as well as city and county-owned beaches, can be found in Table 5.

- **Data from programs to monitor fish tissue for toxic contaminants**

Annual, routine monitoring for bioaccumulative toxics in Iowa fish tissue is conducted as part of three long-term programs: (1) U.S. EPA Region VII's *Regional Ambient Fish Tissue Monitoring Program*, (2) water quality studies of the Des Moines River near Saylorville and Red Rock reservoirs conducted by Iowa State University under contract with the U.S. Army Corps of Engineers, and (3) water quality studies of the Iowa River near Coralville Reservoir conducted by the University of Iowa also under contract with the U.S. Army Corps of Engineers.

Since 1980, annual fish collection and analysis activities in Iowa have been conducted by IDNR as part of the U.S. EPA Region VII's *Regional Ambient Fish Tissue (RAFT) Monitoring Program*. Each year in late summer, IDNR fisheries biologists collect fillet samples of both bottom-feeding fish (common carp or channel catfish) and predator fish (usually largemouth bass, crappie, or walleye) from approximately 30 locations on rivers and lakes in Iowa. Selection of sample sites is based on the level of fishing use and date of most recent fish tissue sampling. Currently, samples are analyzed for 19 pesticides, four organic compounds, and four metals. The RAFT program also involves (1) monitoring for trends in levels of toxics in bottom feeding fish (common carp, *Cyprinus carpio*) at ten fixed sites on Iowa's larger rivers as well as (2) follow-up monitoring designed to verify the existence of high contaminant levels and to determine whether the issuance of consumption advisories is justified. Annual reports for RAFT monitoring in Iowa can be found at <http://www.igsb.uiowa.edu/wqm/Biological/RAFT.html>.

Iowa State University (Department of Civil Engineering, Environmental Engineering Section) conducts annual fish contaminant monitoring for bottom-feeding fish (common carp) at Saylorville and Red Rock reservoirs as part of a U.S. Army Corps of Engineers water quality monitoring program (see <http://home.eng.iastate.edu/~dslutz/dmrwqn/dmrwqn.html>). The University of Iowa and Iowa State University have conducted fish contaminant monitoring as part of a similar program at Coralville Reservoir.

Fish contaminant monitoring is also conducted on an occasional basis as part of special studies of water quality. For example, the Kansas City District of the U.S. Army Corps of Engineers periodically conducts fish contaminant monitoring at Rathbun Reservoir in southern Iowa. Also, fish contaminant monitoring was conducted over a 13-year period (1988-2000) in Pool 15 of the Upper Mississippi River near Davenport, Iowa, in response to a PCB contamination problem.

- **Reports of pollutant-caused fish kills**

IDNR routinely receives reports of fish kills that are investigated by IDNR staff from the Fisheries Bureau and/or the Compliance & Enforcement Bureau. Information from the reports of these kills, including location, the cause and source of the kill, the size of waterbody affected, and the number of fish killed, is entered into the IDNR Fish Kill Database (see <http://programs.iowadnr.gov/fishkill/default.aspx>).

- **Data from public water supplies on the quality of surface water sources and finished water**

The IDNR Environmental Services Division administers the public drinking water program in Iowa under delegation of authority from the U.S. Environmental Protection Agency. As required by the Safe Drinking Water Act of 1996, IDNR prepares an annual report of violations of national primary (finished) drinking water standards by public water supplies in the state (reports are available at <http://www.iowadnr.gov/water/drinking/reports.html>). For the 2010 assessment/listing cycle, reports for 2006 through 2008 were reviewed for violations (IDNR/WQB 2007, 2008, 2009). In addition, several public water supplies using surface water sources in Iowa have generated long-term databases for the quality of raw water used at their facilities. For example, the municipal water supplies at Cedar Rapids and Des Moines routinely collect data on levels of toxic contaminants in the Cedar River

and the Raccoon/Des Moines rivers, respectively, which can influence their water treatment processes.

Since 1994, Syngenta, Inc. has sponsored a voluntary program to monitor levels of atrazine in Iowa several impoundments used as a source of potable water for a municipal water supply. During the period 2006-2008, this program included surface water supplies for nine Iowa municipalities and their respective source waters. Although surface water supplies for all nine communities were monitored in 2006, monitoring ended at Lower Centerville Reservoir (Centerville), Home Pond (Lamoni), and Little River Lake (Leon) after 2006, and monitoring ended at Rathbun and Corydon reservoirs after 2007:

Water Supply Monitored for atrazine by Syngenta:	Primary Source Water	Monitored in 2006?	Monitored in 2007?	Monitored in 2008?
Centerville	Lower Centerville Reservoir	Yes	No	No
Chariton	Lake Morris	Yes	Yes	Yes
Creston	Three Mile Lake	Yes	Yes	Yes
Lamoni	Home Pond	Yes	No	No
Leon	Little River Lake	Yes	No	No
Montezuma	Diamond Lake	Yes	Yes	Yes
Osceola	West Lake Osceola	Yes	Yes	Yes
Rathbun Land & WQ Alliance	Rathbun Reservoir	No	Yes	No
Rathbun Regional Water Alliance	Corydon Reservoir	Yes	Yes	No
Winterset	Cedar Lake	Yes	Yes	Yes

Data from this monitoring program are available via U.S. EPA at the following website: http://www.epa.gov/pesticides/reregistration/atrazine/atrazine_update.htm. Although not part of Syngenta's voluntary monitoring program, atrazine monitoring continued at Corydon Reservoir in Wayne County through 2007. Although no longer used as a source of a public drinking water supply, this reservoir was monitored in agreement with the Iowa DNR as part of the atrazine TMDL for this waterbody.

- **Data from special studies of water quality and aquatic communities**

Special/intensive studies of water quality are typically conducted over a finite time period and are targeted toward understanding or characterizing specific water quality issues. This type of study differs from “routine” monitoring that is conducted over a long time frame and that typically generates information necessary to describe general water quality conditions. The sampling protocol for intensive studies is site-specific and is based on the contaminant(s) of concern. These studies typically require multiple samples per site over a relatively short time frame. If the contaminants of concern have significant seasonal or daily variation, season of the year and time of day variation are accounted for in sampling design. The number of sampling sites, sampling frequency and parameters vary depending on the study.

Each year, a number of special water quality studies are conducted in the state; these studies include monitoring conducted in support of TMDL development. Results of special studies may be summarized in the form of a published document, an unpublished report, or may exist only as raw data. For example, recent studies on the South Fork Iowa River watershed (Tomer et al. 2008) and on the streams of the Mesquaki Indian Settlement (Littin and McVay 2008) contain data useful for assessing support of designated uses. Surveys of aquatic communities are occasionally conducted by IDNR staff as part of special studies. Special water quality studies conducted by colleges and universities as part of undergraduate and graduate projects are also potential sources of water quality data and other water-related information.

- **Best professional judgment of IDNR staff**

IDNR utilizes observations of professional staff of the IDNR bureaus of Fisheries and Wildlife, as well as professional staff in other agencies, to assess support of aquatic life uses in certain types of Iowa waterbodies that have historically lacked chemical, physical, and/or biological water quality data. For example, due to the lack of relevant criteria for assessing wetland quality, water quality assessments for these waterbodies have been based primarily on observations of biologists in the IDNR Wildlife Bureau. Although limited wetland water quality sampling was conducted during the 2006-2008 period, and although several wetland assessments were based on results of this monitoring, the majority of wetland assessments remains based primarily on best professional judgment.

- **Results of volunteer monitoring**

The Iowa volunteer monitoring program (IOWATER) was established in 1999 by the IDNR. This program provides training, equipment and supplies to volunteers for monitoring streams throughout Iowa. Data generated by the IOWATER program can be found on the IDNR web site at <http://www.iowater.net/database/online.asp>. A review of the IOWATER database by IDNR staff in 2002 showed considerable variation in data quality within this database. Due to the often unexplained variation, IDNR staff decided not to use results of volunteer monitoring for Section 305(b) assessments. In addition, Iowa's credible data law passed in 2000 resulted in state regulations that place restrictions on the use of volunteer data for purposes of adding waterbodies to Iowa's Section 303(d) list; these regulations became effective in 2003. These regulations can be found under "Volunteer Monitoring Data Requirements" in the *Iowa Water Quality Standards* (Chapter 61.10, IAC; <http://www.iowadnr.gov/water/standards/files/chapter61.pdf>). These restrictions include a requirement for preparation of a monitoring plan by the volunteer monitor and review and approval of this plan by IDNR before the volunteer data can be used for purposes of Section 303(d) listing. If, however, volunteer monitors encounter and document instances of gross pollution such that water quality conditions that appear to violate Iowa's narrative water quality standards at IAC 61.3(2), IDNR will consider use of this information for purposes of Section 303(d) listing as described in the section of this methodology on "overwhelming evidence of impairment." IDNR staff that direct the IOWATER program are consulted to help identify instances of gross pollution discovered through IOWATER monitoring. Also, any data collected by volunteer monitors that meet Iowa's credible data requirements will be considered for identifying Section 303(d) impairments.

Identifying impairments:

As specified in U.S. EPA's regulations for TMDLs (40 CFR 130.7), sources of existing and readily available water quality-related data and information to be considered as part of Section 303(d) listing include but are not limited to the following:

- the state's most recent CWA Section 305(b) report;
- CWA Section 319 nonpoint source assessments;
- dilution calculations, trend analyses, or predictive models for determining the physical, chemical or biological integrity of streams, rivers, lakes, and estuaries; and
- water quality-related data and information from local, State, Territorial, or Federal agencies [in Iowa, especially the U.S. Geological Survey's National Water Quality Assessment Program

(NAWQA) and National Stream Quality Accounting Network (NASQAN)), tribal governments, members of the public, and academic institutions].

The majority of information used by IDNR to develop the Section 303(d) list of impaired waters (IR Category 5) is taken from the most recent Section 305(b) assessments for the state of Iowa. As noted in this methodology, IDNR staff attempt to utilize water quality data and related information from a variety of sources. IDNR has not, however, used results of dilution calculations or predictive models to add waterbodies to Iowa's Section 303(d) list. Due to the importance of data quality and quantity in developing accurate assessments, and due to requirements of Iowa's credible data law that require site-specific, high-quality data upon which to base listings, only a subset of the available 305(b) information is used for purposes of placing waters into Category 5. The process of determining whether or not data from the above data sources are appropriate for placing waterbodies in Category 5 is described below.

Types of Assessments: *Evaluated and Monitored:*

For purposes of developing Section 305(b) assessments, the existing and readily available water quality data described above are used to make two types of water quality assessments: "evaluated" and "monitored." As described in guidelines for Section 305(b) reporting (U.S. EPA 1997, pages 1-5 and 1-9 [see <http://www.epa.gov/owow/monitoring/guidelines.html>]),

Evaluated waters are

those for which the use support decision is based on water quality information other than current site-specific data such as data on land use, location of sources, predictive modeling using estimated input values, and some questionnaire surveys of fish and game biologists. As a general rule, if an assessment is based on older ambient data (e.g., older than five years), the State should also consider it "evaluated."

For example, water quality assessments based on results from only a few grab samples or on professional judgment of local biologists, in the absence of any supporting data, would be considered "evaluated" assessments.

Monitored waters are

those for which the use support decision is principally based on current, [five years old or less] site-specific ambient monitoring data believed to accurately portray water quality conditions. Waters with data from biosurveys should be included in this category along with waters monitored by fixed-station chemical/physical monitoring or toxicity testing. To be considered "monitored" based on fixed station chemical/physical monitoring, waters generally should be sampled quarterly or more frequently.

Although EPA's guidelines for integrated reporting (U.S. EPA 2005, 2006) do not distinguish between "monitored" and "evaluated" assessments, Iowa DNR feels that the distinction remains important for determining the relative scientific strength and confidence of the water quality assessments developed. In addition, this distinction allows IDNR to better target assessed waters for additional monitoring, and is the basis for identifying waters in need of additional monitoring. Thus the on-line Iowa assessment database (ADBNet [<http://programs.iowadnr.gov/adbnnet/index.aspx>]) is designed to track "monitored" versus "evaluated" assessments while still complying with the integrated reporting format recommended by U.S. EPA (2005).

In terms of the ability of Section 305(b) assessments to characterize current water quality conditions, IDNR considers evaluated assessments as having relatively lower confidence while monitored assessments are of relatively higher confidence. This approach is consistent with guidance from U.S. EPA (U.S. EPA 1997). IDNR considers monitored assessments as sufficiently accurate to be appropriate for both Section 305(b) assessments and Section 303(d) listing (i.e., for placing waters into Category 5 of the integrated report). The lower confidence evaluated assessments, however, are viewed as appropriate only for Section 305(b) reporting. Thus, any waters "evaluated" as "impaired" are placed in IR Categories 2b or 3b (i.e., categories for potentially impaired waterbodies with insufficient information for determining whether uses are met). Such waters are added to Iowa's list of "waters in need of further investigation" (WINOFI list) as provided for in Iowa's credible data law and will be considered for follow-up monitoring to better determine current water quality conditions and the existence of any impairments.

Magnitude of Impairment:

In addition to IDNR's retention of the distinction between "monitored" and "evaluated" waters, IDNR tracks the magnitude of the impairment in its Assessment Database (ADBNet). These impairment magnitudes include the following:

Fully supported/threatened: Water continues to fully support the designated use but an adverse water quality trend is evident such that the water will likely fail to fully support the designated use by the time of the next listing cycle.

Partially supported: A slight to moderate impairment suggested by occurrence in the lower impairment range. The following examples would result in an impairment magnitude of "partially supported": water quality criteria violation frequency greater than 10% but less than 25%; score

for only one of the two indexes of biotic integrity in the impairment range; one pollutant-caused fish kill during the triennial period; lower tier of fish consumption advisories (one meal/week) in effect; geometric mean for *E. coli* greater than the respective criterion but are less than eight times the criterion.

Not supported: A severe impairment suggested by occurrence in the middle to upper impairment range (e.g., water quality criteria violation frequency greater than 25%; scores for both indexes of biotic integrity (fish and aquatic macroinvertebrates) in the impairment range; more than one pollutant-caused fish kill during the triennial period; upper tier of fish consumption advisories (“do not eat”) in effect; geometric mean for *E. coli* greater than eight times the respective criterion.

Tracking the relative severity of impairments allows IDNR to more efficiently prioritize impaired waters for TMDL development and allows more accurate reporting on water quality conditions in the state.

Data quantity considerations (“data completeness” guidelines):

For purposes of Section 303(d) listing in Iowa (i.e., placing waters in Category 5), data quantity issues are addressed in this methodology. Beginning with Iowa's Section 305(b) report for 1990, IDNR staff developed “data completeness” guidelines to avoid basing water quality assessments on inadequate amounts of water quality data and to reduce errors in assessments (for example, incorrectly concluding that an impairment exists). For the various parameters used to develop water quality assessments, these guidelines establish the minimum number of data points needed over a given assessment period to adequately determine whether the applicable water quality standards are being met. Assessments that meet these data completeness guidelines are of relatively high confidence and are considered “monitored.” Assessments based on an insufficient amount of data to meet these guidelines are of relatively low confidence and are thus considered “evaluated.” IDNR's interpretations of the terms “evaluated” and “monitored” are identical to those of U.S. EPA (1997). IDNR's Section 305(b) data completeness guidelines are presented in Table 6. The significance of data completeness guidelines and Iowa's credible data law to Iowa's Section 305(b) water quality assessments and Section 303(d) listings is summarized in Figure 1.

Data quality considerations (“credible data” requirements):

As defined by U.S. EPA, *data quality objectives* are qualitative and quantitative statements that clarify objectives, define appropriate types of data, and specify levels of potential decision errors that will be used as the basis for establishing the quality and quantity of data needed to support decisions. In this

context, Iowa's credible data law (Attachment 1) defines the appropriate types of data for developing the state's Section 303(d) listings. These objectives are as follows:

- "*Credible data*" means scientifically valid chemical, physical, or biological monitoring data collected under a scientifically accepted sampling and analysis plan, including quality control and quality assurance procedures.
- Data dated more than five years before the department's date of listing or other determination under section 455B.194, subsection 1 (Iowa's credible data law), shall be presumed not to be credible data unless the department identifies compelling reasons as to why the data is credible.

As stated in the 2001 Iowa Code, Section 455B.194, subsection 1, the department shall use "credible data" when doing any of the following:

- Developing and reviewing any water quality standard.
- Developing any statewide water quality inventory or other water assessment report. (Note: Iowa's Section 305(b) assessments are not subject to the provisions of Iowa's credible data law.)
- Determining whether any water of the state is to be placed on or removed from any Section 303(d) list.
- Determining whether any water of the state is supporting its designated use or other classification. (Note: the credible data law does not require the use of credible data for establishment of a designated use or other classification of a water of the state.)
- Determining any degradation of a water of the state under 40 CFR 131.12.
- Establishing a total maximum daily load (TMDL) for any water of the state.

The credible data law has occasionally been criticized as being an obstacle to the addition of impaired waters to Iowa's Section 303(d) list. This criticism is directed at the requirement that data older than five years are presumed not to be credible. Because, however, all water quality data are reviewed biennially and assessed for Section 303(d) impairments as the data become available, and because most water quality data in Iowa are generated Iowa DNR, its designees, or other government agencies, the credible data requirements rarely influence IDNR's listing decisions. Thus, such criticism is largely unfounded.

Rationale for any decision not to use existing and readily available data for Section 303(d) listings:

IDNR reviews all existing and readily available water quality-related data and information for purposes of water quality reporting and impaired waters listing as required by Sections 305(b) and 303(d) of the Clean Water Act (see section on [Sources of Existing and Readily Available Water Quality Data](#) in this methodology). Certain categories of water quality information, however, do not meet requirements of either Iowa's credible data law or IDNR's data completeness guidelines for water quality assessments and impaired waters listings. The ultimate reasons for not using certain "existing and readily available data" are (1) the need for reasonably accurate assessments of water quality and (2) the desire to add only waterbodies to the state's Section 303(d) list (Category 5) that are actually "impaired." Placing waters on the state's Section 303(d) list on the basis of inaccurate and/or incomplete data increases the risk that the department's limited resources, including staff time and monitoring dollars, will be used unwisely. Examples of water quality information that typically would not be considered appropriate as the basis for Section 303(d) listing include the following:

- **Best professional judgment of IDNR staff:** IDNR utilizes observations of professional staff of the IDNR bureaus of Fisheries and Wildlife, as well as professional staff in other agencies for purposes of water quality (Section 305(b)) reporting. Best professional judgment is used to assess support of aquatic life uses for certain types of Iowa waterbodies that have historically lacked chemical, physical, and/or biological water quality data (primarily wetlands). To be added to Iowa's list of impaired waters (Category 5), all assessments of impairment based solely on best professional judgment will be further investigated to better document any failure to meet water quality standards. Past experience with assessments based primarily on best professional judgment has demonstrated that such follow-up investigations are necessary to (1) better determine whether a Section 303(d) water quality impairment actually exists and (2) more accurately identify the causes and sources of any existing impairment. Field biologists and other field staff are extremely knowledgeable regarding the water resources they manage but are much less knowledgeable regarding the intent and constraints of Clean Water Act Section 303(d) listing. Waters assessed as "impaired" based only on the basis of best professional judgment will be added to Categories 2b or 3b of the Integrated Report; these two categories comprise the list of "waters in need of further investigation" (WINOFI list) as provided for in Iowa's credible data law.
- **Data or information older than five years from the end of the most recent Section 305(b) reporting cycle:** Data dated more than five years before the end of the current (2010) Section 305(b) data consideration period (the end of calendar year 2008) are presumed under state law to be "not credible" unless IDNR identifies compelling reasons as to why these older

data are credible. This provision of Iowa's credible data law was based on, and is consistent with, U.S. EPA's (1997) recommendation that data older than five years should not be used to make the type of water quality assessment (a "monitored" assessment) that is believed to accurately portray site-specific water quality conditions. Data older than five years, however, may be used for identifying water quality trends for any water of the state for which credible data exist. Historically, data older than five years have been routinely used for Section 305(b) reporting in Iowa but have not been used to identify new Section 303(d) listings. All such assessments are considered "evaluated" and are thus of relatively lower confidence than "monitored" assessments which are based primarily on recent, site-specific ambient monitoring.

As the data upon which non-303(d) assessments are based age beyond five years—and if more recent data are not available—the assessment type is changed from "monitored" (higher confidence) to "evaluated" (lower confidence) as part of the biennial Section 305(b) assessment process. Once placed in IR Category 5 (i.e., on the state's Section 303(d) list), however, a waterbody will not be moved to a non-TMDL category without "good cause" as defined by U.S. EPA regulations at 40 CFR 130.7 (e.g., a TMDL for the waterbody is approved by EPA or new monitoring data suggest that the impairment no longer exists). U.S. EPA regulations do not consider the age of data as a "good cause" for removing a Section 303(d) impairment.

The issue of "old data" is seldom relevant to Section 303(d) listing in Iowa. Water quality data are used for developing the biennial Section 305(b) assessments as they become available and are thus considered for Section 303(d) listing when the data most likely represent current water quality conditions. This process occurs long before the data age beyond their ability to accurately represent current water quality conditions. As the data age beyond five years, the Section 305(b) assessment type is changed from "monitored" to "evaluated" to reflect the lowered level of confidence in assessments based on older data that potentially may not represent current water quality conditions. Any non-303(d) Section 305(b) assessments based on data that have aged beyond 10 years are not included in the current assessment cycle, but the previous assessments based on these data remain in IDNR's on-line assessment database (Iowa ADBNet [<http://programs.iowadnr.gov/adbnet/index.aspx>]).

- **Data that do not meet "completeness guidelines" developed for Section 305(b) reporting:** In order to improve the accuracy of water quality assessments, IDNR has

identified “data completeness guidelines” for using results of routine water quality monitoring for Section 305(b) reporting (Table 6). These guidelines identify the numbers of samples needed for water quality assessments that can support Section 303(d) listings (i.e., *monitored* assessments). These guidelines also identify assessments appropriate only for Section 305(b) reporting (i.e., *evaluated* assessments). These criteria were first developed for Iowa’s 1990 Section 305(b) report and are designed to improve--within the constraints of (1) resources available for monitoring and (2) the designs of existing monitoring networks--the accuracy of Section 305(b) water quality assessments. The improvement in assessment accuracy increases the confidence with which waterbodies are added to Iowa’s Section 303(d) list. Although IDNR ambient water quality monitoring networks and networks of other agencies are designed to produce sufficient data to meet Iowa’s “completeness guidelines,” not all monitoring activities are so-designed. Thus, the use of these guidelines will eliminate certain data from consideration for Section 303(d) listing. Any waterbodies assessed as “impaired” only on the basis of incomplete data, however, will be placed in IR Categories 2b or 3b and will be added to the state list of “waters in need of further investigation” as provided for in Iowa’s credible data law.

- **Results of volunteer monitoring that do not meet requirements specified in Iowa’s credible data legislation and/or Section 305(b) data completeness guidelines:** Results from volunteer monitoring can only be used for Section 303(d) listing if requirements of Iowa’s credible data law are met or if overwhelming evidence of impairment is indicated. To be considered for Section 303(d) listing, IDNR rules [IAC 61.10 through IA 61.13 (455B)] require that volunteer monitoring must be supported by an IDNR-approved sampling and analysis plan that includes quality control and quality assurance procedures. Waterbodies assessed as “impaired” only the basis of volunteer data from non-qualified volunteers will not be added the Iowa’s Section 303(d) list but may be added to the state list of waters in need of further investigation. If, however, results of volunteer monitoring show the existence of gross pollution such that Iowa’s narrative criteria are violated, such waters can be added to Iowa’s Section 303(d) list due to overwhelming evidence of impairment.
- **Results of habitat assessment:** Although detailed information on the quality of aquatic habitats is collected as part of biological monitoring conducted for the IDNR/SHL stream biocriteria and REMAP projects, this information is not directly used to identify Section 303(d) impairments of aquatic life uses. IDNR does, however, incorporate observations on the quality of aquatic habitat into Section 305(b) water quality assessments and biologically-based

Section 303(d) listings. This information is also used in the identification of causes and sources of impairments of aquatic life uses identified through biological monitoring.

- **Assessments of headwater stream segments.** As explained below, Section 303(d) impairments based on results of chemical/physical water quality monitoring on headwater stream segments will be added to Iowa's Section 303(d) list. Due to the lack of a calibrated biological assessment protocol, however, impairments based on results of biological monitoring in headwater segments will not be placed on the Section 303(d) list but will be placed into IR Categories 2b or 3b and added to Iowa's list of waters in need of further investigation.

The aquatic environment of most of Iowa's small headwater streams is one of extremes ranging from flood-flow to no-flow; from completely frozen in winter to extremely warm water temperatures in summer. Due to their position in relation to sources of groundwater, many headwater stream reaches experience no-flow conditions at least once per year. These extremes are sometimes reflected in results of water quality monitoring and biological assessments that suggest impairment. For example, as streams move toward no-flow conditions during summer due to low amounts of precipitation, chemical water quality can degrade drastically, especially regarding levels of dissolved oxygen and pH. As stream flow ceases and the only remaining water exists as isolated and shrinking pools, violations of water quality criteria for dissolved oxygen and/or pH become more common, often with sufficient frequency to suggest impairment of aquatic life uses. Also, due to seasonally reoccurring intermittent flow, the types of aquatic life that inhabit general use streams are often only those able to withstand extremes environmental conditions (the so-called "pioneer species"). Consequently, headwater stream segments tend to have water quality and biological diversity that are low relative to the larger and more ecologically stable stream environments.

Historically, Iowa's headwater stream reaches were typically not designated for protection of either primary contact recreation or aquatic life uses but were instead classified only for protection of "general uses" such as livestock and wildlife watering, aquatic life, noncontact recreation, crop irrigation, and industrial, agricultural, domestic and other incidental water withdrawal uses (Table 9). According to the *Iowa Water Quality Standards* (Section 61.3(2)), general use waters are protected by narrative criteria designed to prevent aesthetically objectionable/nuisance conditions, and other forms of gross pollution attributable to pollution sources (IAC 2006). In contrast, Class A and Class B waters are also protected by numeric

criteria designed to protect human health from recreationally-related waterborne diseases and to protect aquatic life from chronically toxic as well as acutely toxic conditions.

Due to changes in the *Iowa Water Quality Standards* that became effective in March 2006, all perennially-flowing streams and intermittent streams with perennial pools are now presumed to be capable of supporting the highest level of primary contact recreation use and the highest level of aquatic life use. This approach to applying designated uses is called the “rebuttable presumption” (see <http://www.iowadnr.com/water/standards/protectedflow.html>). Under this approach, the Class A1 (primary contact recreation) use is presumptively applied to all of Iowa’s perennial rivers and streams and intermittent streams with perennial pools, and the Class B(WW1) aquatic life use is presumptively applied to all of Iowa’s perennial rivers and streams and intermittent streams with perennial pools unless the water is already designated for Class B(WW2) or Class B(WW3) uses in Iowa’s surface water classification (see http://www.iowadnr.com/water/standards/files/06mar_swc.pdf). A “use attainability analysis” or UAA must be conducted, including field investigations, to determine whether the presumptively-applied use is, in fact, the appropriate designated use for the stream segment in question. For more information on UAAs, please see the IDNR fact sheet, *Assessing Iowa Stream Uses* (http://www.iowadnr.com/water/files/uaa_factsheet.pdf).

Assessments of headwater stream segments based on chemical/physical water

quality data: Because the distinction between a truly intermittent (and thus, general use-only) stream and an “intermittent stream with perennial pools” is currently poorly defined, monitoring data from all currently non-designated and formerly “general use” headwater stream segments will be assessed against the presumptively-applied Class A1/Class B(WW1) water quality criteria for purposes of Section 305(b) assessments and Section 303(d) listings. Any Section 303(d) impairments identified for a presumptively designated stream segment will be placed into state-defined Category 5p (i.e., “5-presumptive”) of Iowa’s Integrated Report. IDNR staff that prepare Iowa’s Section 303(d) list will coordinate with IDNR Water Quality Standards Section staff to determine, to the degree possible, whether UAAs have been conducted for the presumptively-impaired stream segments. If the appropriate uses have been determined through a UAA, the impairment will be placed in IR Category 5a (pollutant-caused impairment) as appropriate.

Assessments of headwater stream segments based on biological data: Biological monitoring is occasionally conducted on Iowa’s headwater stream segments (i.e., having

watersheds draining less than about 10 square miles). Thus, the use of biological assessment methods developed for the larger, more stable, and more diverse streams to assess headwater segments will likely overstate the existence of impairment. For this reason, headwater stream segments that show impairment based on a failure to meet regional expectations for aquatic biota (fish or aquatic macroinvertebrates) of Class B(WW2) streams, will not be added to Iowa's Section 303(d) list of impaired waters. The assessment type for these waters will be considered "evaluated" (indicating an assessment with relatively lower confidence) as opposed to "monitored" (indicating an assessment with relatively higher confidence). Such waters will be placed in either IR Category 2b-u or Category 3b-u (i.e., potentially impaired based on un-calibrated assessment) and will be added to the state's list of "waters in need of further investigation" as provided for in Iowa's credible data law. Once on this list, the assessments can be reviewed to better determine the nature of the water quality problems suggested by biological monitoring and to determine whether follow-up monitoring is justified. See Attachment 2 of this methodology for additional information on IDNR's approach for biological assessment of Iowa's headwater streams. IDNR staff continue to pursue development of an assessment protocol for headwater streams segments.

List of waters in need of further investigation:

Although not appropriate for identifying Category 5 (Section 303(d)) waters, the above types of water-related information can be used for Section 305(b) water quality assessments and thus can be used to place waterbodies on a separate list of Iowa waterbodies in need of further investigation (WINOFI list). As provided for in Iowa's credible data law, the WINOFI list is not part of the Section 303(d) process in Iowa and includes waterbodies where limited information suggests, but does not conclusively or credibly demonstrate, a water quality impairment. The state's WINOFI list is comprised of those waterbodies assessed (evaluated) as potentially "impaired"; that is, the assessment of a designated use in these waterbodies as "impaired" is based on less than complete information; thus, the assessment is of relatively low confidence and is not appropriate for addition to the list of Section 303(d) waterbodies. These potentially-impaired waters are thus placed in subcategories 2b and 3b of the Integrated Report which comprises the list of waters in need of further investigation. Category 2 of the IR is for waters where at least one designated use is fully supported but insufficient information is available to assess the remaining uses; Category 3 is for waters where information is not sufficient to assess any designated use.

If the results of further investigative monitoring demonstrate with credible data that a water quality impairment exists, the affected waterbody can be added to Iowa's Section 303(d) list (IR Category 5). Through this process, 50 WINOFI waters were added to Iowa's 2004 Section 303(d), five WINOFI waters were added to Iowa's 2006 list, four waters were added to the 2008 list, and nine waters were added to Iowa's draft 2010 list.

Overwhelming evidence of impairment:

Situations exist where reliable information can accurately indicate a Section 303(d) impairment of designated beneficial uses even though this information does not meet the IDNR requirements for Section 303(d) listing (Table 6). Such waterbodies would be considered for addition to IR Category 5 (=Section 303(d) list) of Iowa's integrated assessment/listing report. The following are examples of instances where overwhelming evidence justifies determination of impairment in the absence of complete data:

- Presence of reoccurring, man-made circumstances that result in acutely toxic conditions for aquatic life. For example, the addition of untreated septic waste to a stream via an illegal connection to a storm sewer such that the aquatic community is being severely impacted would constitute overwhelming evidence of impairment.
- Man-made alterations of hydrology, flow, or habitat that degrade the quality of aquatic habitats as reflected in significant, adverse deviations in biotic integrity from the reference condition or from the pre-modification aquatic communities. For example, an illegal channel change that adversely affects the aquatic community of a stream reach would constitute overwhelming evidence of impairment.
- Chronic de-watering of a considerable section of a waterbody related to man-made alterations of local hydrology. For example, an illegal water withdrawal for irrigation that severely impacts or eliminates the aquatic life of a stream or river constitutes overwhelming evidence of impairment.
- Presence of exotic species (e.g., common carp or purple loosestrife (*Lythrum salicaria*)) at levels that are believed to impair one or more designated uses. For example, the infestation of a wetland with the invasive exotic plant purple loosestrife such that the value of a wetland for use by waterfowl is degraded constitutes overwhelming evidence of impairment.
- Summer median trophic state index (Carlson 1977, 1991) values for chlorophyll-a or secchi depth that are based on less than three years of data but that are more than five TSI points greater than the

TSI value used to identify impairment with a complete dataset (a “complete dataset” is three or more years of data resulting from three to five samplings per year). For example, if a lake’s median based summer chlorophyll-a TSI value from one year’s monitoring (minimum of three samples) exceeds the IDNR’s trigger value of TSI = 65 by more than five points, the lake would be assessed as Section 303(d) impaired due to overwhelming evidence of impairment (for more information on IDNR’s use of Carlson’s trophic state index, see Attachment 3 of this methodology).

- The *E. coli* geometric mean of at least five samples collected at regular intervals over a summer recreational season, and that meet credible data requirements, would exceed Iowa’s geometric mean criterion even if the remainder of the 10 samples needed for a high-confidence (“monitored”) assessment all had less than the IDNR’s detection level for *E. coli* (i.e., 10 orgs/100 ml).

How water quality data and other water-related information are summarized to determine whether waters are Section 303(d) “impaired”:

- **Physical, chemical, and bacterial data from fixed station water quality monitoring networks:**

These types of data are used with methods for Section 305(b) water quality assessments developed by U.S. EPA, with some of these methods being modified by IDNR (see Tables 6 through 12).

Conventional Parameters: U.S. EPA’s (1997) Section 305(b) assessment guidelines specify that aquatic life uses of surface waters with more than 10% of samples in violation of state water quality criteria for conventional parameters (for example, dissolved oxygen, pH, temperature) should be assessed as “impaired.” This assessment approach is sometimes referred to as “the 10 percent rule”. IDNR has historically not used the 10-percent rule to assess water quality with datasets of less than 10 samples due to the large degree of uncertainty associated with basing impairment decisions on small datasets. The IDNR requirement for at least 10 samples was based on the resultant improvement in the ability of U.S. EPA’s recommended assessment approach to accurately identify an impairment based on a critical value of 10% violation. For example, at sample sizes less than 10, the probability of incorrectly concluding that impairment exists (Type 1 error) with U.S. EPA’s approach is approximately 60%; with 10 samples, the probability of this type of error decreases to approximately 30% (Smith et al. 2001). Despite this approach, the percentage of a Type I error remains high (30 percent). In addition, comparison of raw

percentages to water quality criteria have often been problematic in that they seem to give a contradictory signal of impairment. The most common scenario is the following: more than 10 percent of samples exceed the criterion for pH or dissolved oxygen (thus indicating "impairment") while all other water quality indicators suggest "full support."

Recently, however, alternative approaches have been developed that (1) avoid the need to compare raw percentage values to state criteria to identify impairments and (2) incorporate estimates of the numbers of samples and the corresponding number of violations that represent a significant exceedance of the 10 percent rule. The state of Nebraska (NDEQ 2006), drawing on information from Lin et al. (2000), adopted an assessment approach where the sample sizes and the corresponding number of violations needed to identify a significant exceedance of the 10%-rule with greater than 90 percent confidence are specified. This approach is based on the binomial method for estimating the probability of committing Type I and Type II errors. Nebraska's 2010 assessment methodology is available at <http://www.deq.state.ne.us/>. Table 1 from the Nebraska methodology (*Sample size and number of exceedances required to determine an impaired beneficial use (10% exceedance)*) is included in this methodology as Table 12. IDNR first used this binomial-based approach for identifying impairments based on violations of the 10% rule for the 2006 assessment/listing cycle and has continued to use this approach.

Toxic parameters: U.S. EPA (1997) guidelines state that, for toxic parameters (e.g., ammonia, toxic metals, pesticides), more than one violation of an acute or chronic water quality criterion over a three-year period suggests impairment of aquatic life uses. IDNR has historically used these U.S. EPA guidelines for identifying impairments due to toxic parameters. Based on discussions in 2007 with other states in U.S. EPA Region 7 (i.e., NE and KS) and with U.S. EPA headquarters staff, however, IDNR's approach for identifying impairments due to violations of chronic criteria was changed for the 2008 listing cycle. Impairments due to violations of chronic criteria for toxic parameters were identified for waterbodies where significantly greater than 10 percent of the samples exceed a chronic criterion over a three-year period. Identification of impairments due to violations of acute criteria for toxics remained based on the occurrence of more than one violation of a toxic criterion over a three-year period. This approach will also be used for the 2010 listing cycle.

U.S. EPA (1997, 2002) also developed separate assessment methodologies for using results of fixed station and other ambient monitoring to determine support of fish consumption, primary contact recreation, and drinking water uses. IDNR has largely adopted U.S. EPA's assessment methodology as recommended for primary contact and fish consumption uses (see Table 11). IDNR has modified U.S. EPA's (1997, 2002) Section 305(b) water quality assessment guidelines for assessing drinking water uses with data for nitrate in surface water sources (see Table 11). Also, IDNR has developed assessment methods for toxic data types and assessment categories for which U.S. EPA does not provide specific assessment methods (e.g., using fish kill information).

Chloride, sulfate, and total dissolved solids: Prior to rulemaking efforts by Iowa DNR in 2009, the *Iowa Water Quality Standards* did not contain criteria for protection of aquatic life from either chloride or sulfate. The only related parameter with a numeric criterion was total dissolved solids (TDS): Iowa's general use criteria specified that levels of TDS should not exceed 750 mg/l in any Iowa lake, impoundment, or stream with a flow rate equal to or greater than three times the flow rate of upstream point source discharges. Based on information supplied to IDNR from wastewater permittees, the TDS criterion was changed in 2004 to a site-specific approach: This approach specified an in-stream threshold for TDS of 1,000 mg/l. If a facility's discharge exceeded 1,000 mg/l TDS, toxicity testing would then be required to ensure that the level of TDS being discharged was not toxic to aquatic life. Results of this testing would be used to establish an effluent limit that would be included in the NPDES permit for the facility.

An IDNR rulemaking effort in 2009 resulted in proposed aquatic life criteria for chloride and sulfate (see http://www.iowadnr.gov/water/standards/files/ws_fact.pdf). These new criteria are seen as better indicators of aquatic life health than the previous criterion for TDS which is a measure of all ionic constituents in waters including chloride and sulfate. Although these hardness-dependent criteria were not in effect during the 2006-2008 data gathering period used for the 2010 assessment/listing cycle, IDNR nonetheless compared the existing ambient monitoring data for chloride and sulfate to these criteria. This evaluation was conducted to determine the potential for future Section 303(d) impairments due to chloride and sulfate. For more information on criteria for, see the IDNR's (2009) water quality standards review for chloride, sulfate, and TDS (http://www.iowadnr.gov/water/standards/files/ws_review.pdf).

- **Data from biological monitoring being conducted by IDNR in cooperation with SHL**

Benthic macroinvertebrate and fish sampling data from the IDNR/SHL stream biocriteria and REMAP sampling sites are used to identify impairments of warmwater stream aquatic life uses. IDNR uses a benthic macroinvertebrate index of biotic integrity (BMIBI) and a fish Index of biotic integrity (FIBI) to summarize biological sampling data. The BMIBI and FIBI combine several quantitative measurements or “metrics” that provide a broad assessment of stream biological conditions. A metric is a characteristic of the biological community that can be measured reliably and responds predictably to changes in stream quality. The BMIBI and FIBI each contain twelve metrics that relate to species diversity, relative abundance of sensitive and tolerant organisms, and the proportion of individuals belonging to specific feeding and habitat groups. The metrics are numerically ranked and their scores are totaled to obtain an index rating from 0 (poor) – 100 (optimum).

Qualitative scoring ranges of poor, fair, good, and excellent have been established that reflect the biological community characteristics found at each level. The category of “poor” indicates an impairment of the aquatic life use. The category of “fair,” however, may or may not indicate impairment. A framework for using these data to assess support of aquatic life uses was first developed for Iowa’s 2000 Section 305(b) reporting cycle.

This same basic framework has been used for subsequent reporting/listing cycles.

Several modifications to the process of identifying Section 303(d) biological impairments were made for the 2010 cycle including a more rigorous approach for identifying Section 303(d) biological impairments (see Table 1). A detailed description of these changes and the framework used for the 2010 cycle is included in this methodology as Attachment 2.

- **Data from the IDNR-sponsored lake monitoring conducted by Iowa State University and the University of Iowa Hygienic Laboratory**

The IDNR–sponsored statewide lake water quality monitoring program began in 2000 and continued through 2008. Each of 131 lakes was sampled at least three times during summer seasons to assess seasonal variability of chemical, physical, and biological parameters (e.g., plankton populations). Samples were taken at the deepest point in each lake basin.

Due to year-to-year variability in lake water quality, state limnologists participating in the U.S. EPA Region 7 technical assistance group (RTAG) for nutrient criteria development have recommend that the combined data from at least three years of monitoring results from this type of lake survey is needed to identify nutrient-related water quality

impairments. Thus, IDNR uses median water quality values from a three to five-year period to calculate a trophic state index (TSI) (Carlson 1977). Median-based TSI values are used with the lake assessment framework described in Attachment 3 to determine the existence of an impairment. This framework is based on using the TSI as a numeric translator for Iowa's existing narrative water quality criteria protecting against aesthetically objectionable conditions and/or nuisance aquatic life. For the 2010 reporting/listing cycle, lake data for the five-year period from 2004 through 2008 will be used to identify lake water quality impairments. The 2010 assessment/listing cycle is the fifth such biennial cycle in which the trophic state index has been used to identify Section 303(d) impairments at Iowa lakes.

- **Data from monitoring of bacterial indicators in rivers, lakes, and beach areas**

In July 2003, Iowa DNR adopted criteria for *E. coli* in place of the previous criterion for fecal coliform bacteria into the *Iowa Water Quality Standards* (Table 8). This change was a response to a long-standing recommendation from U.S. EPA. In addition, a proposal was made to subdivide the current Class A (primary contact) use designation to three designations:

- Class A1 (primary contact recreation) (same as the previous Class A designation),
- Class A2 (secondary contact recreational use),
- Class A3 (children's recreational use).

With these revisions, the state of Iowa now considers Class A1 and Class A3 waters with geometric mean levels of *E. coli* greater than 126 organisms per 100 ml to present an unacceptable risk of waterborne disease to swimmers, water skiers, and other persons using surface waters for primary body contact recreational activities where ingestion of water is likely to occur (Section 61.3(3), *Iowa Water Quality Standards*). In addition, Class A2 waters with geometric mean levels of *E. coli* greater than 630 organisms per 100 ml present an unacceptable risk of waterborne disease to persons using surface waters for secondary body contact recreational activities (Section 61.3(3), *Iowa Water Quality Standards*).

In the context of Section 305(b) reporting, U.S. EPA (1997, 2002) recommended that support of primary contact recreation uses be based on (1) a comparison of the geometric mean of at least five samples collected over a 30-day period to state water quality criteria

for indicator bacteria (fecal coliforms, *E. coli*, and/or enterococci) and (2) the percentage of samples that exceed a single-sample maximum value. In cases where the geometric mean exceeded the state water quality criterion, or significantly more than 10% of the samples exceed the single-sample maximum criterion (Table 12), the primary contact uses should be assessed as “impaired.”

Temporal correlation of *E. coli* samples: Several *E. coli* datasets that were reviewed for violations of Iowa’s Class A water quality criteria for the 2010 listing cycle contained *E. coli* data for multiple samples collected on the same day or for samples collected on consecutive days. A recent study of temporal variations in *E. coli* concentrations in the Raccoon River in central Iowa showed a temporal correlation of *E. coli* concentrations within a span of four days (Schilling et al. 2009). Failure to account for this correlation could result in calculations of geometric means that were biased due to inclusion of temporally correlated repeated measures of either high levels or low levels of bacteria in samples collected within this four-day period. Thus, for the 2010 listing cycle, mean (average) values were calculated for multiple *E. coli* samples that collected within a four-day period. This average value was considered an independent estimate of the bacterial concentration during that four-day period, and this average was then used to calculate the geometric mean for the dataset being reviewed.

Identifying bacterial impairments in lakes versus rivers:

Differences in monitoring frequencies at different types of Iowa waterbodies require that different approaches be used when developing assessments of support of contact recreation uses. For example, Iowa’s river reaches and some of its lakes designated for primary contact recreation are typically sampled once per month as part of ambient water quality monitoring activities; none of these river or lake stations are monitored more than twice per month. Thus, a maximum of two samples—and more typically, one sample—of indicator bacteria are collected from these stations during any 30-day period. This amount of data is not sufficient for use with U.S. EPA’s recommended approach for using the geometric mean for assessing support of contact recreation uses (i.e., calculate a geometric mean based on at least five samples collected over a 30-day period).

Swimming beaches at selected Iowa lakes, however, are monitored more frequently than rivers. Beginning in 1999, IDNR began a weekly sampling program during summer months (May through September) at swimming beaches of selected state-owned lakes.

This program now includes weekly sampling at public beaches of approximately 35 state-owned lakes. This program generates the minimum amount of data needed for use with U.S. EPA's recommended assessment method (i.e., five samples collected over a 30-day period).

Thus, because of these varying monitoring frequencies, IDNR uses different procedures to determine the level of use support of the Class A (contact recreation) uses at lake beaches versus river reaches and non-beach areas of lakes. The assessment approaches used for the less frequently monitored rivers and the more frequently monitored lake beaches are discussed below.

Rivers and non-beach areas of lakes: To be assessed as "fully supported" the designated Class A1 primary contact uses, the following conditions should be met: (1) the geometric mean of at least ten samples collected during the recreational seasons (March 15 to November 15) of the current data gathering period (calendar years 2006 through 2008) should not exceed the respective water quality criterion of 126 *E. coli* organisms per 100 ml of *E. coli* and (2) no more than 10 percent of these samples should exceed Iowa's single sample maximum allowable density of 235 *E. coli* organisms per 100 ml. Iowa's 2010 assessment methodology will continue to use the binomial assessment approach for implementing the 10-percent rule that accounts for uncertainty in the use of small sample sizes to identify impairments (see Lin et al. 2000). In addition, no swimming area closures can have been issued during the three-year assessment period. While not entirely consistent with the assessment approach recommended by U.S. EPA (1986, 1997, 2002), the IDNR approach appropriately uses the available monitoring data while incorporating the basic elements of U.S. EPA's recommended approach.

Class A2 (secondary contact recreation) uses will be assessed in a similar manner: (1) the geometric mean of at least ten samples collected during the recreational seasons (March 15 to November 15) of the current data gathering period (calendar years 2006 through 2008) should not exceed the respective water quality criterion of 630 *E. coli* organisms per 100 ml of *E. coli* and (2) no more than 10 percent of these samples, as determined with the binomial method (Lin et al. 2000), should exceed Iowa's Class A2 single sample maximum allowable density of 2,880 *E. coli* organisms per 100 ml.

Lake beaches: Two types of bacteria-related water quality information are used to assess support of the Class A1 (primary contact recreation) use at Iowa's lake beaches: (1) data for indicator bacteria from weekly beach monitoring and (2) information on the closure of beach areas for swimming.

In general, the same approach used for the 2002, 2004, 2006, and 2008 Section 305(b) reporting cycles and Section 303(d) lists will be used for the 2010 cycle: If either of the following exist, the Class A uses of that beach and lake would be assessed as "not supporting" and would be a candidate for Section 303(d) listing:

1. A geometric mean of at least five samples collected over a 30-day period during the recreation season exceeds the Iowa geometric mean water quality criterion of 126 *E. coli* / 100.
2. Significantly more than 10% of the sample values collected during a given recreation season exceed Iowa's single-sample maximum criterion of 235 *E. coli* / 100 ml. Statistical methods developed by Lin et al. (2000) for determining significant exceedances of the 10 percent rule will be used in the process of identifying impairments based on violations of Iowa's single-sample maximum criterion.
3. In the event that a beach was closed to swimming during either year, the Class A1 uses would be assessed as "not supporting." Levels of indicator bacteria that result in IDNR's posting of signs at beaches warning about increased health risk associated with swimming—including both the "Caution: Water Quality Advisory" and the "Water Quality Notice" signs—do not constitute impairment of the Class A1 uses. Neither of these signs are intended to indicate closure of beaches but are posted to warn swimmers of the potential for an increased health risk from swimming. See <http://www.igsb.uiowa.edu/wqm/Beaches/BeachAdvisoryPolicy.htm> for a description of IDNR's beach advisory policy.

For additional information on how IDNR determines support of primary contact and secondary contact recreation uses, see Table 11.

- **Data from programs to monitor fish tissue for toxic contaminants**

The existence of, or potential for, a fish consumption advisory has been, and remains, the basis for Section 305(b) assessments of support of fish consumption uses in Iowa's rivers and lakes. If a waterbody is covered by a consumption advisory, the fish consumption use is assessed as "impaired" (Table 11). Prior to 2006, IDNR used action levels for PCBs, mercury, and chlordane published by the U.S Food and Drug Administration to determine whether consumption advisories should be issued for fish caught as part of recreational fishing in Iowa. In recent years, however, many states have abandoned the use of the FDA action levels in favor of a more protective "risk-based" approach. Thus, in late 2005, the Iowa Department of Public Health (IDPH), in an effort to make Iowa's advisory protocol more protective and more compatible with the various protocols used by adjacent states, developed a revised advisory system for Iowa that covers these contaminants (see Table 13, IDPH (2007) and <http://www.iowadnr.gov/fish/news/consump.html> for more information on Iowa's revised fish consumption advisory protocol).

Other than the changes to a risk-based advisory levels and the addition of a "restricted consumption" category, Iowa's advisory protocol remains the same:

- Decisions to issue consumption advisories remain based on results of annual fish contaminant monitoring conducted either as part of the USEPA/IDNR Regional Ambient Fish Tissue (RAFT) monitoring program or as part of other fish tissue contaminant monitoring programs in Iowa.
- Due to the large amount of variation in contaminant levels within fish populations, two consecutive samplings exhibiting contaminant levels greater than IDNR/IDPH advisory levels are needed to justify issuance of an advisory.
- Similarly, two consecutive samplings showing that contaminant levels are less than the IDNR/IDPH advisory levels are needed to remove an advisory.

In general, these "consecutive" samples are collected in consecutive years as part of the annual U.S. EPA Region VII/IDNR Regional Ambient Fish Tissue (RAFT) monitoring program or as part of special follow-up studies conducted by IDNR. The need to schedule

follow-up samplings one year after the first sampling is related to the length of time required for sample analysis and data reporting. Samples of Iowa fish tissue for RAFT monitoring are typically collected by IDNR biologists in late summer; samples are sent to the U.S. EPA Region VII laboratory in Kansas City for analysis in early fall. Results from this analysis are supplied to IDNR in late spring or early summer of the following year. Decisions to conduct follow-up sampling at a given site are thus based on results of the previous year's sampling. Waterbodies covered by consumption advisories are re-sampled periodically as part of RAFT "follow-up" monitoring to identify any changes in contaminant levels and to justify the need to continue or rescind the advisory.

- **Reports of pollutant-caused fish kills**

Occurrence of a single pollutant-caused fish kill, or a fish kill of unknown origin, on a waterbody or waterbody reach during the most recent three-year period (2007-2009) indicates a severe stress to the aquatic community and suggests that the aquatic life uses should be assessed as "impaired". If a cause of the kill was not identified during the IDNR investigation, or if the kill was attributed to non-pollutant causes (e.g., winterkill), the assessment type will be considered "evaluated." Such assessments, although suitable for Section 305(b) reporting, either are inappropriate for state Section 303(d) listing (no pollutant load to allocate) or lack the degree of confidence to support addition to the state's Section 303(d) list of impaired waters (IR Category 5). Waterbodies affected by such fish kills will be placed in IR subcategories 2b or 3b and will be added to the state list of waters in need of further investigation.

If, however, a cause of the kill is identified, and the cause is either known, or suspected, to be a "pollutant", the assessment type is considered "monitored" and the affected waterbody is a candidate for Section 303(d) listing. Waterbodies affected by this type of kill will be handled as follows:

- TMDLs will not be developed for kills caused by a one-time illegal or unauthorized release of manure or other toxic substance where enforcement actions were taken. The rationale for this approach is as follows:

(1) A consent order has been issued to the party responsible for the kill and monetary restitution sought for the fish killed. A consent order is issued in settlement of an administrative order or as an alternative to issuing an

administrative order. A consent order indicates that IDNR has voluntarily entered into a legally enforceable agreement with the other party. IDNR feels that these enforcement actions are more appropriate, efficient, and effective for addressing a spill-related impairment than is the TMDL process.

(2) No daily load allocation process is possible with a pollutant that is discharged only once.

Such waterbodies will be placed into Integrated Report subcategory 4d as defined by IDNR. In this way, the impairment status of the affected waterbody remains highlighted.

- Fish kills attributed to a pollutant, but where a source of the pollutant was not identified, will be placed into Integrated Report subcategory 5b. The intent of placing these waterbodies into Category 5 is not to necessarily require a TMDL but to keep the impairment highlighted due to the potential for similar future kills from the unaddressed causes and/or sources.
- Fish kills attributed to authorized discharges (i.e., a wastewater discharge meeting permit limits) are considered for Section 303(d) listing (subcategory 5a) as the existing, required pollution control measures are not adequate to address this impairment, and a TMDL is needed.

Note on de-listing waters impaired by fish kills: Iowa DNR historically used the following rationale to de-list fish kill waters:

- If no fish kill has occurred over the last three years, the toxic impact that caused the kill had likely dissipated. Thus, good cause for de-listing exists, and the assessment should be considered “evaluated” (i.e., of lower confidence), and the affected waterbody should be moved from IR Category 5 and be placed into IR Categories 2b or 3b (list of waters in need of further investigation).
- If no fish kill had occurred over the last six years, any impact from the fish kill upon which the impairment was based likely has long-ago dissipated. Thus, the

assessment that was based on the fish kill will not be included in either the new Section 305(b) assessments or Section 303(d) listings. The affected waterbody was moved to IR Category 3a (no information available to assess support of beneficial use).

This rationale was based on (1) acceptance of no more than a once-in-three year acutely toxic water quality impact (U.S. EPA 1994) and (2) conclusions from an IDNR study on recovery of aquatic communities from fish kill events in Iowa (Wilton 2002). Taken together, these two references suggested that a three-year period was more than sufficient an aquatic community to fully recover from a one-time toxic event.

For the 2006 listing cycle, however, U.S. EPA disagreed with IDNR on this de-listing rationale and decided to no longer accept IDNR's de-listing decisions for fish kill waters. U.S. EPA's recommendation for an acceptable de-listing strategy was to conduct biological monitoring at the affected waters to demonstrate that the aquatic community had recovered from the fish kill event. U.S. EPA maintained that without such follow-up field evaluations, IDNR lacked sufficient justification for de-listing a fish kill impairment. Thus, beginning with the 2006 Section 303(d) cycle, the following approach has been used for the de-listing of fish kill impairments in Iowa:

Fish kill-impairments identified on wadeable streams will remain in IR category 5 and on Iowa's Section 303(d) lists until IDNR biological monitoring has been conducted and "full support" of aquatic life uses is indicated (i.e., the "new data" rationale for de-listing impairments). Because monitoring resources are not available to conduct follow-up biological monitoring on most fish kill-affected waters; most IR Category 5 fish kill impairments will remain indefinitely. Also, because IDNR lacks biological assessment protocols for intermittent streams, non-wadeable streams/rivers, and for lakes, the fish kill-related impairments for these waterbody types will remain on Iowa's Section 303(d) list until such assessment protocols are developed and until biological monitoring is conducted in the affected water.

For IR Category 4d waters (i.e., a fish kill-impaired water where enforcement actions were taken against the party responsible for the kill), if no additional fish kills have been reported over at least the last five years, any impact from the fish

kill upon which the impairment was based likely has long-ago dissipated. The IR category for the kill will be changed from 4d to 2b or 3b (potentially impaired) and added to the state list of waters in need of further investigation. If no additional kills have been reported for an additional five year period, the IR category will be changed from 2b/3b to 3a (water not assessed).

If no additional fish kills have been reported for a 2b/3b water over the last ten years, the IR category for the kill will be changed from 2b/3b (potentially impaired) to 3a (insufficient information to assess support of use).

Iowa DNR's 2010 listing/de-listing methodology for fish kills is summarized in Table 14.

- **Data from the statewide survey of freshwater mussels from 1998-1999**

Information from *Statewide Assessment of Freshwater Mussels (Bivalva: Unionidae) in Iowa Streams: Final Report* (Arbuckle et al. 2000) will again be used for the 2010 IR to assess support of aquatic life uses of Iowa streams and rivers. Only a few localized mussel surveys have been conducted since the statewide survey of Arbuckle et al. (2000). Thus, unless special studies produce additional mussel survey data, the majority of the assessments based on Arbuckle et al. (2000) will not change for the 2010 assessment and listing cycle. The methodology used to develop these assessments is as follows.

The survey conducted by Arbuckle et al. (2000) involved re-sampling of sites visited in the mid-1980s by Frest (1987). For purposes of identifying candidates for Section 303(d) listing, the number of mussel species reported for a given waterbody by Frest was compared to the number of species reported for the same waterbody by Arbuckle et al. The degree to which the aquatic life use was supported was based on the percent change in the number of mussel species from the 1984-85 period to the 1998-99 period. If the mean waterbody species richness (SR) was four or greater in the 1984-1985 survey period, then the following assessment approach using percent change from the 1984-85 to 1998-99 survey periods was used to identify candidates for Section 303(d) listing:

If species richness (SR) in 1984-85 is ≥ 4 , and the percent decline in SR from 1984-85 to 1998-99 is:	Then use support category is:	Integrated Report Category
< 25%	Fully Supporting	1

26-50%	Fully Supporting <u>or</u> Fully Supporting / Threatened with a declining trend (potentially "impaired")	1 or 5b
51%-75%	Partially Supporting ("impaired")	5b
> 75%	Not Supporting ("impaired")	5b

The decision to consider only those sites having four or more species reported in the 1984-85 survey is based on (1) a review of the historical distributions of freshwater mussels in Iowa as shown by Cummings and Mayer (1992) and (2) the framework (i.e., percent decline approach) described in table above. For the Iowa ecoregions that show historical presence of a stream/river community of freshwater mussels (i.e., all ecoregions except 47e and the portions of ecoregions 47f and 40 in the Missouri River drainage), a species richness of approximately four appears to characterize average species richness from the 1984-85 survey by Frest. The decision to identify a waterbody as impaired due to a decline in species richness between the 1984-85 and 1998-99 survey periods is based on quartiles (i.e., from a 25% to 50% decline: "fully supported/threatened with a declining trend"; from a 50% to 75% decline, "partially supported"; more than a 75% decline, "not supported"). Any decision to add a waterbody to the state list of impaired waters based on a percent decline of between 26 and 50 percent will be made on a case-by-case basis, with impairment and listing more likely as the percent decline approaches 50 percent. Using four species as a minimum for this assessment approach allows for some apparent decline between the survey periods without identifying the waterbody as "impaired." Such declines may be due to problems with sampling efficiency as opposed to the actual elimination of species.

As presented by Arbuckle et al. (2000), the potential causes of declines in species richness of Iowa's freshwater mussels include siltation, destabilization of stream substrate, stream flow instability, and high in-stream levels of nutrients (phosphorus and nitrogen). Their study also suggested the importance of stream shading provided by riparian vegetation to mussel species richness. For purposes of Section 305(b) reporting and Section 303(d) listing, the following causes and sources will be identified for all waters assessed as "impaired" due to declines in the mussel community: siltation from agricultural and natural sources; flow modification due to hydromodification of the watershed; and nutrients from agricultural and natural sources. Because site-specific causes and sources of these impairments were not identified, any waters assessed as impaired due to declines in the freshwater mussel community will be placed into

subcategory 5b. As is typical for Section 305(b) water quality assessments, the sources of impairment identified for Iowa's freshwater mussel community are only *potential* sources. The logistics of a statewide water quality assessment process does not often allow precise site-specific determinations of pollutant sources. More accurate information on sources would typically be gathered during the stressor identification phase of TMDL development.

- **Data from public water supplies on the quality of raw and finished water**

Data for the quality of raw (untreated) water from a surface water source will be used with the methodology for identifying impairments in Class C (drinking water use) waters described in Table 11. Three types of contaminants are considered as part of Section 305(b) assessments to determine the degree to which the designated Class C uses are supported: metals, pesticides, and inorganics (nitrate). Impairment of Class C uses for these classes of toxic contaminants will be determined as follows:

Data for metals or pesticides in the raw water source:

For sampling frequencies of quarterly or more frequent, moving annual average values will be compared to the respective Class C criterion (see Table 7). The type of annual averages calculated will depend on the sampling frequency for the data collected (e.g., monthly vs. quarterly). If any moving annual average exceeds the Class C criterion, the Class C uses will be assessed as impaired (not supported). When calculating moving annual averages, non-detect values will be set equal to the IDNR ambient monitoring non-detect level. Situations where non-detect levels exceed water quality criteria will be handled on a case-by-case basis.

Data for inorganics (i.e., nitrate) in the raw water source:

No more than 10 percent of the samples violate the maximum contaminant level (MCL) for nitrate. The methodology of Lin et al. (2000) will be used to determine whether significantly more than 10 percent of the samples exceed the MCL.

Impairments related to the quality of finished (treated) water will be determined through review of annual IDNR public drinking water program compliance reports (e.g., IDNR/WQB 2007, 2008, 2009) available at <http://www.iowadnr.gov/water/drinking/reports.html>). Information from these reports on violations of Class C water quality criteria and issuance of drinking water advisories will be

used with methods described in Table 11 to determine the existence of impairment of drinking water uses.

- **Data from special studies of water quality and aquatic communities**

Results of special water quality studies that meet all requirements of Iowa's "credible data" law, including the availability of a quality assurance project plan (or equivalent plan or methodology for sampling and analysis), will be considered on a case-by-case basis. IDNR will review all relevant quality assurance/project plans for special studies prior to the decision to use study results for purposes of Section 303(d) listing. Results from special studies that meet "credible data" requirements will be compared to water quality criteria as specified in the *Iowa Water Quality Standards* with the methods described in this document.

- **Results of volunteer monitoring that meet "credible data" requirements**

Results of volunteer monitoring that meet all requirements of Iowa's "credible data" law, including the availability of a DNR-approved quality assurance project plan (or equivalent plan or methodology for sampling and analysis), will be considered on a case-by-case basis. IDNR will review all relevant quality assurance/project plans for volunteer monitoring studies prior to the decision to use study results for purposes of Section 303(d) listing. Results from volunteer monitoring studies that meet "credible data" requirements will be compared to the appropriate water quality criteria as specified in the *Iowa Water Quality Standards* with the methods described in this document.

Exclusion (de-listing) of waters from the 2008 Section 303(d) list:

According to U.S. EPA regulations (40 CFR 130.7), a state must demonstrate "good cause" for exclusion of previously impaired waterbodies. According to these regulations, "good cause" includes, but is not limited to, more recent or accurate data; more sophisticated water quality modeling; flaws in the original analysis that led to the water being listed; or changes in conditions; e.g., new control equipment or the elimination of discharges. Thus, the following can be used to demonstrate good cause for not listing a waterbody on the Section 303(d) list or to decrease the scope of impairment to a listed waterbody:

- **More recent or accurate data.** Additional monitoring data or information from a waterbody may demonstrate that it now meets applicable water quality standards. These data must be generated from monitoring studies and programs consistent with Iowa's credible data law and

must be in sufficient quantity to be used with Section 305(b) water quality assessment procedures (see Table 6). Special conditions include the following:

1. Chlorophyll-a and Secchi depth: For Iowa lakes, median-based trophic state index (TSI) values for both chlorophyll-a and Secchi depth must be 63 or less for two consecutive summer seasons before a lake can be removed from the state's Section 303(d) list (IR Category 5) (see Attachment 3 of this methodology for more information). A TSI value of 63 indicates a chlorophyll-a concentration of approximately 27 ug/l and a Secchi depth of approximately 0.8 meters.
 2. Indicator bacteria: For waters with contact recreation uses assessed as impaired by indicator bacteria—and assuming that sufficient and credible new data are available—geometric mean levels must be less than the applicable state water quality criterion for two consecutive listing cycles (i.e., five consecutive years) prior to de-listing. Also, the percentage of samples that exceed the state's single-sample maximum E. coli criterion must be less than 10 percent for two consecutive listing cycles. Requiring that geometric means and single-sample maximum values meet applicable water quality criteria for two consecutive listing cycles is designed to avoid impairment flip-flopping that can occur with high-variability and weather-influenced parameters such as indicator bacteria.
 3. Atrazine: For waters with drinking water uses assessed as impaired by atrazine, all moving annual averages must be less than the atrazine MCL for two consecutive Section 303(d) listing cycles before a de-listing due to more recent data. Atrazine in surface waters, and especially in lakes, can exhibit wide fluctuation from year to year. IDNR assessment/listing staff will review the historic atrazine data to determine any trends in levels and to determine whether de-listing is justified.
- **Flaws in original analysis or errors in listing**. Errors in the data or flaws in assessment procedures used to list the waterbody invalidate the basis for listing. Changes in assessment methodology can be considered as correcting flaws in analysis or errors in listing.
 - **New conditions**. Examples of new conditions include revised water quality standards, the elimination of discharges, and new control equipment such that a listed waterbody no longer meets the criteria for Section 303(d) listing.

All waters removed from Iowa's 2008 Section 303(d) list will be summarized in a table posted at the Iowa DNR impaired waters web site (<http://www.igsb.uiowa.edu/wqm/ImpairedWaters/303d.html>). For any waterbody listed on the 2008 Section 303(d) list and not included on the 2010 list, a waterbody-specific rationale for the exclusion or de-listing will be incorporated into Iowa DNR's on-line Section 305(b) assessment database (ADBNet: <http://programs.iowadnr.gov/adbnet/index.aspx>).

Waterbodies added to an Iowa 303(d) list will be placed on subsequent lists unless (1) there are sufficient credible data to reassess the waterbody and demonstrate that 303(d) listing is not appropriate or (2) some other "good cause" is demonstrated for not including the water on the 303(d) list. Age of data alone is not an adequate justification for not including a previously-listed water on a new list of impaired waters. This provision is especially relevant to waterbodies included on lists based on results of one-time surveys (e.g., results of biological assessments conducted as part of biocriteria development or faunal surveys (e.g., freshwater mussels)). For example, if a waterbody was added to Iowa's 2004 303(d) list based on a biological assessment conducted in 2002, this waterbody should remain on Iowa's subsequent 303(d) lists until (1) a TMDL is completed, (2) additional monitoring is conducted that shows "full support" of aquatic life uses, or (3) a flaw in the original data analysis or assessment is discovered.

In addition, lack of sufficient new data to develop a "monitored" assessment for a previously-listed waterbody is not adequate justification for excluding a waterbody from Section 303(d) listing. For example, if a routinely-monitored waterbody was added to Iowa's 2004 303(d) list based on a "monitored" assessment showing violations of the Iowa water quality criterion for indicator bacteria, this waterbody should remain on Iowa's impaired waters lists until (1) adequate data are available to develop a high-confidence ("monitored") assessment, (2) the newly developed assessment shows "full support" of the impaired use, or (3) there is some other "good cause" for de-listing this impairment.

Prioritization and scheduling of waters for TMDL development:

CWA Section 303(d) requires that each "state shall establish a priority ranking for such waters, taking into account the severity of the pollution and the uses to be made of such waters." A system of prioritization for waterbodies included in Category 5 of the Integrated Report has been developed by the IDNR based on several factors. Included in these factors are the required elements of "the severity of the pollution and the uses to be made of such waters." The methods developed are described below; these methods are the same as used to prioritize waterbodies on Iowa's 2002 through 2008 Section 303(d) lists. These criteria are a guide. Other factors, such as best professional judgment of IDNR staff, results of volunteer monitoring, and public comments, may also be considered when prioritizing waters.

If a waterbody meets any one criterion in a priority category, that does not necessarily mean the water will be prioritized as such, since many waters fit some criteria from all categories.

Priorities	Applicable Criteria
-------------------	----------------------------

High	
-------------	--

- | | |
|--|---|
| | <ul style="list-style-type: none">▪ Waters where sufficient water quality information exists to understand and analyze causes and effects of the problems and opportunities are available to correct or substantially improve water quality;▪ Waters with imminent human health or aquatic health problems;▪ Waters with documented widespread local support for water quality improvement; or▪ Waters where state or federally threatened or endangered species are impacted. |
|--|---|

Medium	
---------------	--

- | | |
|--|--|
| | <ul style="list-style-type: none">▪ Waters where sufficient water quality information exists to understand and analyze causes and effects of the problems; however, opportunities are not immediately available to correct or substantially improve water quality; or▪ Waters where local support for TMDL development is expected but not known. |
|--|--|

Low	
------------	--

- | | |
|--|--|
| | <ul style="list-style-type: none">▪ Waters where insufficient water quality information exists to understand and analyze causes and effects of the problems and limited opportunities are available, at this time, to correct or substantially improve water quality;▪ Waters with no evident local support for water quality improvements. |
|--|--|

Addressing interstate inconsistencies in Section 303(d) lists:

Inconsistency in the Section 303(d) listings of border rivers and other interstate waters is a long-standing national problem (see GAO 2002). IDNR faces potential listing consistency issues with the following states and rivers that border Iowa: South Dakota (Big Sioux River), Nebraska (Missouri River), Missouri (Des Moines River), and Illinois and Wisconsin (Upper Mississippi River). Thus, IDNR will either (1) request and/or review the draft 303(d) lists of, or (2) consult directly with, states with which Iowa shares border waters.

The Upper Mississippi River Basin Association's *Water Quality Task Force* has provided, and continues to provide, a forum for improving listing consistency for the Upper Mississippi River for the states of Illinois, Iowa, Minnesota, Missouri and Wisconsin (see UMRBA-WQTF 2004). In addition to the face-to-face consultations provided in the UMRBA Water Quality Task Force, interstate consistency can also be

addressed through viewing web-available integrated reports and Section 303(d) lists of adjacent states. For the 2010 listing cycle, integrated reporting web sites for Nebraska and South Dakota were visited to, as much as possible, resolve interstate listing issues:

- Nebraska waterbodies of relevance to interstate coordination are (1) the Missouri River downstream from its confluence with the Platte River (NDEQ waterbody MT1-10000) and (2) the Missouri River from the Platte River upstream to its confluence with the Big Sioux River (NDEQ waterbody NE1-10000) (see NDEQ 2010 and <http://www.deq.state.ne.us/>).
- The South Dakota waterbodies of the Big Sioux River of relevance to interstate coordination are as follows (see SDENR 2010 and <http://denr.sd.gov/des/sw/surfacewaterquality.aspx>):
 - SD BS-R-Big-Sioux_17: mouth to Indian Creek
 - SD BS-R-Big-Sioux_16: Indian Creek to near Alcester
 - SD BS-R-Big-Sioux_15: near Alcester to Fairview
 - SD BS-R-Big-Sioux_14: near Fairview to Ninemile Creek
 - SD BS-R-Big-Sioux_13: Ninemile Creek to near Brandon (partial: to IA/MN state line)

Where the listing in another state is different than in Iowa, the IDNR will review the assessment data, supporting information, and assessment methodology that support the listing in the other state. These data will be reviewed and applied to Iowa's Section 303(d) listing methodology outlined in this document. If a listing from another state for a border river is based on water quality standards that are consistent with the *Iowa Water Quality Standards*, the Iowa listing will be changed to reflect that listing.

IDNR will also review the Section 303(d) listings from adjacent states for waters that either enter Iowa from Minnesota or leave Iowa into Minnesota or Missouri (e.g., the Cedar River in Mitchell County and the Chariton River in Appanoose County), or that are shared with Iowa by either state (e.g., Tuttle Lake in Emmet County). In terms of waters flowing into the state of Iowa from the state of Minnesota, the following basin assessments were reviewed at the Minnesota Pollution Control Agency's 305(b) assessment [web site](#): Cedar, Des Moines, Minnesota, and Missouri. In terms of waters flowing from the state of Iowa into the state of Missouri, the Missouri DNR's web site for impaired waters (<http://www.dnr.mo.gov/env/wpp/waterquality/303d.htm>) was reviewed for impairments that might affect Iowa's impairment decisions.

Where Section 303(d) listing decisions differ across a state line, the supporting assessment data and methodology will be requested from the appropriate state. IDNR will review these data using Iowa's Section 303(d) listing methodology outlined in this document to determine whether modifications to Iowa's Section 303(d) list are justified.

This process of reviewing Section 303(d) listings for waters that border or are shared with adjacent states is designed to reduce between-state inconsistencies in Section 303(d) listings and to provide a basis for cooperation on future development of TMDLs for these interstate waters.

Public participation:

A draft of this methodology was provided to the public for review and comment as part of the public comment period for the draft 2010 Section 303(d) list. The draft methodology was available in hard copy by contacting the IDNR. The draft was also available at the IDNR website at <http://www.igsb.uiowa.edu/wqm/wqa/303d.html>. Comments on the draft methodology were received for a period of thirty days.

The methods used to assess water quality, however, are always changing, due both to recommendations from U.S. EPA and due to changes at the state level (e.g., changes in the *Iowa Water Quality Standards*). Thus, IDNR will accept comments at any time regarding this methodology.

References

- Arbuckle, K.E., J.A. Downing, and D. Bonneau. 2000. Statewide assessment of freshwater mussels (Bivalva: Unionidae) in Iowa streams: final report. Iowa Department of Natural Resources.
- Carlson, R.E. 1977. A trophic state index for lakes. *Limnol Oceanogr.* 22:361-369.
- Carlson, R.E. 1991. Expanding the trophic state concept to identify non-nutrient limited lakes and reservoirs. In: *Proceedings of a National Conference on Enhancing the States' Lake Management Programs, Monitoring and Lake Impact Assessment*, Chicago. pp. 59-71.
- Cummings, K.S. and C.A. Mayer. 1992. *Field Guide to Freshwater Mussels of the Midwest. Manual 5*, Illinois Natural History Survey. 194 p.
- Foreman, K.L. 2007. Water quality monitoring in the Hoover Creek watershed, 2004-2006. Iowa Geological Survey, Technical Information Series 53. Iowa Department of Natural Resources. 50 p.
- Frest, T.J. 1987. Mussel survey of selected interior Iowa streams. University of Northern Iowa. Final Report to Iowa Department of Natural Resources and U.S. Fish & Wildlife Service.
- GAO. 2002. Water quality: inconsistent state approaches complicate Nation's efforts to identify its most polluted waters. Report GAO-02-186. United States General Accounting Office. 40 pp.
- IAC. 2008. Chapter 567-61: water quality standards. Iowa Administrative Code [effective date 06/11/2008].
- IDNR. 2009. Water quality standards review: chloride, sulfate, and total dissolved solids. Iowa Department of Natural Resources. 79 p.
- IDNR. 2010. Iowa's beach monitoring program, 2008. Water Fact Sheet 2010-2. Watershed Monitoring and Assessment Section, Iowa Department of Natural Resources. 4 p.
- IDNR/WQB. 2007. State of Iowa, public drinking water program: 2006 annual compliance report. Water Supply Engineering & Operations sections, Water Quality Bureau, Environmental Services Division, Iowa Department of Natural Resources. 29 p. + appendices.
- IDNR/WQB. 2008. State of Iowa, public drinking water program: 2007 annual compliance report. Water Supply Engineering & Operations sections, Water Quality Bureau, Environmental Services Division, Iowa Department of Natural Resources. 30 p. + appendices.
- IDNR/WQB. 2009. State of Iowa, public drinking water program: 2008 annual compliance report. Water Supply Engineering & Operations sections, Water Quality Bureau, Environmental Services Division, Iowa Department of Natural Resources. 32 p. + appendices.
- IDNR/WRS. 2001. Stream watershed assessment to support development of total maximum daily loads (TMDLs): project report. Water Resources Section, Environmental Protection Division, Iowa Department of Natural Resources. 86 p.

- IDPH. 2007. Fish consumption advisory protocol in Iowa. Iowa Department of Public Health. 8 p.
- Langel, R.J. and M.M. Kilgore. 2006. Water quality monitoring in the Yellow River watershed, 2005. Iowa Geological Survey, Technical Information Series 50. Iowa Department of Natural Resources. 85 p.
- Lin, P., D. Meeter, and X. Niu. 2000. A nonparametric procedure for listing and delisting impaired waters based on criterion exceedances. Technical Report, prepared by Department of Statistics, Florida State University, submitted to the Florida Department of Environmental Protection. 21 pgs.
- Littin, G.R. and J.C. McVay. 2008. Water-quality and biological assessment of the Iowa River and tributaries within and contiguous to the Meskwaki Settlement of the Sac and Fox Tribe of the Mississippi in Iowa, 2006–07: U.S. Geological Survey Scientific Investigation Report, 2009–5105, 41 p.
- Lutz, D.S. and B.C. Francois. 2007. Annual report: water quality studies—Red Rock and Saylorville reservoirs, Des Moines River, Iowa: January 10, 2006–December 5, 2006. Submitted to Department of the Army, Rock Island District, Corps of Engineers, Rock Island, IL. Environmental Engineering Section, Department of Civil Engineering, Engineering Research Institute, Iowa State University, Ames. 395 p.
- Lutz, D.S. and B.C. Francois. 2008. Annual report: water quality studies—Red Rock and Saylorville reservoirs, Des Moines River, Iowa: January 9, 2007–December 4, 2007. Submitted to Department of the Army, Rock Island District, Corps of Engineers, Rock Island, IL. Environmental Engineering Section, Department of Civil Engineering, Engineering Research Institute, Iowa State University, Ames. 398 p.
- Lutz, D.S. and C.J. Steffen. 2009. Annual report: water quality studies—Red Rock and Saylorville reservoirs, Des Moines River, Iowa: January 15, 2008–December 4, 2008. Submitted to Department of the Army, Rock Island District, Corps of Engineers, Rock Island, IL. Environmental Engineering Section, Department of Civil Engineering, Engineering Research Institute, Iowa State University, Ames. 393 p.
- MPCA. 2010. 2010 integrated report: general report to the Congress of the United States pursuant to Section 305(b) of the 1972 Clean Water Act: water years 2008-2009. Minnesota Pollution Control Agency. Accessed at <http://www.pca.state.mn.us/index.php/view-document.html?qid=5968> on April 7, 2011.
- NDEQ. 2006. Methodologies for waterbody assessments and development of the 2006 integrated report for Nebraska. Nebraska Department of Environmental Quality, Water Quality Division. 21 pgs. plus appendix.
- NDEQ. 2010. 2010 water quality integrated report. Water Quality Division, Nebraska Department of Environmental Quality. Accessed at <http://www.deq.state.ne.us/> on April 7, 2011.
- Schilling, K.E., T. Hubbard, J. Luzier, and J. Spooner. 2006. Walnut Creek watershed restoration and water quality monitoring project: final report. Iowa Geological Survey, Technical Information Series 49. Iowa Department of Natural Resources. 124 p.
- Schilling, K.E., Y-K. Zhang, D.R. Hill, C.S. Jones, C.F. Wolter. 2009. Temporal variations of *Escherichia coli* concentrations in a large Midwestern river. *Journal of Hydrology*: 365:79-85.

- SDENR. The 2010 South Dakota integrated report for surface water quality assessment. South Dakota Department of Environment and Natural Resources. Accessed at <http://denr.sd.gov/documents/10irfinal.pdf> on April 7, 2011.
- Smith, E.P., K. Ye, C. Hughes, and L. Shabman. 2001. Statistical assessment of violations of water quality standards under Section 303(d) of the Clean Water Act. *Environmental Science & Technology* 35:606-612.
- Tomer, M.D., T.B. Moorman, and C.G. Rossi. 2008. Assessment of the Iowa River's South Fork watershed: Part 1. water quality. *Journal of Soil and Water Conservation*, 63(6):360-370.
- UMRBA-WQTF. 2004. Upper Mississippi River water quality: the states' approaches to Clean Water Act monitoring, assessment, and impairment decisions. Upper Mississippi River Basin Association, Water Quality Task Force. 75 p.
- U.S. EPA. 1986. Quality criteria for water, 1986. Office of Water Regulations and Standards, U.S. Environmental Protection Agency, Washington D.C., EPA 440/5-86-001. 451 p.
- U.S. EPA. 1994. Water quality standards handbook: second edition. U.S. Environmental Protection Agency, EPA 823-B-94-005a.
- U.S. EPA. 1997. Guidelines for the preparation of the comprehensive state water quality assessments (305(b) reports) and electronic updates. Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency, Washington, D.C.
- U.S. EPA. 2002. Consolidated assessment and listing methodology: toward a compendium of best practices. First edition, July 2002. U.S. Environmental Protection Agency.
- U.S. EPA. 2003. Guidance for 2004 assessment, listing, and reporting requirements pursuant to Sections 303(d) and 305(b) of the Clean Water Act, July 21, 2003. Watershed Branch, Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency. 32 p.
- U.S. EPA. 2005. Guidance for 2006 assessment, listing, and reporting requirements pursuant to Sections 303(d), 305(b), and 314 of the Clean Water Act, July 29, 2005. Watershed Branch, Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency. 70 p. plus appendices. (<http://www.epa.gov/owow/tmdl/2006IRG/>)
- U.S. EPA. 2006. Information concerning 2008 Clean Water Act Section 303(d), 305(b), and 314 integrated reporting and listing decisions. Memorandum of October 12, 2006 from Diane Regas, Director, Office of Wetlands, Oceans and Watersheds to Water Directors of Regions 1-10. 17 p. (http://www.epa.gov/owow/tmdl/2008_ir_memorandum.html)
- U.S. EPA. 2009. Information concerning 2010 Clean Water Act Sections 303(d), 305(b), and 314 integrated reporting and listing decisions. Memorandum of May 5, 2009 from Suzanne Schwartz, Acting Director, Office of Wetlands, Oceans and Watersheds to Water Directors of Regions 1-10. 8 p plus appendix (<http://www.epa.gov/owow/tmdl/guidance/final52009.pdf>).
- Wilton, T. 2002. Stream fish kill follow-up assessment: fish community sampling results. Water

Resources Section, Environmental Protection Division, Iowa Department of Natural Resources.
21 p.

Table 1. Summary of changes in Iowa DNR's Section 303(d) listing methodology between the 2008 and 2010 listing cycles.		
Change in Methodology:	2008 Listing Cycle	2010 Listing Cycle
Assigning magnitudes of bacterial impairments	Considered single-sample maximum impairments as "partially supported"; those based on geometric means as "not supported"	Magnitudes are assigned based on magnitude of geometric mean
Revision of biological assessment and listing protocol	One sampling of a stream of watershed size within calibration range was sufficient to identify biological impairment	Two independent samplings are needed to identify biological impairment
Creation of IR subcategories to better track biological assessments on streams/rivers having watershed sizes both within and outside of the calibration range of IDNR's biological assessment protocol.	No distinction between assessments conducted within or outside of bioassessment calibration range.	Addition of six IR subcategory suffixes for calibrated and un-calibrated biological assessments for IR subcategories 2b and 3b, and creation of a 5b subcategory for waters needing additional sampling
Changes in water quality criteria for toxics (e.g., metals and pesticides)	Criteria for toxics as approved in the <i>Iowa Water Quality Standards</i> prior to 2009 were used to identify impairments	Revised criteria in the <i>Iowa Water Quality Standards</i> as approved by U.S. EPA in 2009 were used to identify impairments

Table 2. Summary of U.S. EPA's "integrated reporting" format as used for Iowa's 2010 Section 305(b) and Section 303(d) cycle.		
Integrated Report Category	Source of Category	Description of Category
1	U.S. EPA	All designated uses are met.
2a	U.S. EPA	Some of the designated uses are met but there is insufficient data to determine if remaining designated uses are met.
2b	IDNR	At least one use assessed as supported with at least one other use potentially impaired based on an "evaluated" assessment. This subcategory, along with subcategory 3b, forms the state list of waters in need of further investigation.
2b-c	IDNR	Potential biological impairment on stream with watershed size within calibration range of assessment protocol. At least one other use is assessed as "fully supported," but the aquatic life use of a stream segment with a watershed size within the calibrated range of IDNR's biological assessment protocol has been assessed as potentially impaired.
2b-u	IDNR	Potential biological impairment on stream with watershed size outside of calibration range of assessment protocol. At least one other use is assessed as "fully supported," but the aquatic life use of a stream segment with a watershed size outside the calibrated range of the biological assessment protocol has been assessed as potentially impaired.
3a	U.S. EPA	Insufficient data to determine whether any designated uses are met.
3b	IDNR	Insufficient data exist to determine whether any designated uses are met, but at least one use is potentially impaired based on an "evaluated" assessment. This subcategory, along with subcategory 2b, forms the state list of waters in need of further investigation.
3b-c	IDNR	Potential biological impairment on stream with watershed size within calibration range of assessment protocol. The aquatic life use of a stream segment within the calibrated range of the biological assessment protocol has been assessed as potentially impaired; no other uses are assessed due to lack of water quality information;

Table 2. Summary of U.S. EPA's "integrated reporting" format as used for Iowa's 2010 Section 305(b) and Section 303(d) cycle.		
Integrated Report Category	Source of Category	Description of Category
<i>Continued:</i>		
3b-u	IDNR	Potential biological impairment on stream with watershed size outside of calibration range of assessment protocol. The aquatic life use of a stream segment with a watershed size outside the calibrated range of the biological assessment protocol has been assessed as potentially impaired; no other uses are assessed due to lack of water quality information;
4a	U.S. EPA	Water is assessed as impaired or threatened but a TMDL is not needed because a TMDL has been completed.
4b	U.S. EPA	Water is assessed as impaired but a TMDL is not needed because other required control measures are expected to result in attainment of water quality standards in a reasonable period of time.
4c	U.S. EPA	Water is assessed as impaired but a TMDL is not needed because the impairment or threat is not caused by a "pollutant."
4d	IDNR	Water is assessed as impaired due to a pollutant-caused fish kill but a TMDL is not needed because enforcement actions were taken against, and monetary restitution sought from, the party responsible for the kill.
5a	U.S. EPA	Water is assessed as impaired or threatened by a pollutant stressor and a TMDL is needed [along with IR categories 5b and 5p, the state's Section 303(d) list].
5b	IDNR	Water is assessed as impaired or threatened based on results of biological monitoring or a fish kill investigation where specific causes and/or sources of the impairment have not yet been identified [along with IR categories 5a and 5p, the state's Section 303(d) list].
5b-t	IDNR	Tentative biological impairment: The aquatic life uses of a stream segment with a watershed size within the calibration range of the IDNR biological assessment protocol are assessed as Section 303(d)-impaired based on <u>only one</u> of the two biological sampling events needed to confirm the existence of a biological impairment.
5b-v	IDNR	Verified biological impairment: The aquatic life uses of a stream with a watershed size within the calibration range of IDNR biological assessment protocol are assessed as Section 303(d)-impaired based on results of the required two or more biological sampling events in multiple years within the previous five years needed to confirm the existence of a biological impairment.
5p	IDNR	A presumptively-applied designated use is assessed as 303(d) impaired or threatened. [Along with IR categories 5a and 5b, the state's Section 303(d) list.]

Table 3. Comparison of Iowa DNR's assessment reaches for the Upper Mississippi River to those agreed upon in 2004 by the Upper Mississippi River Basin Association (UMRBA) as part of the memorandum of understanding on interstate assessment reaches developed by the UMRBA Water Quality Task Force.

IDNR Waterbody ID Number	Waterbody Description	Length (miles)	UMRBA Assessment Reach	Segment Description	Length (miles)*	Hydrologic Unit Code (HUC)
IA 03-SKM-0010-1	Iowa/Missouri state line (Des Moines R.) to Sugar Cr. nr. Ft. Madison	17.3	Flint-Henderson	Des Moines R. to Iowa R.	74.75	07080104
IA 03-SKM-0010-2	Sugar Cr. to Skunk R.	19.5				
IA 02-ICM-0010-1	Skunk R. to water supply intake at Burlington	8.75				
IA 02-ICM-0010-2	Burlington water supply intake to Iowa R.	29.2				
IA 01-NEM-0010-1	Iowa R. to L&D 15 at Davenport	49.3	Copperas-Duck	Iowa R. to Lock & Dam 13 at Clinton	89.3	07080101
IA 01-NEM-0010-2	L&D 15 to L&D 14 at LeClaire	10.7				
IA 01-NEM-0010-3	L&D 14 to Wapsipinicon R.	13.1				
IA 01-NEM-0010-4	Wapsipinicon R. to L&D 13 at Clinton	16.2				
IA 01-NEM-0020-1	L&D 13 to Catfish Cr. at Dubuque	54.0	Apple-Plum	Lock & Dam 13 to Lock & Dam 11	59.68	07060005
IA 01-NEM-0020-2	Catfish Cr. to L&D 11 at Dubuque	5.68				
IA 01-NEM-0030-1	L&D 11 to L&D 10 at Guttenberg	30.9	Grant-Maquoketa	Lock & Dam 11 to Wisconsin R.	46.0	07060003
IA 01-NEM-0030-2	L&D 10 to Wisconsin R.	15.1				
IA 01-NEM-0040-1	Wisconsin R. to L&D 9 at Harpers Ferry	19.0	Coon-Yellow	Wisconsin R. to Root R.	42.9	07060001
IA 01-NEM-0040-2	L&D 9 to IA/MN state line	23.9				

*The length of the UMRBA assessment reaches was adjusted to correspond to the total mileage in the respective IDNR assessment reaches.

Table 4. Monitoring stations on the Iowa portion of the Upper Mississippi River and associated tributaries sampled from 2004 through 2008 as part of the USGS Long-Term Resource Monitoring Program (LTRMP)						
No.	Waterbody, Location	Designated Uses**	Waterbody ID Number	County	Dates of First and Last Sampling	LTRMP Station No.
1.	Catfish Cr., near mouth,	A1,B(WW1)	IA 01-TRK-0100_1	Dubuque	Mar. 25, 1998 to Sep. 21, 2004	CF00.3M
2.	Elk R., near mouth	A1,B(WW1)	IA 01-MAQ-0030_1	Clinton	Sep. 20, 1997 to Sep. 20, 2004	ER02.4M
3.	Maquoketa R., near mouth	A1,B(WW1)	IA 01-MAQ-0050_1	Jackson	May 5, 1993 to Nov. 12, 2008	MQ02.1M
4.	Mill Cr. near mouth	A1,B(WW2)	IA 01-TRK-0030_1	Jackson	Mar. 26, 1998 to Sep. 20, 2004	MC01.0M
5.	Upper Mississippi R. at Le Claire	A1,B(WW1)	IA 01-NEM-0010_2	Scott	May 19, 1993 to Sep. 22, 2004	M497.2B
6.	Upper Mississippi R. at Camanche	A1,B(WW1)	IA 01-NEM-0010_4	Clinton	May 5, 1993 to Sep. 22, 2004	M511.4B
7.	Upper Mississippi R. L&D 12 tailwater, Bellevue	A1,B(WW1)	IA 01-NEM-0020_1	Jackson	Oct. 15, 1990 to Nov. 10, 2008	M556.4A
8.	Upper Mississippi R, L&D 11 tailwater, Dubuque	A1,B(WW1)	IA 01-NEM-0020_2	Dubuque	May 6, 1993 to Sep. 21, 2004	M582.5B
9.	Upper Mississippi R, L&D 10 tailwater, Guttenberg	A1,B(WW1)	IA 01-NEM-0030_2	Clayton	Jun. 22, 1998 to Sep. 21, 2004	M615.2B
10.	Upper Mississippi River at Gordon's Bay Landing	A1,B(WW1)	A 01-NEM-0040_1	Allamakee	Sep. 21, 2001 to Jun. 16, 2006	M646.9X
11.	Upper Mississippi R. Big Slough at Lansing Bridge	A1,B(WW1)	IA 01-NEM-0040_2	Allamakee	Apr. 19, 1996 to Jun. 16, 2006	M663.4E
12.	Rock Cr., near mouth	A1,B(WW2)	IA 01-MAQ-0010_1	Clinton	Jun. 11, 1996 to Nov. 12, 2008	RK00.1M
13.	Rock Cr., upstream PCS Nitrogen	A1,B(WW2)	IA 01-MAQ-0010_2	Clinton	Jun. 11, 1996 to Nov. 12, 2008	RK03.7M
14.	Shrickers Slough	A1,B(WW1)	IA 01-MAQ-0005-L_0	Clinton	May 5, 1993 to Nov. 10, 2008	M508.1F
15.	Tete de Mortes Cr.	A1,B(WW1)	IA 01-TRK-0090_1	Jackson	Jun. 24, 1997 to Sep. 21, 2004	TM4.1M
16.	Turkey R., near mouth	A1,B(WW1)	IA 01-TRK-0200_0	Clayton	Jun. 22, 1998 to Sep. 21, 2004	TK04.8M
17.	Upper Iowa R. near mouth	A1,B(WW1)	IA 01-UIA-0090_0; IA 01-UIA-0100_0	Allamakee	Jun. 26, 1996 to Nov. 12, 2008	UI02.9M
18.	Wapsipinicon R., near mouth,	A1,B(WW1)	IA 01-WPS-0010_1	Clinton	May 5, 1993 to Nov. 12, 2008	WP02.6M
19.	Yellow R, near mouth	A1,B(WW1)	IA 01-YEL-0070_0	Allamakee	Apr. 19, 1996 to June 16, 2006	YL01.5M

**Designated Uses (from *Iowa Water Quality Standards* (IAC 2008)):

Class A1 = primary human contact/recreation;

Class B(WW1) = Waters in which temperature, flow and other habitat characteristics are suitable to maintain warm water game fish populations along with a resident aquatic community that includes a variety of native nongame fish and invertebrate species.

Class B(WW2) = Waters in which flow or other physical characteristics are capable of supporting a resident aquatic community that includes a variety of native nongame fish and invertebrate species. The flow and other physical characteristics limit the maintenance of warm water game fish populations.;

Table 5. Iowa beaches monitored by DNR/SHL or by local cooperators for indicator bacteria during recreational seasons from 2006-2008. Each group of beaches is listed alphabetically by lake name.

State-owned beaches 32 lakes; 37 beaches		City/county-owned beaches 28 lakes and beaches	
Lake Name	County	Lake Name	County
Ahquabi	Warren	Belva Deer	Keokuk
Anita	Cass	Briggs Woods	Hamilton
Backbone	Delaware	Central Park	Jones
Beeds	Franklin	Cold Springs	Cass
Big Creek	Polk	Crystal	Hancock
Big Spirit (Crandall's; Marble)	Dickinson	Don Williams	Boone
Black Hawk	Sac	Easter	Polk
Blue	Monona	Eldred-Sherwood	Hancock
Bob White	Wayne	Fairfield	Jefferson
Brushy Creek	Webster	Gabrielson Park	Buena Vista
Clear Lake (McIntosh Woods; Clear Lake)	Cerro Gordo	Hickory Grove	Story
Darling	Washington	Cornelia	Wright
Geode	Henry	Iowa	Iowa
George Wyth	Black Hawk	Pahoja	Lyon
Green Valley	Union	Little River	Decatur
Keomah	Mahaska	Lost Island	Palo Alto
Lacey Keosauqua	Van Buren	Malone	Clinton
MacBride	Johnson	Meredith Park	Pocahontas
Manawa	Pottawattamie	Ocheyedan Pits	Osceola
Nine Eagles	Decatur	Oldham	Monona
North Twin (east and west)	Calhoun	Orleans Beach	
Pine (south)	Hardin	Peterson	Story
Pleasant Creek	Linn	Pollmiller	Lee
Prairie Rose	Shelby	Spring	Greene
Red Haw	Lucas	Swan	Carroll
Rock Creek	Jasper	Treman	Calhoun
Springbrook	Guthrie	West Lake	Clarke
Three Fires	Taylor	Willow	Harrison
Union Grove	Tama		
Viking	Montgomery		
Wapello	Davis		
West Okoboji (Emerson, Gull Point, Pikes Point, Triboji)	Dickinson		

Table 6. Data completeness guidelines for using results of routine ambient water quality monitoring to make “monitored” assessments of designated beneficial uses for Section 305(b) water quality assessments in Iowa. “Monitored” assessments are used to place waters in Category 4 (impaired but TMDL not required) and Category 5 (impaired and TMDL required, the Section 303(d) list) of Iowa’s Integrated List/Report). Descriptions of “data required” have been modified to reflect the data gathering timeframe of the 2010 Section 303(d) listing cycle.

DESIGNATED USE	TYPE OF INFORMATION	DATA REQUIRED
Aquatic Life	Data for levels of toxics in waterbodies	Data collected quarterly or more frequently during calendar years 2006-2008; a minimum of 10 samples is needed.
	Data for levels of conventional pollutants (DO, pH, temp.)	Data collected monthly or more frequently during calendar years 2006-2008; a minimum of 10 samples is needed.
	Data from DNR biocriteria sampling at reference, test, and watershed sites.	At least two valid fish index of biotic integrity (IBI) or macroinvertebrate IBI’s for calibrated segments sampled during the most recent 5 complete calendar years (see Attachment 2 for more information).
	Data from the ISU/Iowa DNR statewide lake survey	Data collected at least 3 times per summer for at least 3 years (minimum of 9 samples).
	Results of fish kill investigations	Reports of pollutant-caused fish kills from 2006-2009.
Fish Consumption	Data for site-specific levels of toxic contaminants in fish tissue	All data on levels of toxic contaminants in fish tissue during the period covered by the 2010 assessment cycle (2006-2008).
Primary Contact Recreation	Data for levels of indicator bacteria (<i>E. coli</i>) from river waterbodies or non-beach areas of publicly-owned lakes or flood control reservoirs	Data collected monthly or more frequently during recreation seasons of 2006-2008; at least 10 temporally independent samples need to be collected.
	Data for levels of indicator bacteria (<i>E. coli</i>) from beach areas of publicly-owned lakes and flood control reservoirs	At least five samples approximately equally spaced over a 30-day period during recreation seasons of 2006-2008.
	Data from the IDNR-sponsored ISU/SHL statewide lake surveys for chlorophyll-a and Secchi depth	Data collected at least 3 times per summer for at least 3 consecutive years (minimum of 9 samples).
Drinking Water	Data for levels of <u>toxics</u>	Data collected quarterly or more frequently during calendar years 2006-2008; a minimum of 10 samples is needed.
	Data for levels of <u>nitrate</u>	Data collected monthly or more frequently during calendar years 2006-2008; a minimum of 10 samples is needed.

*Data that do not meet IDNR’s completeness guidelines can be used to develop “evaluated” (versus “monitored”) assessments for purposes of Section 305(b) water quality reporting. These “evaluated” assessments, however, are of generally lower confidence and are not appropriate for adding waters to IR Categories 4 or 5 (impairment categories) of the Integrated Report (IR). Evaluated assessments are, however, appropriate for adding waters to IR Categories 1, 2 and 3.

Table 7. Summary of Iowa water quality criteria used to make assessments of support of beneficial designated uses of Iowa surface waters for purposes of the 2010 Section 305(b) / Section 303(d) reporting/listing cycles. The criteria listed are only for those parameters used for the 2010 Section 305(b)/303(d) assessment/listing cycle. For a complete list and description of Iowa water quality criteria, see the *Iowa Water Quality Standards (IAC 2008)*; <http://www.iowadnr.gov/water/standards/files/chapter61.pdf>.

PARAMETER	DESIGNATED USE							
	Class A1, A2 and A3: swimmable	Class B(WW1): aquatic life	Class B(WW2) & B(WW3) aquatic life	Class B(CW1): coldwater aquatic life	Class B(CW2): coldwater aquatic life	Class B(LW): aquatic life of lakes and wetland	Class C: source of a water supply	HH (Human Health)
dissolved oxygen (mg/l) 16-hour minimum / 24-hour minimum)	none	5.0 / 5.0	5.0 / 4.0	7.0 / 5.0	7.0 / 5.0	5.0 / 5.0	none	none
temperature (added heat)	none	no increase > 3 C; increase < 1 C / hr; no increase above 32 C	no increase > 3 C; increase < 1 C / hr; no increase above 32 C	no increase > 2 C; increase < 1 C / hr; no increase above 20 C	no increase > 2 C; increase < 1 C / hr; no increase above 20 C	no increase > 2 C; increase < 1 C / hr; no increase above 32 C	none	none
pH	not < 6.5; not > 9. max. change = 0.5 units	not < 6.5; not > 9. max. change = 0.5 units	not < 6.5; not > 9. max. change = 0.5 units	not < 6.5; not > 9. max. change = 0.5 units	not < 6.5; not > 9. max. change = 0.5 units	not < 6.5; not > 9. max. change = 0.5 units	none	none
ammonia-nitrogen (mg/l)	none	criteria are dependent on the pH and temperature of the lake, stream or river; see Tables 3a through 3c of the <i>Iowa Water Quality Standards (IAC 2008)</i> for criteria for Class B(WW1), B(WW2), B(WW3), B(CW1), B(CW2), and B(LW) waters.					none	none
nitrate-nitrogen (mg/l)	none	none	none	none	none	none	10	none

Table 7 (continued)								
PARAMETER	DESIGNATED USE							
	Class A1, A2 & A3: swimmable	Class B(WW1): aquatic life	Class B(WW2) & B(WW3) aquatic life	Class B(CW1): coldwater aquatic life	Class B(CW2): coldwater aquatic life	Class B(LW): aquatic life of lakes and wetland	Class C: source of a water supply	HH (Human Health): fish / fish & water
chloride (mg/l)	none	none	none	none	none	none	250	none
fluoride (ug/l)	none	none	none	none	none	none	4000	none
<i>E. coli</i> (indicator bacteria)	[See Table 8]	none	none	none	none	none	none	none
TOXIC METALS (all values in ug/l; chronic / acute criteria are given for Class B designations; NA = value not applicable)								
Aluminum	None	87 / 750	87 / 750	87 / 1106	none	748 / 983	None	none
Arsenic	none	150 / 340	150 / 340	200 / 360	none	200 / 360	None	50 / 0.18
Cadmium*	none	0.27 / 2.13	0.27 / 2.13	1 / 4	none	1 / 4	5	168 / NA
chromium (VI)	none	11 / 16	11 / 16	40 / 60	none	10 / 15	100	3365 / NA
Copper*	none	9.3 / 14	9.3 / 14	20 / 30	none	10 / 20	none	1000 / 1300
Cyanide	none	5.2 / 22	5.2 / 22	5 / 20	none	10 / 45	none	140 / 140
Lead*	none	3.2 / 81.7	3.2 / 81.7	3 / 80	none	3 / 80	50	None
Mercury	none	0.9 / 1.64	0.9 / 1.64	3.5 / 6.5	none	0.91 / 1.7	none	0.15 / 0.05
Selenium	none	5 / 19.3	5 / 19.3	10 / 15	none	70 / 100	none	170 / 4200
Zinc*	none	120 / 120	120 / 120	200 / 220	none	100 / 110	none	2600 / 740
PESTICIDES (all values in ug/l; chronic / acute / human health criteria (HHC) are given; NA = value not applicable)								
2,4-D	none	none	none	none	none	none	none	100
2,4,5-TP (Silvex)	none	none	none	none	none	none	MCL: 50	none
Alachlor	none	none	none	none	none	none	MCL: 2	none
Atrazine	none	none	none	none	none	none	MCL: 3	none

*Criteria are based on a hardness of 100 mg/l using the respective equations in the Iowa Water Quality Standards (IAC 2008)

(<http://www.iowadnr.gov/water/standards/files/chapter61.pdf>).

Table 7 (continued)								
PARAMETER	DESIGNATED USE							
	Class A1, A2 & A3: swimmable	Class B(WW1): aquatic life	Class B(WW2) & B(WW3) aquatic life	Class B(CW1): coldwater aquatic life	Class B(CW2): coldwater aquatic life	Class B(LW): aquatic life of lakes and wetland	Class C: source of a water supply	HH (Human Health)
Carbofuran	none	none	none	none	none	none	40	none
Chlorpyrifos	none	0.041 / 0.083	0.041 / 0.083	0.041 / 0.083	none	0.041 / 0.083	none	none
DDT+DDD+DDE	none	0.001 / 1.1	0.001 / 1.1	0.001 / 0.9	none	0.001 / 0.55	none	0.0022 / 0.0022
Dieldrin	none	0.056 / 0.24	0.056 / 0.24	0.056 / 0.24	none	0.056 / 0.24	none	0.00054 / 0.00052
Dinoseb	none	none	none	none	none	none	7	none
Lindane	none	NA / 0.95	NA / 0.95	NA / 0.95	none	NA / 0.95	none	1.8 / 0.98
Parathion	none	0.13 / 0.65	0.13 / 0.65	0.13 / 0.65	none	0.13 / 0.65	none	none
Picloram	none	none	none	none	none	none	500	none
Simazine	none	none	none	none	none	none	4	none

Table 8. Summary of Iowa water quality criteria for indicator bacteria (*E. coli*) in surface waters designated in the *Iowa Water Quality Standards* (IAC 2008) for either primary contact recreation, secondary contact recreation, or children's recreational use. The *E. coli* content shall not exceed the following levels when the Class A uses can reasonably be expected to occur.

	Class A1: primary contact recreational use*	Class A2: secondary contact recreational use*	Class A3: children's recreational use*
Geometric Mean (No. of <i>E. coli</i> organisms/100 ml of water)	126	630	126
Sample Maximum (No. of <i>E. coli</i> organisms/100 ml of water)	235	2,880	235

* Criteria apply from March 15 through November 15 (i.e., the "recreational season") except year-round for Class A2 waters that are also designated for Class B(CW1) [coldwater aquatic life] uses.

Table 9. General water quality criteria to protect beneficial general uses for all Iowa surface waters (from the *Iowa Water Quality Standards*, IAC, Section 61.3(2)).

The following criteria are applicable to all surface waters including general use and designated use waters, at all places and at all times, to protect livestock and wildlife watering, aquatic life, noncontact recreation, crop irrigation, and industrial, domestic, agricultural, and other incidental water withdrawal uses not protected by specific numerical criteria in the subrule 61.3(3) of the *Iowa Water Quality Standards* (IAC 2008):

- All waters of the state shall be "free from" the following:
 - substances attributable to point source wastewater dischargers that will settle to form sludge deposits;
 - floating debris, oil, grease, scum and other materials from wastewater discharges or agricultural practices in amounts sufficient to create a nuisance;
 - materials attributable to wastewater discharges or agricultural practices producing objectionable color, odor, or other aesthetically objectionable conditions;
 - substances attributable to wastewater discharges or agricultural practices in concentrations or combinations which are acutely toxic to human, animal, or plant life;
 - substances attributable to wastewater discharges or agricultural practices in quantities which would produce undesirable or nuisance aquatic life;
- The turbidity of a receiving water shall not be increased by more than 25 nephelometric turbidity units by any point source discharge;
- The acceptable level of total dissolved solids (TDS) and constituent cations and anions will be established on a site-specific basis (see *Supporting Document for Iowa Water Quality Management Plans, Chapter IV* (<http://www.iowadnr.com/water/standards/files/04part2.pdf>) [Note: 1000 mg/l (TDS) is the threshold in-stream level at which negative impacts to the uses of the receiving stream may begin to occur.]
- The *Escherichia coli* content of water which enters a sinkhole or losing stream segment, regardless of the waterbody's designated use, shall not exceed a geometric mean of 126 organisms per 100 ml or a sample maximum of 235 organisms/100 ml. No new wastewater discharges will be allowed on watercourses which directly or indirectly enter sinkholes or losing stream segments.

Table 10. Methods for determining support of AQUATIC LIFE USES for general use and designated use surface waters in Iowa for 2010 Section 305(b) reporting and 303(d) listing. For shallow lakes, TSI = trophic state index of Carlson (1977).					
Type of waterbody	Source of Information	Fully Supported	Fully Supported/Threatened	Partially Supporting	Not Supporting
Rivers, streams, lakes & flood control reservoirs	Data from ambient water quality monitoring during current reporting period.	Up to one violation of acute toxicity criteria if grab samples are collected quarterly or more frequently. Chronic criteria for toxics and criteria for conventional pollutants exceeded in $\leq 10\%$ of samples.	Criteria for conventional pollutants or chronic toxicity criteria are exceeded in no more than 10% of samples but levels are trending such that future impairment is likely.	Criteria for conventional pollutants or chronic toxicity criteria exceeded in from 11-25% of samples (90% confidence level).	More than one violation of acute toxicity criteria if samples collected quarterly or more often; or, criteria for conventionals or chronic toxicity criteria exceeded in more than 25% of samples.
Shallow lakes (see Attachment 4)	IDNR water quality monitoring, 2006-08	TSI values for both chlorophyll-a and Secchi depth are < 65	TSI values for both chlorophyll-a or Secchi depth are < 65 but at least one parameter exhibits an adverse trend over time such that impairment is likely to occur.	TSI values for either chlorophyll-a or Secchi depth are equal to or greater than 65 but less than 70.	TSI values for both chlorophyll-a and Secchi depth are equal to or greater than 65, or the TSI value for either parameter is equal to or greater than 70.
Warmwater Streams and Rivers	Stream biocriteria sampling data (see Attachment 2)	Scores for fish or macroinvertebrate indexes of biotic integrity equal or exceed the ecoregion / subecoregion biological impairment criterion.	[Category not used for Section 305(b) reporting in Iowa.]	Scores for <u>one</u> of the indexes of biotic integrity (fish or macroinvertebrate) significantly less than the ecoregion / subecoregion biological impairment criterion.	Scores for <u>both</u> indexes of biotic integrity (fish and macroinvertebrate) significantly less than the ecoregion / subecoregion biological impairment criterion.
Coldwater Streams	Stream biocriteria sampling data (See Attachment 2)	Two or less of the eight biological indicators less than the 25 th percentile of the respective indicator value for Iowa coldwater streams.	[Category not used for Section 305(b) reporting in Iowa.]	From five to six of the eight biological indicators less than the 25 th percentile of the respective indicator value for Iowa coldwater streams.	From seven to eight of the eight biological indicators less than the 25 th percentile of the respective indicator value for Iowa coldwater streams.
Rivers, streams, lakes & flood control reservoirs	Fish kill reports*	No pollutant-caused fish kills reported.	[Category not used for Section 305(b) reporting in Iowa.]	One pollutant-caused fish kill reported.	More than one pollutant-caused reported.

*Sources of fish kills will be reviewed to determine whether the affected waterbody is a candidate for 303(d) listing.

Table 11. Methods for determining support of classified, beneficial uses for FISH CONSUMPTION, PRIMARY CONTACT RECREATION, and DRINKING WATER for surface waters in Iowa for 2010 Section 305(b) reporting and 303(d) listing. Note: TSI = trophic state index of Carlson (1977).					
Type of Waterbody	Source of Information	Fully Supported	Fully Supported/Threatened	Partially Supporting	Not Supporting
FISH CONSUMPTION USES					
Streams, rivers, lakes, & flood control reservoirs	monitoring of levels of toxic contaminants in fish tissue	Waterbody is not covered by an IDNR/IDPH-issued fish consumption advisory	Results of monitoring have not resulted in issuance of an advisory but results of monitoring show an adverse trend suggesting that issuance of an advisory is imminent.	Levels of one or more toxics have exceeded the respective IDNR/IDPH advisory trigger levels in two consecutive samplings and a “one meal/week” advisory is in effect for the general population	Levels of one or more toxics have exceeded the respective IDNR/IDPH advisory trigger levels in two consecutive samplings and a “do not eat” advisory is in effect for the general population
CLASS A1 and A3 PRIMARY CONTACT RECREATION (SWIMMABLE) USES					
Streams, rivers, lakes, & flood control reservoirs	monthly monitoring data for indicator bacteria	Geometric mean of <i>E. coli</i> samples \leq 126 orgs / 100 ml and \leq 10% of samples exceed 235 orgs/100 ml.	Geometric mean of <i>E. coli</i> samples \leq 126 orgs / 100 ml and \leq 10% of samples $>$ 235 orgs/100 ml but worsening trend suggests that future impairment is likely.	Geometric mean of <i>E. coli</i> samples \leq 126 orgs/100 ml but more than 10% of samples exceed 235 orgs/100 ml (90% confidence level).	Geometric mean of <i>E. coli</i> samples $>$ 1,000 orgs/100.
Lake beaches	weekly monitoring data for indicator bacteria	Geometric mean of at least 5 <i>E. coli</i> samples collected over a 30-day period \leq 126 orgs / 100 ml and \leq 10% of samples exceed 235 orgs/100 ml.	Geometric mean of <i>E. coli</i> samples \leq 126 orgs/100 ml and \leq 10% of samples $>$ 235 orgs/100 ml but worsening trend suggests that future impairment is likely.	Seasonal geometric mean of <i>E. coli</i> samples \leq 126 orgs/100 ml but $>$ 10% of samples exceed 235 orgs/100 ml (90% confidence level).	Geometric mean of at least 5 <i>E. coli</i> samples over a 30-day period $>$ 1,000 orgs / 100 ml.
Lakes (see Attachment 3)	ISU & SHL ambient lake monitoring, 2004-08	TSI values for both chlorophyll-a and Secchi depth are $<$ 65	TSI values for both chlorophyll-a or Secchi depth are $<$ 65 but at least one parameter exhibits an adverse trend over time such that impairment is likely to occur.	TSI values for either chlorophyll-a or Secchi depth are equal to or greater than 65 but less than 70.	TSI values for both chlorophyll-a and Secchi depth are equal to or greater than 65, or the TSI value for either parameter is equal to or greater than 70.
Streams, rivers, lakes, & flood control reservoirs	Closure* of beaches and other swimming areas	No swimming area closures in effect during the biennial reporting period	Geometric mean of <i>E. coli</i> samples \leq 126 orgs / 100 ml but worsening trend suggests that future impairment is likely.	One swimming area closure of less than one week duration during the biennial reporting period	More than one swimming area closure, or one swimming area closure of more than one week duration during the biennial period

*Elevated levels of indicator bacteria at beaches of Iowa's state-owned lakes can trigger the posting of a “swimming is not recommended” sign. The posting of this sign, however, does not mean that the beach is closed. IDNR can, and will, close beaches in case of an emergency health risk such as a wastewater bypass, spill of a hazardous chemical, or a localized outbreak of an infectious disease (see IDNR 2010: *Iowa's beach monitoring program, 2008*).

Table 11. (continued).

Type of Waterbody	Source of Information	Fully Supported	Fully Supported/Threatened	Partially Supporting	Not Supporting
CLASS A2 SECONDARY CONTACT RECREATION (SWIMMABLE) USES					
Streams, rivers, lakes, & flood control reservoirs	monthly monitoring data for indicator bacteria	Geometric mean of <i>E. coli</i> samples \leq 630 orgs / 100 ml and \leq 10% of samples exceed 2,880 orgs/100 ml.	Geometric mean of <i>E. coli</i> samples \leq 630 orgs / 100 ml and \leq 10% of samples $>$ 2,880 orgs/100 ml but worsening trend suggests that future impairment is likely.	Geometric mean of <i>E. coli</i> samples \leq 630 orgs/100 ml but more than 10% of samples exceed 2,880 orgs/100 ml (90% confidence level).	Geometric mean of <i>E. coli</i> samples $>$ 1000 orgs/100.
DRINKING WATER USES					
Waterbodies designated for use as a source of potable water (=raw water source)	ambient monitoring data for <u>toxics</u>	Average levels of toxic metals or pesticides are less than human health criteria (HH) or maximum contaminant levels (MCLs).	Average levels of toxic metals or pesticides \leq HH criteria or MCLs, but the average levels of at least one toxic is trending upward toward its respective HH criteria or MCL; waterbody is considered "impaired"	[category not used for Section 305(b) reporting]	Average level of toxic metals or pesticides greater than the MCL.
Waterbodies designated for use as a source of potable water (=raw water source)	ambient monitoring data for <u>atrazine</u>	All moving annual average levels of atrazine are less than the maximum contaminant level (MCL) of 3 ug/l.	All moving annual average levels are less than the MCL, but average levels are trending upward toward the MCL; waterbody is considered "impaired"	[category not used for Section 305(b) reporting]	One or more of the moving annual average levels exceed the MCL.
Waterbodies designated for use as a source of potable water (=raw water source)	ambient monitoring data for <u>nitrate</u>	No more than 10% of samples violate the maximum contaminant level (MCL) for nitrate.	No more than 10% of samples violate the MCL for nitrate but nitrate levels are trending upward such that impairment is likely.	From 11-25% of samples violate the MCL for nitrate (90% confidence level).	More than 25% of samples exceed the MCL for nitrate.
Municipal drinking water (=finished water)	public water supplies using surface waters	No drinking water supply closures or advisories in effect; water not treated beyond reasonable levels	[Category not used for Section 305(b) reporting or 303(d) listing.]	One drinking water advisory lasting 30 days or less per year, or other problems not requiring closure but affecting treatment costs	One or more drinking water supply advisory lasting more than 30 days per year, or one or more drinking water supply closures per year

Table 12. Sample size and number of exceedances required to determine an impaired beneficial use (10% exceedance) to maintain a greater than 90 percent confidence level as reported by Lin et al. (2000) (table excerpted from NDEQ 2006).

Minimum number of exceedances required to maintain a >90% confidence that a designated use is impaired (10% exceedance).					
Sample Size (n)	Number of observations exceeding required to define an impaired use	Confidence Level	Sample Size (n)	Number of observations exceeding required to define an impaired use	Confidence Level
10	3	0.930	56	10	0.951
11	3	0.910	57	10	0.945
12	4	0.974	58	10	0.940
13	4	0.966	59	10	0.933
14	4	0.956	60	10	0.927
15	4	0.944	61	10	0.920
16	4	0.932	62	10	0.913
17	4	0.917	63	10	0.905
18	4	0.911	64	11	0.948
19	5	0.965	65	11	0.943
20	5	0.957	66	11	0.938
21	5	0.948	67	11	0.932
22	5	0.938	68	11	0.926
23	5	0.927	69	11	0.920
24	5	0.915	70	11	0.913
25	5	0.902	71	11	0.906
26	6	0.960	72	12	0.947
27	6	0.953	73	12	0.942
28	6	0.945	74	12	0.937
29	6	0.936	75	12	0.931
30	6	0.927	76	12	0.926
31	6	0.917	77	12	0.920
32	6	0.906	78	12	0.913
33	7	0.958	79	12	0.907
34	7	0.952	80	13	0.946
35	7	0.945	81	13	0.942
36	7	0.937	82	13	0.937
37	7	0.929	83	13	0.931
38	7	0.920	84	13	0.926
39	7	0.911	85	13	0.920
40	7	0.900	86	13	0.914
41	8	0.952	87	13	0.908
42	8	0.946	88	13	0.901
43	8	0.939	89	14	0.941
44	8	0.932	90	14	0.937
45	8	0.924	91	14	0.932
46	8	0.916	92	14	0.927
47	8	0.907	93	14	0.921
48	9	0.954	94	14	0.915
49	9	0.948	95	14	0.910
50	9	0.942	96	14	0.903
51	9	0.936	97	15	0.941
52	9	0.929	98	15	0.937
53	9	0.922	99	15	0.932
54	9	0.914	100	15	0.927
55	9	0.906			


Table 13. Summary of Iowa's protocol for issuing fish consumption advisories. Issuance of an advisory requires two consecutive samplings that show contaminant levels above advisory trigger levels. This protocol was developed by the Iowa Department of Public Health in cooperation with IDNR (IDPH 2007).

Parameter	Contaminant Concentrations in fish fillets:		
	Unrestricted consumption	Limit consumption to one meal per week	Do not eat
PCBs	0 to 0.2 ppm	0.2 to 2.0 ppm	> 2.0 ppm
Mercury	0. to 0.3 ppm	0.3 to 1.0 ppm	> 1.0 ppm
Chlordane	0. to 0.6 ppm	0.6 to 5.0 ppm	> 5.0 ppm

Table 14. Placement of fish kill-affected waters into IR categories for Iowa's 2010 Integrated Reporting cycle.

Year of kill:	Years without a reported kill:	Pollutant-caused kill; no restitution sought	Pollutant-caused kill; restitution sought	No cause identified; or non-pollutant/natural kill
2009	0	5a/5b	4d	2b/3b
2008	1	5a/5b	4d	2b/3b
2007	2	5a/5b	4d	2b/3b
2006	3	5a/5b	4d	2b/3b
2005	4	5a/5b	4d	2b/3b
2004	5	5a/5b	4d	2b/3b
2003	6	5a/5b	2b/3b	2b/3b
2002	7	5a/5b	2b/3b	2b/3b
2001	8	5a/5b	2b/3b	2b/3b
2000	9	5a/5b	2b/3b	2b/3b
1999	10	5a/5b	2b/3b	2b/3b
1998	11	5a/5b	3a	3a
1997	12	5a/5b	3a	3a

Figure 1. Use of water quality data and information for Iowa's 2010 Integrated Report (Section 305(b)/303(d) report/list).



Attachment 1.
Excerpt from Senate File 2371: Iowa's credible data legislation

PAG LIN

1 1 SENATE FILE 2371
1 2
1 3 AN ACT
1 4 RELATING TO THE ESTABLISHMENT OF A WATER QUALITY INITIATIVE
1 5 PROGRAM BY THE DEPARTMENT OF AGRICULTURE AND LAND STEWARD-
1 6 SHIP AND THE DEPARTMENT OF NATURAL RESOURCES, DEFINING
1 7 AND PROVIDING FOR THE USE OF **CREDIBLE DATA** FOR QUALITY CONTROL
1 8 AND ASSURANCE PROCEDURES, AND PROVIDING FOR OTHER PROPERLY
1 9 RELATED MATTERS, AND PROVIDING AN APPLICABILITY DATE.
1 10
1 11 BE IT ENACTED BY THE GENERAL ASSEMBLY OF THE STATE OF IOWA:
1 12

7 21 Sec. 9. Section 455B.171, Code 1999, is amended by adding
7 22 the following new subsections:

7 23 NEW SUBSECTION. 10A. "**Credible data**" means scientifically
7 24 valid chemical, physical, or biological monitoring data
7 25 collected under a scientifically accepted sampling and
7 26 analysis plan, including quality control and quality assurance
7 27 procedures. Data dated more than five years before the
7 28 department's date of listing or other determination under
7 29 section 455B.194, subsection 1, shall be presumed not to be
7 30 credible data unless the department identifies compelling
7 31 reasons as to why the data is credible.

7 32 NEW SUBSECTION. 14A. "Historical data" means data
7 33 collected more than five years before the department's date of
7 34 listing or other determination under section 455B.194,
7 35 subsection 1.

8 1 NEW SUBSECTION. 19A. "Naturally occurring condition"
8 2 means any condition affecting water quality which is not
8 3 caused by human influence on the environment including, but
8 4 not limited to, soils, geology, hydrology, climate, wildlife
8 5 influence on the environment, and water flow with specific
8 6 consideration given to seasonal and other natural variations.

8 7 NEW SUBSECTION. 31A. "Section 303(d) list" means any list
8 8 required under 33 U.S.C. } 1313(d).

8 9 NEW SUBSECTION. 31B. "Section 305(b) list" means any
8 10 report or list required under 33 U.S.C. } 1315(b).

8 11 NEW SUBSECTION. 39A. "Total maximum daily load" means the
8 12 same as in the federal Water Pollution Control Act.

8 13 Sec. 10. NEW SECTION. 455B.193 QUALIFICATIONS FOR
8 14 COLLECTION OF CREDIBLE DATA.

8 15 For purposes of this part, all of the following shall
8 16 apply:

8 17 1. Data is not credible data unless the data originates
8 18 from studies and samples collected by the department, a
8 19 professional designee of the department, or a qualified
8 20 volunteer. For purposes of this subsection, "professional
8 21 designee" includes governmental agencies other than the
8 22 department, and a person hired by, or under contract for
8 23 compensation with, the department to collect or study data.

8 24 2. All information submitted by a qualified volunteer
8 25 shall be reviewed and approved or disapproved by the
8 26 department. The qualified volunteer shall submit a site
8 27 specific plan with data which includes information used to
8 28 obtain the data, the sampling and analysis plan, and quality
8 29 control and quality assurance procedures used in the
8 30 monitoring process. The qualified volunteer must provide
8 31 proof to the department that the water monitoring plan was
8 32 followed. The department shall review all data collected by a
8 33 qualified volunteer, verify the accuracy of the data collected
8 34 by a qualified volunteer, and determine that all components of
8 35 the water monitoring plan were followed.

9 1 3. The department shall retain all information submitted
9 2 by a qualified volunteer submitting the information for a
9 3 period of not less than ten years from the date of receipt by
9 4 the department. All information submitted shall be a public
9 5 record.

9 6 4. The department shall adopt rules establishing
9 7 requirements for a person to become a qualified volunteer.
9 8 The department of natural resources shall develop a
9 9 methodology for water quality assessments as used in the
9 10 section 303(d) listings and assess the validity of the data.

9 11 Sec. 11. NEW SECTION. 455B.194 CREDIBLE DATA REQUIRED.

9 12 1. The department shall use credible data when doing any
9 13 of the following:

9 14 a. Developing and reviewing any water quality standard.

9 15 b. Developing any statewide water quality inventory or
9 16 other water assessment report.

9 17 c. Determining whether any water of the state is to be
9 18 placed on or removed from any section 303(d) list.

9 19 d. Determining whether any water of the state is
9 20 supporting its designated use or other classification.

9 21 e. Determining any degradation of a water of the state
9 22 under 40 C.F.R. } 131.12.

9 23 f. Establishing a total maximum daily load for any water
9 24 of the state.

9 25 2. Notwithstanding subsection 1, credible data shall not
9 26 be required for any section 305(b) report and credible data
9 27 shall not be required for the establishment of a designated
9 28 use or other classification of a water of the state.

9 29 3. This section shall not be construed to require credible
9 30 data as defined in section 455B.171, subsection 10A, in order
9 31 for the department to bring an enforcement action for an
9 32 illegal discharge.

9 33 Sec. 12. NEW SECTION. 455B.195 USE OR ANALYSIS OF
9 34 CREDIBLE DATA.

9 35 1. For any use or analysis of credible data described in
10 1 section 455B.194, subsection 1, all of the following shall
10 2 apply:

10 3 a. The use of credible data shall be consistent with the
10 4 requirements of the federal Water Pollution Control Act, 33
10 5 U.S.C. } 1251 et seq.

10 6 b. The data quality for removal of water of the state from
10 7 any list of impaired waters including any section 303(d) list
10 8 shall be the same as the data quality for adding a water to
10 9 that list.

10 10 c. A water of the state shall not be placed on any section
10 11 303(d) list if the impairment is caused solely by violations
10 12 of national pollutant discharge elimination system program
10 13 permits or stormwater permits issued pursuant to section
10 14 455B.103A and the enforcement of the pollution control
10 15 measures is required.

10 16 d. A water of the state shall not be placed on any section
10 17 303(d) list if the data shows an impairment, but existing
10 18 technology-based effluent limits or other required pollution
10 19 control measures are adequate to achieve applicable water
10 20 quality standards.

10 21 e. If a pollutant causing an impairment is unknown, the
10 22 water of the state may be placed on a section 303(d) list.
10 23 However, the department shall continue to monitor the water of
10 24 the state to determine the cause of impairment before a total
10 25 maximum daily load is established for the water of the state
10 26 and a water of the state listed with an unknown status shall
10 27 retain a low priority for a total maximum daily load
10 28 development until the cause of the impairment is determined
10 29 unless the department, after taking into consideration the use
10 30 of the water of the state and the severity of the pollutant,
10 31 identifies compelling reasons as to why the water of the state
10 32 should not have a low priority.

10 33 f. When evaluating the waters of the state, the department
10 34 shall develop and maintain three separate listings including a
10 35 section 303(d) list, a section 305(b) report, and a listing
11 1 for which further investigative monitoring is necessary. The
11 2 section 305(b) report shall be a summary of all potential
11 3 impairments for which credible data is not required. If
11 4 credible data is not required for a section 305(b) report, the
11 5 placement of a water of the state on any section 305(b) report
11 6 alone is not sufficient evidence for the water of the state's
11 7 placement on any section 303(d) list. When developing a
11 8 section 303(d) list, the department is not required to use all
11 9 data, but the department shall assemble and evaluate all
11 10 existing and readily available water quality-related data and
11 11 information. The department shall provide documentation to
11 12 the regional administrator of the federal environmental
11 13 protection agency to support the state's determination to list
11 14 or not to list its waters.

11 15 g. The department shall take into consideration any
11 16 naturally occurring condition when placing or removing any
11 17 water of the state on any section 303(d) list, and
11 18 establishing or allocating responsibility for a total maximum
11 19 daily load.

11 20 h. Numerical standards shall have a preference over
11 21 narrative standards. A narrative standard shall not
11 22 constitute the basis for determining an impairment unless the
11 23 department identifies specific factors as to why a numeric
11 24 standard is not sufficient to assure adequate water quality.

11 25 i. If the department has obtained credible data for a
11 26 water of the state, the department may also use historical
11 27 data for that particular water of the state for the purpose of
11 28 determining whether any trends exist for that water of the
11 29 state.

11 30 2. This section shall not be construed to require or

11 31 authorize the department to perform any act listed in section
11 32 455B.194, subsection 1, not otherwise required or authorized
11 33 by applicable law.

11 34 Sec. 13. LEGISLATIVE STUDY. The legislative council is
11 35 requested to establish an interim study relating to the use of
12 1 plant nutrients on Iowa soil. The committee is directed to
12 2 submit its findings, with any recommendations, in a report to
12 3 the general assembly not later than January 15, 2001.

12 4 Sec. 14. **APPLICABILITY OF SECTION 303(d) LISTS.** This Act
12 5 takes effect July 1, 2000. However, any requirements under
12 6 this Act which apply to a section 303(d) list shall not apply
12 7 for the section 303(d) list for the year 2000, but any
12 8 requirements shall take effect for all section 303(d) lists
12 9 created after the year 2000 list.

12 10

12 11

12 12

12 13

MARY E. KRAMER
President of the Senate

12 14

12 15

12 16

12 17

12 18

BRENT SIEGRIST
Speaker of the House

12 19

12 20

12 21 I hereby certify that this bill originated in the Senate and
12 22 is known as Senate File 2371, Seventy-eighth General Assembly.

12 23

12 24

12 25

12 26

MICHAEL E. MARSHALL
Secretary of the Senate

12 27

12 28 Approved

, 2000

12 29

12 30

12 31

12 32 THOMAS J. VILSACK

12 33 Governor

Attachment 2

GUIDELINES FOR DETERMINING SECTION 305(B) AQUATIC LIFE USE SUPPORT (ALUS) USING STREAM BIOCRITERIA SAMPLING DATA FOR THE 2006 SECTION 305(B) REPORTING AND SECTION 303(D) LISTING CYCLES

Introduction:

Since the late 1980s, U.S. EPA has encouraged states to develop and adopt narrative and biological criteria (biocriteria) for surface waters. Biocriteria are narrative or numeric expressions that describe the best attainable biological integrity (reference condition) of aquatic communities inhabiting waters of a given designated aquatic life use (U.S. EPA 1990a). Supported by a water quality planning grant from the U.S. EPA Region VII, geographers of the U.S. EPA Corvallis Environmental Research Laboratory collaborated with DNR staff to revise and subdivide the ecoregions in Iowa (see Omernik et al. 1993; Griffith et al. 1994). As part of this effort, a list of candidate stream reference sites was generated. Reference sites are located on the least impacted streams within an ecoregion or subecoregion. Reference sites can thus serve as benchmarks to which water quality-impaired streams can be compared. A pilot reference site sampling study was conducted in 1994 to develop standardized data collection procedures for assessing the quality of aquatic habitat and for sampling benthic macroinvertebrate and fish communities (Wilton 1996). Approximately 100 reference sites were sampled during the initial reference site sampling period 1994-1998; an additional 75 sites were sampled with the biocriteria sampling protocol as part of test site sampling and sampling for watershed projects. These data, as well as more recent reference site sampling data from 1999-2004, were used to develop and calibrate indicators of stream biological integrity (Wilton 2004) and biological assessment criteria used in assessments of aquatic life use support for the 2006 Section 305(b) report. For a discussion of the process used to calculate the bioassessment criteria, please see the addendum to this attachment.

The bioassessment indicators were originally calibrated for assessing support of Class B(LR) and Class B(WW) warmwater aquatic life uses in wadeable stream segments. The indicators were not calibrated for small headwater "General Use" streams or nonwadeable

warmwater rivers having watershed drainage areas ≥ 500 mi². In the absence of specifically calibrated indicators for these types of warmwater lotic systems, the current indicators and criteria have been applied; however, these assessments are considered “evaluated” rather than “monitored” assessments to reflect a greater degree of uncertainty in the assessment conclusions. Separate indicators and guidelines described later in this section have been developed for determining the level of support for the Class B(CW) coldwater aquatic life uses designated for trout streams of northeastern Iowa.

Uses designated for individual stream and river reaches in Iowa are summarized in the “Water Use Designations” portion of the *Iowa Water Quality Standards* (IAC 2003); definitions of designated uses [e.g., Class B(WW), Class B(LR), and Class B(CW)] are presented in the *Iowa Water Quality Standards* (IAC 2003).

The Iowa DNR uses a Benthic Macroinvertebrate Index of Biotic Integrity (BMIBI) and a Fish Index of Biotic Integrity (FIBI) to summarize biological sampling data. The BMIBI and FIBI combine several quantitative measurements or “metrics” that provide a broad assessment of stream biological conditions. A metric is a characteristic of the biological community that can be measured reliably and responds predictably to changes in stream quality. The BMIBI and FIBI each contain twelve metrics that relate to species diversity, relative abundance of sensitive and tolerant organisms, and the proportion of individuals belonging to specific feeding and habitat groups. The metrics are numerically ranked and their scores are totaled to obtain an index rating from 0 (poor) – 100 (optimum). Qualitative scoring ranges of poor, fair, good, and excellent have been established that reflect the biological community characteristics found at each level (Table 2-1a, 2-1b). These qualitative ranges are general interpretative guidelines only. To assess support of aquatic life uses, sample site IBI scores are compared against Biological Impairment Criteria (BIC) (Table 2-2), which more specifically reflect reference conditions defined by ecoregion and habitat class.

Determining Support of General Use, Class B(LR) and B(WW) Aquatic Life Uses

Aquatic life use assessments for the 2006 Integrated Report cycle were updated using biological assemblage sampling data from 2003 and 2004. The primary types and sources of data are: a) benthic macroinvertebrate and fish assemblage data collected as part of the

DNR/SHL stream biocriteria project; b) fish assemblage data collected by staff of the DNR Fisheries Bureau. Before making assessments, data completeness and quality are evaluated. "Comparable" data are considered as having completeness and quality that is comparable to biocriteria project data used to develop reference biotic indexes and impairment criteria. These data are used to make "monitored" (higher confidence) assessments. "Tentative" data are considered as having lesser or uncertain levels of completeness and quality documentation. These data are used to make "evaluated" (lower confidence) assessments.

To determine the level of aquatic life use support for a stream sampling site, the BMIBI and/or FIBI scores from that stream are compared against index levels measured at reference stream sites located in the same ecological region. Reference sites are also stratified by habitat class in certain ecoregions where statistically significant differences have been found between reference sites having abundant coarse substrates and riffle habitat versus those lacking these habitat characteristics. A set of biological assessment criteria were specifically developed for the 2006 305(b) report using stream reference site data from 1994-2004. The 25th percentile values of the reference site BMIBI and FIBI index scores within a given ecoregion or habitat class were used as the biological impairment criteria (BIC) for 305(b) assessment purposes (Table 2-2). Use of the reference 25th percentile as an impairment threshold is consistent with biocriteria development guidance (U.S. EPA 1996), and has demonstrated efficacy in state bioassessment programs (Yoder and Rankin 1995). Biotic index performance evaluation in Iowa found little or no overlap of index interquartile ranges between reference sites and test (impacted) sites, which suggests that reference 25th percentile levels are appropriate for assessing biological impairment.

Generally, a stream is considered biologically impaired if one or both of its index scores are significantly lower than the BIC. An uncertainty adjustment value (UAV) equal to 8 BMIBI points or 7 FIBI points is applied in cases where single sample data are used to assess aquatic life use support status. The UAV reflects the typical year-to-year IBI scoring variation observed among least disturbed reference sites throughout Iowa. It is used to identify stream segments that are within a reasonable margin of error from the lower 25th percentile of reference site IBI scores and may be considered a higher priority for follow-up sampling in order to better determine the status of aquatic life uses.

“Monitored” assessments are those for which biocriteria project comparable data are available to assess a “calibrated” stream segment, which is defined as wadeable streams designated as B(LR) or B(WW) in 2004 and have a watershed drainage area < 500 square miles. “Evaluated” assessments are generally of two kinds: 1) cases in which data of lesser or uncertain comparability are used to assess a “calibrated” segment; 2) cases where biotic index data are used to assess “uncalibrated” segments (i.e., general use segments or non-wadeable river segments having watershed drainage area $\geq 500 \text{ mi}^2$).

Aquatic Life Use Support Guidelines

The following guidelines are used to make aquatic life use status recommendations on the basis of biological sampling data only. In many cases, water quality monitoring data are also available to evaluate aquatic life use status from the perspective of chemical and physical water quality standards attainment. In these cases, a weight of evidence approach is taken to make adjustments and assign the most appropriate aquatic life use status category.

Fully Supporting “Monitored”

- Assessments for calibrated stream segments having comparable data consisting of at least one valid BMIBI score and at least one valid FIBI score, and the single score(s) or the average(s) of multiple scores equal or exceed the BIC.

Fully Supporting “Evaluated”

- Assessments for calibrated segments having comparable data consisting of at least one valid BMIBI or FIBI score but not both index scores, and the single score and/or the average of multiple scores for that index equal or exceed the BIC; OR.
- Assessments for calibrated segments having tentative data consisting of at least one valid BMIBI score and/or FIBI score, and the single score(s) and/or the average(s) of multiple scores equal or exceed the BIC; OR.
- Assessments for uncalibrated segments having comparable or tentative data consisting of at least one valid BMIBI score and/or FIBI score, and the single score(s) or the average(s) of multiple scores equal or exceed the BIC.

Partially Supporting “Monitored”

- Assessments for calibrated segments having comparable data consisting of at least one valid BMIBI score and/or FIBI score.
 - If valid score(s) for only one index, the single score plus the applicable UAV is less than the BIC or the average of multiple scores is less than the BIC;
OR,
 - If valid score(s) for both indexes, then: 1) the single score(s) plus the applicable UAV(s) is less than the BIC or the average(s) of multiple scores is less than the BIC, and 2) the single score or the average of multiple scores for at least one index does not fall in the qualitative range indicating “poor” biocondition (see Tables 2-1a and 2-1b).

Partially Supporting “Evaluated”

- Assessments for uncalibrated segments having comparable or tentative data consisting of at least one valid BMIBI score and/or FIBI score.
 - If valid score(s) for only one index, the single score plus the applicable UAV is less than the BIC or the average of multiple scores is less than the BIC;
OR,
 - If valid score(s) for both indexes, then: 1) the single score(s) plus the applicable UAV(s) is less than the BIC or the average(s) of multiple scores is less than the BIC, and 2) the single score or the average of multiple scores for at least one index does not fall in the qualitative range indicating poor biocondition.
- Assessments for calibrated segments having tentative data consisting of at least one valid BMIBI score and/or FIBI score.
 - If valid score(s) for only one index, the single score plus the applicable UAV is less than the BIC or the average of multiple scores is less than the BIC;
OR,
 - If valid score(s) for both indexes, then: 1) the single score(s) plus the applicable UAV(s) is less than the BIC or the average(s) of multiple scores is less than the BIC, and 2) the single score or the average of multiple scores for at least one index does not fall in the qualitative range indicating poor biocondition.

Not Supporting “Monitored”

- Assessments for calibrated segments having comparable data consisting of at least one valid BMIBI score and one valid FIBI score, and both of the following conditions are true: 1) the single score(s) plus the applicable UAV(s) is less than the BIC or the average(s) of multiple scores is less than the BIC; and 2) the single score or the average of multiple scores for both the BMIBI and the FIBI fall in the qualitative range indicating poor biocondition.

Not Supporting "Evaluated"

- Assessments for uncalibrated segments having comparable or tentative data consisting of at least one valid BMIBI score and one valid FIBI score, and both of the following conditions are true:
 - 1) the single score(s) plus the applicable UAV(s) is less than the BIC or the average(s) of multiple scores is less than the BIC;
 - 2) the single score or the average of multiple scores for both the BMIBI and the FIBI fall in the range indicating poor biocondition.
- Assessments for calibrated segments having tentative data consisting of at least one valid BMIBI score and one valid FIBI score, and both of the following are true:
 - 1) the single score(s) plus the applicable UAV(s) is less than the BIC or the average(s) of multiple scores is less than the BIC,
 - 2) the single score or the average of multiple scores for both the BMIBI and the FIBI fall in the qualitative range indicating poor biocondition.

Abbreviations and terms: **ALUS**, Aquatic Life Use Support; **BIC**, Biological Impairment Criteria/Criterion; **BMIBI**, Benthic Macroinvertebrate Index of Biotic Integrity; **FIBI**, Fish Index of Biotic Integrity; **UAV**, Uncertainty Adjustment Value [8 pts. BMIBI, 7 pts. FIBI]

Calibrated - Stream segments designated as B(LR) or B(WW) in 2004 and have a watershed drainage area < 500 square miles.

Uncalibrated - General use segments or non-wadeable river segments having watershed drainage area \geq 500 mi².

Comparable - Data considered as having completeness and quality that is comparable to biocriteria project data used to develop reference biotic indexes and impairment criteria.

Tentative - Data considered as having lesser or uncertain levels of completeness and quality documentation.

Table 2-1(a). BMIBI qualitative scoring ranges.

Biological Condition Rating	Characteristics of Benthic Macroinvertebrate Assemblage
76-100 (Excellent)	High numbers of taxa are present, including many sensitive species. EPT taxa are very diverse and dominate the benthic macroinvertebrate assemblage in terms of abundance. Habitat and trophic specialists, such as scraper organisms, are present in good numbers. All major functional feeding groups (ffg) are represented, and no particular ffg is excessively dominant. The assemblage is diverse and reasonably balanced with respect to the abundance of each taxon.
56-75 (Good)	Taxa richness is slightly reduced from optimum levels; however, good numbers of taxa are present, including several sensitive species. EPT taxa are fairly diverse and numerically dominate the assemblage. The most-sensitive taxa and some habitat specialists may be reduced in abundance or absent. The assemblage is reasonably balanced, with no taxon excessively dominant. One ffg, often collector-filterers or collector-gatherers, may be somewhat dominant over other ffgs.
31-55 (Fair)	Levels of total taxa richness and EPT taxa richness are noticeably reduced from optimum levels; sensitive species and habitat specialists are rare; EPT taxa still may be dominant in abundance; however, the most-sensitive EPT taxa have been replaced by more-tolerant EPT taxa. The assemblage is not balanced; just a few taxa contribute to the majority of organisms. Collector-filterers or collector-gatherers often comprise more than 50% of the assemblage; representation among other ffgs is low or absent.
0-30 (Poor)	Total taxa richness and EPT taxa richness are low. Sensitive species and habitat specialists are rare or absent. EPT taxa are no longer numerically dominant. A few tolerant organisms typically dominate the assemblage. Trophic structure is unbalanced; collector-filterers or collector-gatherers are often excessively dominant; usually some ffgs are not represented. Abundance of organisms is often low.

Table 2-1(b). Fish Index of Biotic Integrity (FIBI) qualitative scoring guidelines.

<p>71-100 (Excellent)</p>	<p>Fish (excluding tolerant species) are fairly abundant or abundant. A high number of native species are present, including many long-lived, habitat specialist, and sensitive species. Sensitive fish species and species of intermediate pollution tolerance are numerically-dominant. The three most abundant fish species typically comprise 50% or less of the total number of fish. Top carnivores are usually present in appropriate numbers and multiple life stages. Habitat specialists, such as benthic invertivore and simple lithophilous spawning fish are present at near optimal levels. Fish condition is good; typically less than 1% of the total number of fish exhibit external anomalies associated with disease or stress.</p>
<p>51-70 (Good)</p>	<p>Fish (excluding tolerant species) are fairly abundant to very abundant. If high numbers are present, intermediately tolerant species or tolerant species are usually dominant. A moderately high number of fish species belonging to several families are present. The three most abundant fish species typically comprise two-thirds or less of the total number of fish. Several long-lived species and benthic invertivore species are present. One to several sensitive species are usually present. Top carnivore species are usually present in low numbers and often one or more life stages is missing. Species that require silt-free, rock substrate for spawning or feeding are present in low proportion to the total number of fish. Fish condition is good; typically less than 1% of the total number of fish exhibit external anomalies associated with disease or stress.</p>
<p>26-50 (Fair)</p>	<p>Fish abundance ranges from lower than average to very abundant. If fish are abundant, tolerant species are usually dominant. Native fish species usually equal ten or more species. The three most abundant species typically comprise two-thirds or more of the total number of fish. One or more sensitive species, long-lived fish species or benthic habitat specialists such as Catostomids (suckers) are present. Top carnivore species are often, but not always present in low abundance. Species that are able to utilize a wide range of food items including plant, animal and detrital matter are usually more common than specialized feeders, such as benthic invertivore fish. Species that require silt-free, rock substrate for spawning or feeding are typically rare or absent. Fish condition is usually good; however, elevated levels of fish exhibiting external anomalies associated with disease or stress are not unusual.</p>
<p>0-25 (Poor)</p>	<p>Fish abundance is usually lower than normal or, if fish are abundant, the assemblage is dominated by a few or less tolerant species. The number of native fish species present is low. Sensitive species and habitat specialists are absent or extremely rare. The fish assemblage is dominated by just a few ubiquitous species that are tolerant of wide-ranging water quality and habitat conditions. Pioneering species, introduced species, and short-lived fish species are typically the most abundant types of fish. Elevated levels of fish with external physical anomalies are more likely to occur.</p>

Table 2-2. Biological Impairment Criteria (BIC) used for the assessment of warmwater rivers and streams in the 2006 section 305(b) reporting and section 303(d) listing cycles. For a discussion of how the BIC were derived, please see the addendum to this Attachment.

Ecoregion:	FIBI	BMIBI
40a – Central Irregular Plains	33	41
47 – Western Corn Belt Plains (WCBP) Subregions:		
47(a) – WCBP /Northwest Iowa Loess Prairies	43	54
47(b) – WCBP / Des Moines Lobe (Stable Riffle Habitat*)	53	62
(No Stable Riffle Habitat)	32	62
47(c) – WCBP / Iowan Surface (Stable Riffle Habitat)	65	70
(No Stable Riffle Habitat)	44	52
47(d) – WCBP / Missouri Alluvial Plain	-	-
47(e) – WCBP / Loess Hills and Rolling Loess Prairies	31	54
47(f) – WCBP / Southern Iowa Rolling Loess Prairies (Mississippi Drainage System)	36	51
(Missouri Drainage System)	31	54
52b – Paleozoic Plateau (Driftless Area)	52	61
72d – Central Interior Lowland	-	-

Determining Support of B(CW) [coldwater] Aquatic Life Uses

Nine coldwater streams where biocriteria sampling was done from 1994-1998 were used to establish criteria used to determine the status of Class B(CW) aquatic life use. Eight biological indicators that reflect coldwater stream water quality and habitat suitability were calculated, and a ranking system was used to determine the level of B(CW) use support.

Coldwater stream biological indicators used to determine B(CW) aquatic life use status.

1. Number of sensitive benthic macroinvertebrate taxa.
2. Number of coldwater obligate benthic macroinvertebrate taxa
3. Benthic macroinvertebrate biotic index of organic enrichment.
4. Percent dominance of three most abundant benthic macroinvertebrates.
5. Number of coldwater fish species.
6. Percent abundance of coldwater fish species
7. Presence/absence of trout.
8. Trout reproduction rating for stream.

The degree of B(CW) use support for a given stream site was assessed by determining the number of biological indicator values that ranked below the 25th percentile of indicator values from all nine coldwater stream sampling sites. Sites with ≤ 2 indicators ranking below the 25th percentile level are assessed as fully supporting or fully supporting/threatened (=FS or FS/T); sites with 2-4 indicators ranking below the 25th percentile level are assessed as fully supporting/threatened (=FS/T); sites with 5 or 6 indicators below the 25th percentile level are assessed as partially supporting (=PS); sites with 7 or 8 indicators below the 25th percentile level are assessed as not supporting (=NS).

II. Applying the site assessment results to a Section 305(b) stream segment.

- a) Stream segment assessments derived from a single sampling event. When data from one sampling event at one sampling site are the only data available, the assessment result for that site (e.g., fully supporting/threatened) is applied to the entire stream segment length. Most of the stream segments assessed for Section 305(b) reporting with results of 1997-2002 biocriteria sampling belong to this category.

- b) Stream segments with multiple sampling sites. Relatively few stream segments have data from multiple biological sampling sites, and these are examined on a case-by-case basis. In general, when data from multiple sites are available, the lowest assessment result is assigned to the entire stream segment length. For example, if one site assessment result indicates aquatic life use is partially supporting and a second site assessment result is fully supporting/threatened uses, the partially supporting assessment is applied to the entire stream segment. One exception of this is when one or more sites are judged to be unrepresentative of the stream segment as a whole (e.g., mixing zone of wastewater discharge). In this case, only the assessment results from the site or sites that are considered representative are used to make the assessment for the entire stream segment.

III. Identifying causes and sources of impairment.

As defined in guidelines for Section 305(b) reporting (U.S. EPA 1997), causes of water quality impairment are those pollutants and environmental stressors that contribute to the impairment of designated uses in a waterbody. Sources are the activities, facilities or conditions that contribute the pollutants and environmental stressors which result in the impairment of designated beneficial uses. For example, high levels of pesticides (the *cause*) from agricultural activities (the *source*) can impair a waterbody's designated beneficial uses as a source of drinking water.

Causes and sources of impairment are specified for stream segments assessed as either "partially supporting" or "not supporting" aquatic life uses. DNR Watershed Monitoring & Assessment Section staff follow U.S. EPA guidelines and use best professional judgment to identify and assign a magnitude to each cause and source of impairment. DNR staff consider available information about pollution sources and recent events affecting water quality. Summary information from stream physical habitat evaluations are also used to assess causes and sources that are related to habitat alterations. The information reviewed includes floodplain land uses, buffer strip width and vegetation, channel sinuosity and morphometry, bank conditions, sediment composition, stream flow, and instream habitat.

References for Attachment 2:

- Griffith, G.E., J.M. Omernik, T.F. Wilton, and S.M. Pierson. 1994. Ecoregions and subcoregions of Iowa: a framework for water quality assessment and management. *Journal of the Iowa Academy of Science*. 10(1):5-13.
- IAC. 2002. Chapter 567-61: water quality standards. Iowa Administrative Code [effective date 08/14/2002].
- Mundahl, N.D. and T.P. Simon. 1999. Chapter 15:383-416. Development and application of an index of biotic integrity for coldwater streams of the upper Midwestern United States. In Assessing the Sustainability and Biological Integrity of Water Resources Using Fish Communities. CRC Press LLC.
- Omernik, J.M., G.E. Griffith, and S.M. Pierson. 1993. Ecoregions and western cornbelt plains subregions of Iowa. U.S. Environmental Protection Agency, Environmental Research Laboratory, Corvallis, Oregon. 29 p.
- U.S. EPA. 1990. Biological criteria – national program guidance for surface waters. EPA-440/5/5-90-004. U.S. Environmental Protection Agency, Office of Water Regulations and Standards. Washington D.C.
- U.S. EPA 1996. Biological criteria: technical guidance for streams and small rivers. revised edition. EPA-822-B-96-001. U.S. Environmental Protection Agency, Office of Water, Washington, D.C. 162p.
- U.S. EPA. 1997. Guidelines for the preparation of the comprehensive state water quality assessments (305(b) reports) and electronic updates. Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency, Washington, D.C.
- Wilton, T.F. 1996. Pilot study of biocriteria data collection procedures for wadeable streams in Iowa: final report. Iowa Department of Natural Resources, Environmental Protection Division, Water Resources Section. 18p.
- Wilton, T.F. 2004. Biological assessment of Iowa's wadeable streams. Iowa Department of Natural Resources, Environmental Services Division, TMDL and Water Quality Assessment Section. Des Moines, Iowa. 267p.
- Yoder, C.O., and E.T. Rankin. 1995. Chapter 9:109-144. Biological criteria program development and implementation in Ohio. In Biological assessment and criteria: tools for water resources planning. W.S. Davis and T.P. Simon, editors. CRC Press, Inc

Addendum to Attachment 2:

Establishment of Biological Impairment Criteria (BIC) for Determining Support of Warmwater Stream Aquatic Life Designated Uses

September 2007

Introduction

This document describes the rationale, procedures, and results from the recalculation of Biological Assessment Criteria (BIC) used in the 2006 biennial 305(b)/303(d) Integrated Report. Supplemental information describing sampling protocols, biotic index development, ecoregions and reference sites can be found in the IDNR stream bioassessment project report (Wilton 2004). Procedures for determining the support status of designated aquatic life uses are described in the 305(b)/303(d) Integrated Report assessment methodology (IDNR 2007). With minor modifications, the existing bioassessment framework has been used for 305(b) reporting and 303(d) impaired waters listings since the 2000 assessment cycle.

To determine the support status of warmwater stream aquatic life uses, the Benthic Macroinvertebrate Index of Biotic Integrity (BMIBI) and the Fish Index of Biotic Integrity (FIBI) sample scores from a given segment are compared to applicable Biological Impairment Criteria (BIC). The BIC are statistically derived from index scores sampled at reference stream sites located in the same ecological region (Figure 1). Reference sites are chosen to represent least disturbed stream habitats that support healthy biological aquatic communities. Reference data have been used to define best aquatic life use expectations through calibration of the BMIBI and FIBI and establishment of Biological Assessment Criteria (BIC). Wadeable stream reference sites are generally sampled in a five-year rotational schedule. The first cycle of reference site sampling was conducted from 1994-1998. A few additional reference sites were sampled in 1999, and the second cycle of reference site sampling was conducted from 2000-2004.

BIC Re-calculation Rationale

Since the reference site network and bioassessment approach is relatively new in Iowa, it was reasoned that the inclusion of recent data would help insure that reference biological conditions appropriately reflect a full range of climatic and hydrologic conditions affecting stream aquatic communities. Given the cyclic nature of drought and wet years in the Midwest, it was believed that averaging of reference sampling results from a decade of sampling (1994-2004) would more appropriately reflect the natural variations in stream biological conditions and provide a robust data set for stream biological assessment. Therefore, additional results of reference site sampling from 2002-2004 were added to the previous data set of 1994-2001 data in order to update the BIC (Table 1) for the 2006 listing cycle. The additional data, in most cases, increased the number of BMIBI and FIBI samples representing each reference site from one to two samples.

Methods

A consistent approach was followed in calculating the BIC for the 2006 and previous assessment cycles. Currently, 95 reference sites are recognized by the IDNR for stream bioassessment purposes. Only data from the 87 warmwater reference sites were used to calculate the BIC. Data from eight coldwater

reference sites were disregarded. Approximately 180 valid BMIBI and FIBI scores obtained during the normal July – October sampling index period were included in the BIC calculations. The respective BMIBI and FIBI scores from each site were averaged and the site averages were compiled by ecoregion. Statistical summaries of average reference site IBI scores are reported in Table 2 and Table 3. In response to previous findings (Wilton 2004), additional statistical tests were performed to examine for differences between habitat and benthic sampling gear groupings within certain ecoregions (Tables 4-6).

IDNR has chosen the 25th percentile values of the reference site BMIBI and FIBI index scores within a given ecoregion or habitat class to represent the biological impairment criteria (BIC) for 305(b)/303(d) biological assessment purposes (Table 1). Use of the reference 25th percentile as an impairment threshold is consistent with biocriteria development guidance (U.S. EPA 1996), and has demonstrated efficacy in state bioassessment programs (Yoder and Rankin 1995). Evaluation of biotic index performance in Iowa found little or no overlap of index interquartile ranges between reference sites and test (impacted) sites, which suggests that reference 25th percentile levels are appropriate for assessing biological impairment (Wilton 2004).

Recalculation Results

Table 1 provides a comparison of the BIC used in the 2006 IR assessment with the BIC used in the 2002 and 2004 assessment cycles. For the BMIBI, two BIC were raised, one was kept equal, and seven BIC were lowered in relation to the 2002/2004 BIC. Separate BIC were established by sampling gear type within ecoregion 47c after statistical analysis found a significant difference in BMIBI scores among sites sampled using the Hess sampling device (riffle habitat) versus sites sampled using Hester-Dendy artificial substrates (Table 4; rank sum test $p < 0.05$). This separation resulted in both the largest BIC increase (11 points; Hess sites) and the largest decrease (7 points, Artificial Substrate sites) from the 2002/2004 BIC.

For the FIBI, four of the 2006 BIC were raised, four were kept equal, and four were lowered in relation to the 2002/2004 BIC. The largest BIC increase was 3 points (47a) and the largest decrease was 7 points (52b). Riffle and non-riffle sites within ecoregion 47f were combined to calculate a single BIC after statistical testing failed to show a difference in FIBI scores among these groups (Addendum Table 6; rank sum test $p > 0.05$).

While most of the changes in BIC were small, more of them were lowered than raised or kept the same. This trend has prompted follow-up examination of trends in reference site sampling data. For example, it was determined that approximately 60% of reference sites had higher BMIBI or FIBI scores from the 1994-1998 period (cycle 1) compared with scores from the 2000-2004 sampling period (cycle 2). Mean site paired differences (cycle 1- cycle 2) of 3.9 points for the BMIBI and 4.1 points for the FIBI were both significantly greater than zero (paired t-test, $p < 0.05$), thus indicating an overall decline in BMIBI and FIBI scores. This trend is cause for concern that reference conditions might be deteriorating, and simultaneously points out the value of sustained long-term monitoring projects.

The IDNR bioassessment unit has initiated an investigation of factors that may have contributed to the observed trend. Significant year-to-year differences in the magnitude of changes in IBI levels have been observed (Figure 2) suggesting that climatic variation is a potential contributing factor. Precipitation patterns, for example, can influence the flow regime, habitat and water quality conditions under which the aquatic communities develop. A correlation analysis found the largest changes in FIBI scores between sample cycles 1 and 2 were associated with the largest differences in sample date flow. This relationship might reflect differences in fish distribution or sampling effectiveness that occur under different flow regimes. Additional exploratory analysis found a lack of relationship between the direction or size of reference site changes in BMIBI scores and changes in FIBI scores (Figure 3), which might indicate the two indexes respond to environmental conditions at different spatial and/or temporal scales.

Ecoregion or stream watershed size also were not related with the direction or size of changes in IBI levels. The bioassessment unit is not currently aware of any widespread changes in land use or anthropogenic stressors in reference site watersheds that might explain the declining trend, but will continue to investigate this possibility.

Future Outlook

IDNR considers the development and verification of reference conditions to be an evolving process. Reference sites and reference conditions for bioassessment are the subject of significant research and development work throughout the United States. IDNR will continue to improve its reference condition development process and will utilize new techniques and methods as they become available.

As new data from reference sites is obtained, it will be reviewed and incorporated in each successive biennial Integrated Report. When the next cycle of warmwater reference site sampling is completed, IDNR will again review and update the BIC, if needed. At that time, there will be a minimum of three samples from each reference site covering approximately seventeen years of sampling. Other data, particularly the 2002-2006 (REMAP) random survey of perennial streams will be reviewed to determine whether additional reference sites can be gleaned and the data used to better define reference conditions and BIC. Although no specific timeframe has been set, it is anticipated that Iowa's wadeable stream bioassessment framework and BIC will be reviewed for potential incorporation within Iowa's water quality standards.

References

- Chapman, S.S., J.M. Omernik, G.E. Griffith, W.A. Schroeder, and T.F. Wilton. 2002. Ecoregions of Iowa and Missouri (color poster with map, descriptive text, summary tables, and photographs): Reston, Virginia, U.S. Geological Survey (map scale 1:1,800,000).
- IDNR 2007. (DRAFT) Methodology for Iowa's 2006 water quality assessment, listing, and reporting pursuant to Sections 305(b) and 303(d) of the federal Clean Water Act. April 2007 Watershed Monitoring & Assessment Section, Geological Survey & Land Quality Bureau, Environmental Services Division, Iowa Department of Natural Resources.
- U.S. EPA 1996. Biological criteria: technical guidance for streams and small rivers. Revised edition. EPA-822-B-96-001. U.S. Environmental Protection Agency, Office of Water, Washington, D.C. 162p.
- Wilton, T.F. 2004. Biological assessment of Iowa's wadeable streams. Iowa Department of Natural Resources, Environmental Services Division, TMDL and Water Quality Assessment Section. Des Moines, Iowa. 267p.
- Yoder, C.O., and E.T. Rankin. 1995. Chapter 9:109-144. Biological criteria program development and implementation in Ohio. In Biological assessment and criteria: tools for water resources planning. W.S. Davis and T.P. Simon, editors. CRC Press, Inc

Addendum Table 1. Warmwater stream Biological Impairment Criteria (BIC) for 305(b)/303(d) Integrated Report biological assessments.

Ecoregion	Major Drainage	Riffle?	FIBI BIC '06 ('02-'04)	Bug gear	BMIBI BIC '06 ('02-'04)
40a	All	All	33 (33)	All	41 (46)
47a	All	All	43 (40)	All	54 (53)
47b	All	Yes	53 (55)	All	62 (63)
47b	All	No	32 (32)	All	62 (63)
47c	All	Yes	65 (71)	Hess	70 (59)
47c	All	No	44 (43)	Art. Sub.	52 (59)
47e	All	All	31 (31)	All	54 (56)
47f	MSP	All	36 (41,34)*	All	51 (53)
47f	MO	All	31 (31)	All	54 (56)
52b	All	All	52 (59)	All	61 (61)
72d	All	All	36 (34)	All	51 (53)

* '02-'04 47f BIC: MSP riffle = 41 and MSP non-riffle = 34.

Addendum Table 2. Summary statistics for 1994-2004 warmwater wadeable stream reference site FIBI scores by ecoregion.

Ecoregion	# sites	FIBI mean	FIBI min	FIBI 25th	FIBI median	FIBI 75th	FIBI max
40a	7	40.9	27.0	33.0	37.5	50.0	57.0
47a	6	46.2	42.5	42.9	46.3	49.3	50.0
47b all	20	50.8	28.5	38.3	51.3	61.0	74.5
47b riffle	11	58.8	37.0	52.5	60.0	71.5	74.5
47b non-riffle	9	41.0	28.5	31.8	42.0	48.5	55.5
47c all	20	62.8	38.0	50.2	64.8	76.4	83.0
47c riffle	8	73.1	58.5	64.9	76.6	78.9	83
47c non-riffle	12	55.9	38	44.1	54.3	69.1	76.5
47e	8	36.0	25.5	30.9	37.0	37.9	49.5
47f	17	46.7	23.5	35.5	48.5	54.5	71.0
52b	7	64.9	48.0	52.0	63.5	79.0	81.0
72d	2	45.2	43.0	44.1	45.2	46.3	47.3

Addendum Table 3. Summary statistics for 1994-2004 reference site BMIBI scores by ecoregion.

Ecoregion	# sites	BMIBI mean	BMIBI min	BMIBI 25th	BMIBI median	BMIBI 75th	BMIBI max
40a	7	48.7	34.0	41.0	50.0	56.5	68.0
47a	6	66.5	50.0	53.8	65.5	78.3	88.0
47b	20	65.6	37.5	62.0	68.9	73.4	76.5
47c art subs	9	58.7	47.0	52.3	59.5	65.2	70.5
47c hess/surber	13	73.3	62.0	70.3	72.6	78.0	81.5
47c all sites	20	67.2	47.0	60.6	69.3	73.3	81.5
47e	8	58.7	46.0	53.4	57.5	66.3	70.0
47f	17	59.2	44.0	50.3	62.5	66.8	71.0
52b	7	67.9	54.5	61.0	68.0	75.0	80.5
72d	2	43.8	39.0	41.4	43.8	46.1	48.5

Addendum Table 4. Statistical analysis of 1994-2004 reference site BMIBI scores from select ecoregions by benthic macroinvertebrate sampling gear.


Ecoregion	# sites	BM-IBI mean	BM-IBI min	BM-IBI 25th	BM-IBI median	BM-IBI 75th	BM-IBI max	BM-IBI two sample mean TTest p-value	BM-IBI rank sum p-value
40a art subs	3	38.3	31.0	31.0	34.0	50.0	50.0	0.1186	0.1213
40a hess/surber	6	51.2	41.0	41.0	50.3	59.4	68.0		
47b art subs	9	59.6	37.5	45.0	58.0	74.0	75.0	0.0380	0.1756
47b hess/surber	14	70.0	61.0	63.7	69.4	74.0	76.5		
47c art subs	9	58.7	47.0	52.3	59.5	65.2	70.5	0.00005	0.0005
47c hess/surber	13	73.3	62.0	70.3	72.6	78.0	81.5		
47e art subs	3	58.7	46.0	46.0	62.5	67.5	67.5	0.9966	1.0000
47e hess/surber	5	58.7	52.5	54.3	56.5	64.3	70.0		
47f art subs	6	60.3	44.0	48.5	62.8	71.0	71.0	0.8107	0.7079
47f hess/surber	12	60.0	45.5	51.4	59.5	66.9	69.7		

Addendum Table 5. Statistical analysis of 1994-2004 reference site BMIBI scores from select ecoregions by stream type: riffle or non-riffle. Riffle streams include >10% riffle macrohabitat, >10% cobble substrate and >30% total coarse substrate.

Ecoregion	# sites	BMIBI mean	BMIBI min	BMIBI 25th	BMIBI median	BMIBI 75th	BMIBI max	BMIBI two sample TTest mean p-value	BMIBI rank sum p-value
47b riffle	11	68.9	62.0	64.2	67.3	74	76.5	0.1138	0.2875
47b non-riffle	9	61.5	37.5	49.0	70.5	72.0	75.0		
47c riffle	8	74.4	62.0	72.5	74.5	79.5	81.5	0.0024	0.0014
47c non-riffle	12	62.3	47.0	56.1	63.9	69.4	71.5		
47f riffle	9	59.6	45.5	50.0	62.0	68.0	69.7	0.7650	0.8323
47f non-riffle	7	58.2	44.0	50.0	62.5	63.0	71.0		


Addendum Table 6. Statistical analysis of 1994-2004 reference site FIBI scores from select ecoregions by stream type: riffle or non-riffle. Riffle streams include >10% riffle macrohabitat, >10% cobble substrate and >30% total coarse substrate.

Ecoregion	# sites	FIBI mean	FIBI min	FIBI 25th	FIBI median	FIBI 75th	FIBI max	FIBI two sample mean TTest p-value	FIBI rank sum p-value
47b riffle	11	58.8	37.0	52.5	60.0	71.5	74.5	0.0018	0.0044
47b non-riffle	9	41.0	28.5	31.8	42.0	48.5	55.5		
47c riffle	8	73.1	58.5	64.9	76.6	78.9	83.0	0.0049	0.0062
47c non-riffle	12	55.9	38.0	44.1	54.3	69.1	76.5		
47f riffle	9	49.9	34.0	36.5	51.5	61.0	71.0	0.5399	0.6720
47f non-riffle	7	45.9	25.0	36.0	48.5	53.0	62.0		




- 40(a) - Central Irregular Plains (Loess Flats and Till Plains)
47(a) - Western Corn Belt Plains (Northwest Iowa Loess Prairies) 47(b) - WCBP (Des Moines Lobe)
47(c) - WCBP (Iowan Surface) 47(d) - WCBP (Missouri Alluvial Plain) 47(e) - WCBP (Steeply Rolling Loess Prairies)
47(f) - WCBP (Rolling Loess Prairies) 47(m) - WCBP (Western Loess Hills)
52(b) - Driftless Area (Paleozoic Plateau) 72(d) - Central Interior Lowland (Upper Mississippi Alluvial Plain)

Addendum Figure 1. Ecological regions of Iowa (after Chapman et al. 2002).



Addendum Figure 2. Reference site paired differences of first IBI sample minus second IBI sample. Sample years indicate the years of the first IBI sample and the second IBI sample.



Addendum Figure 3. Reference site paired differences of first IBI sample (1994-1998) minus second IBI sample (1999-2004). Site symbols correspond with the ecoregion in which the site is located.

Attachment 3

**THE USE OF THE TROPHIC STATE INDEX TO IDENTIFY
WATER QUALITY IMPAIRMENTS IN IOWA LAKES FOR THE 2010
SECTION 305(b) REPORTING AND SECTION 303(d) LISTING CYCLES**

Iowa DNR
Watershed Monitoring & Assessment Section
Geological and Water Survey

Updated, April 2011

Table of Contents	Page
Introduction	109
Assessment rationale	110
Identifying water quality impairments at Iowa lakes based on TSI	111
Relevant state water quality criteria	112
Data sources	113
Data requirements for listing	113
Data quantity	113
Data quality	114
Threshold TSI values	114
Assessment categories ("monitored" and "evaluated")	114
Use support categories	115
Management and accessibility of assessments	117
Table 3-1. Changes in temperate lake attributes according to trophic state.	118
Table 3-2. Iowa lakes with TSI values for total phosphorus greater than 70 but with TSIs for chlorophyll-a and Secchi depth of less than 65.	118
Table 3-3. Ranges of TSI values for Iowa lakes.	119
Table 3-4. Summary of ranges of TSI values and measurements for chlorophyll-a and Secchi depth used to define Section 305(b) use support categories.	119
Table 3-5. Narrative descriptions of TSI ranges for Secchi depth, phosphorus, and chlorophyll-a for Iowa lakes.	120
References for Attachment 3.	121

INTRODUCTION

Prior to 2000, relatively little water quality monitoring was conducted on Iowa lakes. Lake surveys in Iowa typically involved sampling in only summer seasons of one year at roughly ten-year intervals (see Bachmann 1965, Bachmann et al. 1980, and Bachmann et al. 1994). This amount of data, although providing a snapshot of lake water quality given the climatic conditions of the specific year of sampling, was not particularly useful for developing a more accurate characterization of lake-specific water quality over the long-term. In addition, due to the general lack of historical data, accurate identification of long-term trends in water quality parameters at most Iowa lakes was not possible. Diagnostic/feasibility studies at Iowa lakes (e.g., Bachmann et al. 1982, Downing et al. 2001), have included more intensive water quality monitoring, but such studies have been conducted on relatively few lakes and are of a relatively short duration (from one to two years). Due to this general lack of data, historical assessments of lake water quality in Iowa, such as those used for Section 305(b) reporting and Section 303(d) listing, had been based primarily on the best professional judgment of Iowa DNR fisheries biologists. The nearly total reliance on best professional judgment, while a valid assessment technique, resulted not only from the lack of routine ambient monitoring at Iowa lakes but also from the lack of state water quality criteria for the parameters that are most likely to indicate lake water quality impairments (e.g., nutrients (nitrogen and phosphorus), chlorophyll, turbidity, and impacts due to the accumulation of sediment in lake basins). Previous (pre-2000) Section 305(b) lake assessments that were based on best professional judgment were supplemented with lake monitoring data as this information was available (e.g., Bachmann et al. 1982, Bachmann et al. 1994).

Beginning in 2000, however, the first routine ambient monitoring program for Iowa lakes was initiated. This statewide lake survey of 131 publicly-owned Iowa lakes was funded by Iowa DNR and was conducted by ISU from 2000 through 2007 and by the State Hygienic Laboratory at the University of Iowa (SHL) from 2005-2008. This study was designed to be a long-term study capable of providing multiple years of data that can be used to better characterize lake water quality than was possible with the limited data from previous surveys.

This lake assessment methodology for Iowa's 2010 integrated (305(b)/303(d)) report involves the use of data from the statewide lake surveys conducted by ISU and the SHL from 2002 through 2008 with Carlson's (1977) trophic state index (TSI) to identify lakes that do not fully meet the narrative criteria in Section 61.3(2) of the *Iowa Water Quality Standards*. This general approach was used for Iowa's 2002, 2004, 2006, and 2008 reporting/listing cycles as well. The existence of any lake impairments suggested by a TSI value will be corroborated by IDNR field (Fisheries Bureau) staff. This approach is consistent with Iowa's credible data law and allows assessment of water quality impacts due to parameters that currently lack numeric criteria in the *Iowa Water Quality Standards*. The use of TSI values for chlorophyll and Secchi depth serves as an interim method of assessing lake water quality in Iowa until numeric criteria for nutrient parameters (phosphorus and nitrogen) and their response variables (chlorophyll-a and turbidity) are adopted into the *Iowa Water Quality Standards*.

ASSESSMENT RATIONALE

The concept of “trophic state” has long been used by limnologists to classify lakes and is based on the chemistry and biology of lakes. Although a number of approaches exist for classifying lakes according to trophic state, and although a number of controversies exist regarding how “trophic state” is defined, the use of this framework has the advantages of historical usage, general acceptance of the trophic state concept (e.g., “eutrophic” indicates nutrient enrichment), and an improved ability to describe lake condition versus a description using a single variable or number (e.g., total phosphorus concentration). Table 3-1 describes the general framework of the lake trophic state concept. For a discussion on the development and variety of trophic state indices, see Chapter 2 (*The Basis for Lake and Reservoir Nutrient Criteria*) in U.S. EPA (2000) (see <http://www.epa.gov/waterscience/criteria/nutrient/guidance/lakes/lakes.pdf>).

Carlson's (1977) trophic state index is a numeric indicator of the continuum of the biomass of suspended algae in lakes and thus reflects a lake's nutrient condition and water transparency. The level of plant biomass is estimated by calculating the TSI value for chlorophyll-a. TSI values for total phosphorus and Secchi depth serve as surrogate measures of the TSI value for chlorophyll. The focus on turbidity in general, and chlorophyll in particular, seems appropriate for assessing the degree to which Iowa lakes support their designated Class A1 (primary contact recreation) uses. Carlson's trophic state index provides a convenient and well-established method for identifying turbidity-related impacts to Iowa lakes. As described in a subsequent paper by Carlson (1991), turbidity, and especially turbidity related to large populations of suspended algae, is a key indicator of the degree to which a lake supports primary contact uses:

[plant] biomass is a proximate measure of the problems that plague lakes. Probably few citizens complain about the productivity of their lake and fewer yet lodge complaints about phosphorus concentrations. A biomass-related trophic state definition places the emphasis of the classification on the problem rather than on any potential cause.

Because of this direct linkage between the perceived level of water quality and turbidity, TSI values for chlorophyll-a and Secchi depth will be used as guidelines to identify Iowa lakes that do not meet Iowa's narrative water quality standards protecting against “aesthetically objectionable conditions”. Both chlorophyll-a and Secchi depth appear applicable to Iowa's narrative water quality criterion protecting against aesthetically objectionable conditions in Iowa surface waters (IAC 2008, 61.3(2)). IDNR field (Fisheries Bureau) staff will be contacted to corroborate that the aesthetically objectionable conditions suggested by the TSI values do, in fact, exist. Because aesthetics are more closely associated with recreational uses than to aquatic life uses of Iowa lakes, impairments based on violations of these narrative criteria are typically applied to Class A1 (primary contact recreation) uses.

For two reasons, TSI values for total phosphorus are not used as the primary basis for assessing support of either primary contact recreation uses or aquatic life uses:

1. *TSI's for total phosphorus are poor predictors of impairment due to either Secchi depth or chlorophyll-a:* The typical use of the TSI for total phosphorus to measure trophic state (and the level of water quality) presumes that the

relationship between total phosphorus and chlorophyll-a will, more or less, hold for the lake being assessed. The production of chlorophyll in Iowa's natural lakes and impoundments, however, is sometimes limited by nutrients other than phosphorus (e.g., nitrogen) and/or high levels of non-algal turbidity in the water column. The result is that lakes with very high levels of total phosphorus that suggest hypereutrophic conditions sometimes have levels of chlorophyll-a and Secchi depth that suggest relatively good water quality (i.e., in the middle to lower eutrophic range). As examples, the Iowa lakes in Table 3-2 are those that have TSI values for total phosphorus in the hypereutrophic range (i.e., greater than 70) but that have TSI values for chlorophyll-a and Secchi depth less than the impairment trigger of TSI=65. Examples of lakes in Iowa with relatively high TSI values for total phosphorus but low values for chlorophyll-a and Secchi depth include West Lake Osceola (Clarke County), Center Lake (Dickinson County), and Red Rock Reservoir (Marion County). Thus, while these lakes have very high levels of total phosphorus that might suggest impairment of designated uses, the levels of chlorophyll-a and Secchi depth are relatively low and do not suggest impairment. Because of this lack of correlation between TSI values for total phosphorus and TSI values for the response variables that define the aesthetically objectionable conditions, TSI values for total phosphorus are not used as the primary basis for determining the level of use support or for identifying water quality impairments at Iowa lakes.

2. The Iowa Water Quality Standards lack water quality criteria—narrative or numeric—that are relevant to impacts of total phosphorus in surface waters. When developing this assessment procedure, careful consideration of Iowa's numeric and narrative criteria in the *Iowa Water Quality Standards* showed that none of these criteria are directly relevant to levels of phosphorus in the water column of a lake. That is, phosphorus is not a toxic substance at ambient levels seen in Iowa waters. In addition, high levels of phosphorus in Iowa lakes do not necessarily lead to either nuisance aquatic life or aesthetically objectionable conditions. For example, lakes with growths of aquatic macrophytes in littoral zone areas can have high levels of phosphorus but have low levels of chlorophyll-a and have good water transparency.

For lakes where assessment information from the IDNR Fisheries Bureau is available, TSI values were also used to supplement assessments of the designated Class B aquatic life uses based on best professional judgment of IDNR fisheries biologists. According to biologists in the IDNR Fisheries Bureau, algal blooms can also cause impairments to aquatic life uses of Iowa lakes through interference with some spawning activities of nest building species, e.g., bluegill, bullhead, crappie and largemouth bass) and lowered levels (sags) of dissolved oxygen that, in extreme cases, can cause fish mortality.

IDENTIFYING WATER QUALITY IMPAIRMENTS AT IOWA LAKES BASED ON TSI:

For purposes of developing water quality assessments for the 2010 Section 305(b) reporting cycle, Carlson's (1977, 1984, 1991) "trophic state index" (TSI) values were calculated using the data generated from 131 Iowa lakes as part of ISU and SHL surveys from 2004 through 2008. Overall (five-year) median values were used to calculate TSI values for total phosphorus, chlorophyll-a, and Secchi depth for each lake;

the ranges of these values are summarized in Table 3-3. The identification of an impairment of the primary contact uses was based on TSI values for chlorophyll-a and/or Secchi depth. The TSI values for the indicator variable of total phosphorus are used primarily to interpret discrepancies between TSI values for chlorophyll-a and Secchi depth.

Relevant state water quality criteria:

The *Iowa Water Quality Standards* (IAC 2008) do not contain numeric criteria for nutrients (e.g., nitrogen or phosphorus), chlorophyll, or turbidity that apply to Class A1 uses. Thus, the assessments of the degree to which these parameters might impair the Class A1 uses are based on a comparison of lake-specific TSI values to the following narrative criteria for general use waters as defined in Section 61.3(2) of the *Iowa Water Quality Standards*:

Such waters shall be free from materials attributable to wastewater discharges or agricultural practices producing objectionable color, odor, or other aesthetically objectionable conditions.

Such waters shall be free from substances, attributable to wastewater discharges or agricultural practices, in quantities which would produce undesirable or nuisance aquatic life;

Examples of *aesthetically objectionable conditions* include poor water transparency caused by blooms of algae or high levels of non-algal turbidity that make the lake less desirable (aesthetically unpleasing) for primary contact recreation. Cyanobacteria blooms can also cause *aesthetically objectionable conditions* due to their ability to create unpleasant floating scums on the water surface or unpleasant odors, both of which can limit the primary contact recreation uses at a lake. In addition, cyanobacteria can be considered a form of *nuisance aquatic life* due to their ability to produce toxins that can adversely affect aquatic life and the uses of the lake for watering by livestock and wildlife. In severe cases, levels of these toxins in lake water can affect human health.

IDNR is aware that some of the *aesthetically objectionable conditions* and/or *undesirable or nuisance aquatic life* at the lakes assessed as "impaired" may not be attributable to either wastewater discharges or agricultural practices. For example, a number of lakes assessed as "impaired" based on TSI values are very shallow (mean depth less than 2 meters) natural lakes of glacial origin with very low watershed-to-surface area ratios. The turbidity-related water quality problems at these lakes, whether caused by algae or suspended inorganic sediments, are due primarily to lack of sufficient water depth to prevent internal nutrient recycling and sediment re-suspension due to either bottom-feeding fish (e.g., common carp) and/or wind/wave action. Regardless, the levels of turbidity (whether of algal or non-algal origin) at these lakes constitute limitations to the use of these lakes for their designated beneficial uses. Thus, these lakes are appropriate for addition to the state list of impaired waters.

Data sources:

The primary data source for assessing the degree to which Iowa lakes support their designated primary contact uses is chlorophyll-a and Secchi depth values generated for 131 Iowa lakes sampled as part of the ISU and SHL surveys from 2004 through 2008. Data for inorganic suspended solids and total phosphorus from these surveys were also used to interpret TSI values and to provide a more complete assessment of lake water quality. Information from the IDNR Fisheries Bureau on recent water quality conditions/problems, the status of fish populations, and on lake history was used where appropriate to supplement assessments based on TSI values for chlorophyll-a and/or Secchi depth and to verify the existence of any "aesthetically objectionable condition" suggested by TSI values. In addition, information on lake phytoplankton communities from the ISU and SHL surveys was used to determine the amount and proportion of cyanobacteria in the water column. The amount of cyanobacteria was used to determine potential impairments due to nuisance aquatic life.

Data requirements for listing:

Data quantity:

In 1990, in order to improve the accuracy and confidence level of water quality assessments, IDNR developed "data completeness guidelines" for using results of routine water quality monitoring for Section 305(b) reporting. These state guidelines identify the numbers of samples needed for water quality assessments that can support Section 303(d) listings (i.e., a *monitored* assessment). Assessments based on less than the recommended number of samples are considered "evaluated"; these assessments are of relatively lower confidence than "monitored" assessments and are thus not appropriate for Section 303(d) impaired waters listing but are appropriate for Section 305(b) water quality reporting. In order to account for the year-to-year variability in lake water quality, state limnologists participating in the U.S. EPA Region 7 nutrient criteria regional technical assistance group (RTAG) (IA, KS, MO, NE) recommend in 2001 that the combined data from at least three years of monitoring conducted from three to five times per year should be used to characterize lake water quality and to identify water quality impairments. This recommendation has been incorporated into IDNR's data completeness guidelines. Thus, for purposes of Iowa's 2010 Integrated Report, overall median water quality values from the five-year period from 2004 through 2008 (approximately 25 samples) will be used to calculate TSI values to determine the existence of an impairment. As is typical in all monitoring networks, special circumstances occasionally prevent either sample collection (e.g., adverse weather conditions) or the reporting of data (e.g., laboratory accidents). For purposes of identifying candidate lakes for Iowa's 2010 impaired waters list, only those lakes with at least 10 samples each for chlorophyll-a and Secchi depth over the 2004-2008 period will be considered to meet IDNR's data completeness guidelines. Assessments for lakes with fewer than 10 samples for this period will be considered "evaluated" and thus will not be used to identify candidate lakes for impaired waters listing. Other lake water quality datasets

appropriate for calculating TSI values will be reviewed to determine compliance with Iowa DNR's data completeness guidelines.

Data quality:

As specified in the 2001 Iowa Code, Section 455B.194, subsection 1, (Iowa's credible data law) the department shall use credible data when determining whether any water of the state is to be placed on or removed from any Section 303(d) list (Category 5 of the Integrated Report). In addition, Iowa's credible data law specifies that data more than five years before the end of the most current Section 305(b) period (the end of calendar year 2008) are presumed under state law to be "not credible" unless IDNR identifies compelling reasons as to why the older data are credible. Data generated by the ISU lake survey and through the SHL lake monitoring network meet all requirements of Iowa's credible data law and can thus be used to add waters to Iowa's 2010 impaired waters list. Other datasets appropriate for calculating TSI values will be reviewed to determine compliance with Iowa's credible data law.

Threshold TSI values:

Similar to Iowa's 2004, 2006, and 2008 reporting/listing cycles, a TSI value of greater than or equal to 65 for either chlorophyll-a or Secchi depth will be used to identify candidate lakes for Category 5 of Iowa's 2010 Integrated Report (see Table 1 for a description of the "Integrated Report" categories). This threshold is similar to that used by the Minnesota Pollution Control Agency for lakes in the Western Corn Belt Plains ecoregion of southern Minnesota (MPCA 2005). Nearly the entire state of Iowa lies in this same ecoregion, the exceptions being (1) the portion of south-central and southeastern Iowa in the Central Irregular Plains ecoregion and (2) the portion of northeastern Iowa in the Driftless Area ecoregion. Lakes with TSI values greater than or equal to 65 are likely to have nutrient or sediment-related water quality problems that contribute to excessive turbidity (algal or non-algal) that impair the Class A1 uses and are thus potential candidates for Section 303(d) listing.

Assessment categories ("monitored" and "evaluated"):

Prior to recent revisions to guidance for state compliance with Sections 305(b) and 303(d) of the Clean Water Act (U.S. EPA 2003, 2005), U.S. EPA (1997) recommended that states identify water quality assessments as one of two types: evaluated or monitored. "Evaluated" assessments are those based on data older than five years or other than site-specific ambient monitoring data (e.g., questionnaire surveys of fish and game biologists [=best professional judgment] or predictive modeling using estimated input values) and thus are of relatively low confidence. In contrast, "monitored" assessments are based primarily on recent, site-specific ambient monitoring data and thus are of relatively high confidence. IDNR has historically not considered waterbodies identified as impaired based on evaluated (lower confidence) assessments as candidates for the state's Section 303(d) list. IDNR has, however, historically considered waterbodies identified as impaired based on monitored (higher confidence) assessments as candidates for the state's Section 303(d) list. In order to maintain continuity with past

assessment procedures, and due to the usefulness of EPA's (1997) recommendation, IDNR continues to (1) identify each assessment of lake water quality as either evaluated or monitored and (2) consider only lakes with recent site-specific data ("monitored" assessments) as candidates for Section 303(d) listing. Similar to listings for other types of waterbodies, however, once a lake is added to the state's Section 303(d) list, the lake will remain on the list until new data or some other good cause suggests that the lake should be removed from Iowa's list. Age of data is not an acceptable reason for removing waters from the state's Section 303(d) list.

Use support categories:

The following are detailed descriptions of the use support categories used for Section 305(b) lake assessments for the 2010 reporting cycle. This approach is the same as that used for previous assessment/listing cycles in Iowa. The TSI values associated with each of these use support categories are summarized in Table 3-4. Any impairments (i.e., "aesthetically objectionable conditions") suggested by TSI values for chlorophyll-a and/or Secchi depth are verified by IDNR field (Fisheries) staff.

Not Supporting and "monitored": candidate for Section 303(d) listing:

If the overall (2004-2008) lake-specific median summer TSI value for either chlorophyll-a or Secchi depth is greater than or equal to 70, then the lake should be assessed as "not supporting" designated uses, and the lake should be considered as a candidate for Section 303(d) listing. These lakes are likely to have severe turbidity-related impacts, of either algal or non-algal origin that (1) interfere with designated uses for primary contact recreation and (2) constitute an aesthetically objectionable condition that violates narrative criteria for general use waters as defined in Section 61.3(2) of the *Iowa Water Quality Standards*. The TSI threshold value for chlorophyll-a and/or Secchi depth is the lower limit that identifies "hypereutrophic" lakes (Table 3-1). Thus, this threshold value provides strong evidence of a water quality impairment.

Partially Supporting and "monitored": candidates for Section 303(d) listing:

If the overall (2004-2008) lake-specific median summer TSI value for either chlorophyll-a or Secchi depth is 65 to 69, then the lake should be assessed as "partially supporting" designated uses, and the lake should be considered as a candidate for Section 303(d) listing. These lakes are likely to have moderate turbidity-related impacts of either algal or non-algal origin that interfere with designated uses for primary contact recreation. TSI values from 65 to 69 are in the middle to upper range between eutrophic and hypereutrophic lakes (Table 3-1). The chlorophyll-a and Secchi depth threshold values for this use support category (65 to 69) are those used by the Minnesota Pollution Control Agency to identify Section 303(d)-impaired lakes in southern Minnesota (MPCA 2005). As such, this threshold is appropriate for identifying impairments in Iowa lakes.

Partially Supporting and “evaluated”: not candidates for Section 303(d) listing:

If the overall (2004-2008) lake-specific median summer TSI value for either chlorophyll-a or Secchi depth is 65 to 69, but the TSI value(s) is based on less than sufficient data, then the lake should be assessed as “partially supporting” designated uses but should not be considered a candidate for Section 303(d) listing. These lakes may have turbidity-related impacts, of either algal or non-algal origin, that may interfere with designated uses for primary contact recreation and/or aquatic life. Thus, while the TSI values for Iowa lakes in this category *may* be impaired for Class A1 uses, insufficient data are available for developing Section 305(b) assessments having the high degree of confidence needed to justify Section 303(d) listing. These lakes will be placed into Integrated Report categories 2b or 3b and will thus be added to Iowa's list of waters in need of further investigation. Note: due to the existence of sufficient data for chlorophyll-a and Secchi depth from lakes in Iowa's ambient lake monitoring program, TSI-based “evaluated” (lower confidence) assessments are extremely rare.

Fully Supporting / Threatened and “monitored”: candidates for Section 303(d) listing:

EPA (2005) recommends that states consider as “threatened” those waters that are currently attaining water quality standards but which are expected to not meet water quality standards by the next listing cycle (i.e., with the next two years). For example, a water should be listed if an analysis demonstrates a declining trend in a specific water quality criterion, and the projected trend will result in a failure to meet a criterion by the date of the next list (i.e., 2012 for purposes of the 2010 assessment cycle); or, segments should be listed if there are proposed activities that will result in violations of water quality standards.

Lakes with overall (2004-2008) summer median TSI values for chlorophyll-a and Secchi depth of less than 65, but that demonstrate adverse trends in either of these parameters such that impairment is likely for the next (2012) reporting/listing cycle, will be considered “fully supported/threatened (impaired)” and considered candidates for addition to IR Category 5 (Section 303(d) list).

Identifying water quality trends in “threatened” lakes: For the majority of Iowa lakes, sufficient data do not exist to determine the existence of water quality trends prior to 2000. This lack of historical data stems from the design of previous statewide surveys of Iowa lakes which involved sampling during only one summer season at approximately 10-year intervals (e.g., see Bachmann et al. 1980, Bachmann et al. 1994). The year-to-year variability in lake data—due largely to climatic factors—makes the existing historical (i.e., pre-2000) data of little use for trend determination. Due, however, to the continuity of the current lake

monitoring program, sufficient data exist since 2000 to begin to identify trends in lake water quality over time. Although this nine-year period provides barely enough data to determine trends, the lake-specific data will be examined to determine the existence of any potential changes in water quality over time.

Fully Supporting (not threatened): "evaluated" or "monitored": not candidates for Section 303(d) listing:

Lakes with overall (2004-2008) summer median TSI values for chlorophyll-a and Secchi depth less than 65 are assessed as "fully supporting" their designated uses for primary contact recreation. These lakes have moderately-good (TSI approaching 65) to sometimes exceptional (TSI < 50) water quality with only brief episodes of marginal water quality conditions. The TSI threshold values for both chlorophyll-a and Secchi depth in this category range from the middle range between eutrophic and hyper-eutrophic lakes to the upper range of mesotrophic lakes. Thus, the range of lake quality in this assessment category is considerable.

The narrative descriptions of these assessments in this database use qualitative characterizations of TSI values (e.g., "good", "poor", "high"; "low") ; Table 3-5 summarizes these characterizations.

MANAGEMENT AND ACCESSIBILITY OF ASSESSMENTS:

The Section 305(b) assessments of the degree of support of the primary contact recreation (Class A1) and aquatic life (Class B(LW) or B(WW)) uses for the 131 lakes sampled as part of the DNR's lake monitoring programs are entered into Iowa DNR's Section 305(b) assessment database (ADBNet; <http://programs.iowadnr.gov/adbnet/index.aspx>).

Table 3-1. Changes in temperate lake attributes according to trophic state (modified from U.S. EPA 2000, Carlson and Simpson 1996, and Oglesby et al. 1987).

TSI Value	Attributes	Primary Contact Recreation	Aquatic Life (Fisheries)
50-60	eutrophy: anoxic hypolimnia; macrophyte problems possible	[none]	warmwater fisheries only; percid fishery; bass may be dominant
60-70	bluegreen algae dominate; algal scums and macrophyte problems occur	weeds, algal scums, and low transparency discourage swimming and boating	Centrarchid fishery
70-80	hyper-eutrophy (light limited). Dense algae and macrophytes	weeds, algal scums, and low transparency discourage swimming and boating	Cyprinid fishery (e.g., common carp and other rough fish)
>80	algal scums; few macrophytes	algal scums, and low transparency discourage swimming and boating	rough fish dominate; summer fish kills possible

Table 3-2. Iowa lakes with overall median TSI values for total phosphorus greater than 70 (=hyper-eutrophic) that have TSI values for chlorophyll-a and Secchi depth that do not suggest impairment of primary contact recreation (i.e., TSI values of less than 65). TSI values are based on data from the Iowa State University and University Hygienic Laboratory surveys of 131 Iowa lakes from 2000 through 2008 (N approximately equal to 38); lakes are ranked by the TSI value for total phosphorus.

Lake Name	County	TSI for total phosphorus	TSI for chlorophyll-a	TSI for Secchi depth
Red Rock Reservoir	Marion	78	51	64
Saylorville Reservoir	Polk	74	58	62
Hannen Lake	Benton	72	63	59
Dog Creek Lake	O'Brien	72	63	62
Lake Keomah	Mahaska	71	63	57
West Lake (Osceola)	Clarke	71	60	62

Table 3-3. Ranges of TSI values for Iowa lakes based on median values from the five years of the statewide survey used for the 2010 assessment/listing cycle (2004-2008) of 131 Iowa lakes by Iowa State University and University of Iowa Hygienic Laboratory. Lakes were sampled approximately three times per summer.

	TSI Values:		
	total phosphorus	chlorophyll-a	Secchi depth
minimum	40	42	38
10 th percentile	57	53	50
25 th percentile	62	59	58
median	67	65	63
75 th percentile	72	69	70
90 th percentile	77	72	74
maximum	88	79	83
mean of medians	67	64	63
standard deviation	9	8	9

Table 3-4. Summary of ranges of TSI values and measurements for chlorophyll-a and Secchi depth used to define Section 305(b) use support categories for the 2004, 2006, 2008, and 2010 reporting cycles.

Level of Support	TSI value	Chlorophyll-a (ug/l)	Secchi Depth (m)
fully supported	≤55	≤12	≥1.4
fully supported / threatened (candidate for Section 303(d) listing)	55 → 65	12 → 33	1.4 → 0.7
partially supported (evaluated: in need of further investigation)	65 → 70	33 → 55	0.7 → 0.5
partially supported (monitored: candidates for Section 303(d) listing)	65 → 70	33 → 55	0.7 → 0.5
not supported (monitored or evaluated: candidates for Section 303(d) listing)	≥70	≥55	≤0.5

Table 3-5. Narrative descriptions of TSI ranges for Secchi depth, phosphorus, and chlorophyll-a for Iowa lakes used for the 2004, 2006, 2008, and 2010 Section 305(b) reporting cycles. These characterizations were used in developing lake-specific assessments that are included in the Iowa DNR's Section 305(b) assessment database (ADBNet).

TSI value	Secchi description	Secchi depth (m)	Phosphorus & Chlorophyll-a description	Phosphorus levels (ug/l)	Chlorophyll-a levels (ug/l)
> 75	extremely poor	< 0.35	extremely high	> 136	> 92
70-75	very poor	0.5 – 0.35	very high	96 - 136	55 – 92
65-70	poor	0.71 – 0.5	high	68 – 96	33 – 55
60-65	moderately poor	1.0 – 0.71	moderately high	48 – 68	20 – 33
55-60	relatively good	1.41 – 1.0	relatively low	34 – 48	12 – 20
50-55	very good	2.0 – 1.41	low	24 – 34	7 – 12
< 50	exceptional	> 2.0	extremely low	< 24	< 7

References for Attachment 3:

- Bachmann, R.W. 1965. Some chemical characteristics of Iowa lakes and reservoirs. *Proceedings of the Iowa Academy of Science*, 72:238-243.
- Bachmann, R.W., T.A. Hoyman, L.K. Hatch, and B.P. Hutchins. 1994. A classification of Iowa's lakes for restoration. Department of Animal Ecology, Iowa State University, Ames, Iowa. 517 p.
- Bachmann, R.W., M.R. Johnson, M.V. Moore, and T.A. Noonan. 1980. Clean lakes classification study of Iowa's lakes for restoration. Iowa Cooperative Fisheries Research Unit and Department of Animal Ecology, Iowa State University, Ames, Iowa. 715 p.
- Bachmann, R.W., R. Lohnes, and D. Bonneau. 1982. Swan Lake restoration (Phase I, final report): a diagnostic and feasibility study for pollution abatement and restoration of Swan Lake, Carroll County, Iowa. Iowa Department of Environmental Quality. 134 p.
- Carlson, R.E. 1977. A trophic state index for lakes. *Limnol Oceanogr.* 22:361-369.
- Carlson, R.E. 1991. Expanding the trophic state concept to identify non-nutrient limited lakes and reservoirs. In: *Proceedings of a National Conference on Enhancing the States' Lake Management Programs, Monitoring and Lake Impact Assessment*, Chicago. pp. 59-71.
- Carlson, R.E. 1984. Discussion: using differences among Carlson's trophic state index values in regional water quality assessment, by Richard A. Osgood. *Water Res. Bull.* 19(2):307-309.
- Carlson, R.E. and J. Simpson. 1996. A coordinator's guide to volunteer lake monitoring methods. North American Lake Management Society, Madison, WI.
- IAC. 2008. Chapter 567-61: water quality standards. Iowa Administrative Code [effective date 06/11/2008].
- MPCA. 2005. Guidance manual for assessing the quality of Minnesota surface waters for determinations of impairment: 305(b) report and 303(d) list. Environmental Outcomes Division, Minnesota Pollution Control Agency. 106 pp. plus appendices.
- Oglesby, R.T., J.H. Leach, and J. Forney. 1987. Potential Stizostedion yield as a function of chlorophyll concentration with special reference to Lake Erie. *Canadian Journal of Fisheries and Aquatic Sciences*, 44(Suppl.):166-170.
- U.S. EPA. 1997. Guidelines for the preparation of the comprehensive state water quality assessments (305(b) reports) and electronic updates. Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency, Washington, D.C.

U.S. EPA. 2000. Nutrient criteria technical guidance manual: lakes and reservoirs. Report No. EPA-822-B00-001, Office of Water, U.S. Environmental Protection Agency, Washington D.C.

U.S. EPA. 2003. Guidance for 2004 assessment, listing, and reporting requirements pursuant to Sections 303(d) and 305(b) of the Clean Water Act, July 21, 2003. Watershed Branch, Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency. 32 p.

U.S. EPA. 2005. Guidance for 2006 assessment, listing, and reporting requirements pursuant to Sections 303(d), 305(b), and 314 of the Clean Water Act, July 29, 2005. Watershed Branch, Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency. 70 p. plus appendices.

Attachment 4

METHODOLOGY FOR ASSESSING THE DEGREE TO WHICH IOWA'S SHALLOW NATURAL LAKES SUPPORT THEIR DESIGNATED AQUATIC LIFE USES FOR THE 2010 INTEGRATED REPORTING CYCLE

Iowa DNR
Watershed Monitoring & Assessment Section
Geological and Water Survey

INTRODUCTION:

Due to the lack of appropriate water quality criteria, and due to the lack of an assessment protocol, IDNR has historically relied on the professional judgment of IDNR biologists to assess Iowa's shallow lakes and wetlands. Although assessed for purposes of Section 305(b) reporting, Iowa's wetlands and shallow lakes have typically not been identified as candidates for Section 303(d) impaired waters listing. That is, without water quality monitoring data, and without an assessment protocol to objectively identify the degree to which a shallow lake or wetland supported its designated aquatic life use, IDNR was unable to develop high-confidence assessments that would support a Section 303(d) listing.

In 2006, the IDNR Watershed Monitoring and Assessment Section initiated routine water quality monitoring on several shallow lakes and wetlands in north-central and northwest Iowa. This monitoring has continued through 2008. Thus, for the 2010 assessment/listing cycle, data generated from 2006-08 for Secchi depth and chlorophyll-a from 11 of Iowa's shallow natural lakes of glacial origin (Table 4-1) will be used with Carlson's (1977) trophic state index to identify the degree to which these shallow lakes support their designated Class B(LW) aquatic life uses. Information from IDNR field staff on the status of aquatic macrophytes and aquatic macroinvertebrates at the shallow lakes monitored will be used to supplement the water quality assessments developed.

ASSESSMENT RATIONALE:

The concept of "trophic state" has long been used by limnologists to classify lakes and is based on the chemistry and biology of lakes. Although a number of approaches exist for classifying lakes according to trophic state, and although a number of controversies exist regarding how "trophic state" is defined, the use of this framework has the advantages of historical usage, general acceptance of the trophic state concept (e.g., "eutrophic" indicates nutrient enrichment), and an improved ability to describe lake condition versus a description using a single variable or number (e.g., total phosphorus concentration). Table 4-2 describes the general framework of the lake trophic state concept. For a discussion on the development and variety of trophic state indices, see Chapter 2 (*The Basis for Lake and Reservoir Nutrient Criteria*) in U.S. EPA (2000) (see <http://www.epa.gov/waterscience/criteria/nutrient/guidance/lakes/lakes.pdf>).

Carlson's (1977) trophic state index is a numeric indicator of the continuum of the biomass of suspended algae in lakes and thus reflects a lake's nutrient condition and water transparency. The level of plant biomass is estimated by calculating the TSI value for chlorophyll-a. TSI values for Secchi depth serves as surrogate measures of the TSI value for chlorophyll. Carlson's trophic state index provides a convenient and well-established method for identifying turbidity-related impacts to Iowa lakes and thus seems appropriate for assessing the degree to which Iowa's shallow lakes support their designated Class B(LW) aquatic life uses.

Because of the direct linkage between turbidity and attainment of aquatic life goals, TSI values for chlorophyll-a and Secchi depth will be used to identify shallow lakes in Iowa that do not fully support their designated Class B(LW) aquatic life uses. IDNR field staff will provide information from surveys for aquatic macrophytes, aquatic macroinvertebrates, and fish populations to supplement the assessment and to corroborate any impairment of aquatic life uses that is identified. IDNR field staff will be contacted to ensure that the TSI-based assessment is consistent with their knowledge of the particular shallow lake.

The connection of water clarity and chlorophyll-a (as interpreted by the trophic state index) at shallow lakes to the *Iowa Water Quality Standards* (IAC 2008) is the attainment of the designated Class B(LW) aquatic life use. This use is defined as follows:

Lakes and wetlands (Class "B(LW)"). These are artificial and natural impoundments with hydraulic retention times and other physical and chemical characteristics suitable to maintain a balanced community normally associated with lake-like conditions (IAC 2008).

The goal of Iowa's shallow lakes management strategy is to use lake management techniques such as lake draw-downs and biomanipulation to shift the lake from a turbid, algae-dominated system with little or no rooted aquatic vegetation and with poor sport fisheries to a clear-water, macrophyte-dominated state that supports a balanced warmwater aquatic community of fish, aquatic macroinvertebrates, and aquatic vegetation (macrophytes) (IDNR 2008). This TSI-based assessment method, with its focus on water clarity, provides an objective measure of the relative success of IDNR's management strategy.

This methodology applies only to shallow lakes and not to wetlands. For purposes of the 2010 assessment/listing cycle, shallow lakes are defined as lakes with maximum depths typically greater than seven feet but less than 15 feet. Shallow lakes typically do not stratify thermally in summer. Abundant rooted aquatic vegetation (macrophytes), including submergent and emergent vegetation, may cover much of a shallow lake. Shallow lakes can support a variety of beneficial uses including boating, fishing, waterfowl production, hunting, aesthetics, and limited swimming. Wetlands have maximum depths typically less than seven feet, often have minimal open water in summer, and are typically not managed as sport fisheries but for waterfowl and wildlife production, hunting, and aesthetics. Wetlands are not managed for swimming uses and lack beach areas.

IDENTIFYING WATER QUALITY IMPAIRMENTS AT SHALLOW LAKES

Overview:

For purposes of developing water quality assessments for the 2010 Section 305(b) reporting cycle, Carlson's (1977) "trophic state index" (TSI) was used with the three years of data generated for 11 Iowa shallow lakes as part of Iowa DNR surveys from 2006 through 2008. Overall (three-year) summer-season median values calculated from this dataset were used to calculate TSI values for chlorophyll-a, and Secchi depth for each lake. The identification of impairments of aquatic life uses was based on the resulting TSI values for chlorophyll-a and/or Secchi depth.

Relevant state water quality criteria:

The *Iowa Water Quality Standards* (IAC 2008) do not contain numeric criteria for nutrients (e.g., nitrogen or phosphorus), chlorophyll, or turbidity that apply to Class B(LW) aquatic life uses. Thus, the assessments of the degree to which these parameters might impair the Class B(LW) uses are based on a determination of whether this use is impaired by turbidity as interpreted through the trophic state index (Carlson 1977).

Data Sources:

Data for chlorophyll-a and Secchi depth collected by IDNR staff from 2006 through 2008 will be used with the trophic state index. IDNR field staff will also provide information on the status of aquatic macrophyte, macroinvertebrate, and fish communities at the shallow lakes assessed.

Data requirements for listing:

Data quantity:

In 1990, in order to improve the accuracy and confidence level of water quality assessments, IDNR developed "data completeness guidelines" for using results of routine water quality monitoring for Section 305(b) reporting. These state guidelines identify the numbers of samples needed for water quality assessments that can support Section 303(d) listings (i.e., a *monitored* assessment). Assessments based on less than the recommended number of samples are considered "evaluated"; these assessments are of relatively lower confidence than "monitored" assessments and are thus not appropriate for impaired waters listing but are appropriate for Section 305(b) water quality reporting.

In order to account for the year-to-year variability in lake water quality, state limnologists participating in the U.S. EPA Region 7 nutrient criteria regional technical assistance group (RTAG) (IA, KS, MO, NE) recommend in 2001 that the combined data from at least three years of monitoring conducted from three to five times per year should be used to characterize lake water quality and to identify water quality impairments.

This recommendation has been incorporated into IDNR's data completeness guidelines.

Thus, for purposes of Iowa's 2010 Integrated Report, overall summer-season median water quality values from the three-year period from 2006 through 2008 will be used to calculate TSI values to determine the existence of a turbidity-related impairment. Only those shallow lakes with at least nine samples for chlorophyll-a and Secchi depth over the 2006-2008 period will be considered to meet IDNR's data completeness guidelines. Assessments for shallow lakes with fewer than nine samples for this period will be considered "evaluated" and thus will not be used to identify candidate lakes for Section 303(d) impaired waters listing.

Data quality:

As specified in the 2001 Iowa Code, Section 455B.194, subsection 1, (Iowa's credible data law) the department shall use credible data when determining whether any water of the state is to be placed on or removed from any Section 303(d) list (Category 5 of the Integrated Report). In addition, Iowa's credible data law specifies that data more than five years before the end of the most current Section 305(b) period (the end of calendar year 2008) are presumed under state law to be "not credible" unless IDNR identifies compelling reasons as to why the older data are credible. Data generated by the IDNR staff as part of the 2006-08 shallow lakes surveys meet all requirements of Iowa's credible data law and can thus be used to shallow lakes to Iowa's 2010 impaired waters list.

Threshold TSI values:

Similar to the approach for assessing lake water quality that Iowa has used since the 2004 reporting/listing cycle, a TSI value of equal to or greater than 65 for either chlorophyll-a or Secchi depth will be used to identify candidate shallow lakes for Section 303(d) listing and addition to Category 5 of Iowa's 2010 Integrated Report (see [Table 2](#) for a description of the "Integrated Report" categories). Lakes with TSI values greater than or equal to 65 are likely to have nutrient or sediment-related water quality problems that contribute to excessive turbidity (algal or non-algal) that impair the Class B(LW) aquatic life uses and are thus potential candidates for Section 303(d) listing. Shallow lakes with TSI values approaching, but not exceeding, 65 will also be considered candidates for Section 303(d) listing if data suggest a worsening water quality trend that threatens full support. This methodology is similar to that used by the Minnesota Pollution Control Agency for lakes in the Western Corn Belt Plains ecoregion of southern Minnesota (MPCA 2005). All of Iowa's natural lakes of glacial origin lie within this ecoregion.

Assessment categories ("monitored" and "evaluated"):

Prior to recent revisions to guidance for state compliance with Sections 305(b) and 303(d) of the Clean Water Act (U.S. EPA 2003, 2005), U.S. EPA (1997) recommended that states identify water quality assessments as one of two types:

evaluated or monitored. Evaluated assessments are those based on data older than five years or other than site-specific ambient monitoring data (e.g., questionnaire surveys of fish and game biologists [=best professional judgment] or predictive modeling using estimated input values) and thus are of relatively low confidence. In contrast, monitored assessments are based primarily on recent, site-specific ambient monitoring data and thus are of relatively high confidence. IDNR has historically not considered waterbodies identified as impaired based on evaluated (lower confidence) assessments as candidates for the state's Section 303(d) list. IDNR has, however, historically considered waterbodies identified as impaired based on monitored (higher confidence) assessments as candidates for the state's Section 303(d) list. In order to maintain continuity with past assessment procedures, and due to the usefulness of EPA's (1997) recommendation, IDNR continues to (1) identify each assessment of lake water quality as either evaluated or monitored and (2) only consider lakes with recent site-specific data ("monitored" assessments) as candidates for Section 303(d) listing.

Use support categories:

The following are detailed descriptions of the use support categories used for Section 305(b) shallow lake assessments for the 2010 reporting cycle. The TSI values associated with each of these use support categories are summarized in Table 4-3. Any impairments suggested by TSI values for chlorophyll-a and/or Secchi depth are verified by IDNR field staff.

Not Supporting and "monitored": candidate for Section 303(d) listing:

If the overall (2006-2008) lake-specific summer-season median TSI value for either chlorophyll-a or Secchi depth based on at least nine samples is greater than or equal to 70, then the lake should be assessed as "not supporting" its designated aquatic life uses, and the lake should be considered as a candidate for Section 303(d) listing. These lakes are likely to have severe turbidity-related impacts, of either algal or non-algal origin that prevent the shallow lake from supporting its Class B(LW) aquatic life use. The TSI threshold value of 70 for chlorophyll-a and/or Secchi depth is the lower limit that identifies "hypereutrophic" lakes (Table 4-2). Thus, this threshold value provides strong evidence of a water quality impairment.

Partially Supporting and "monitored": candidate for Section 303(d) listing:

If the overall (2006-2008) lake-specific median summer TSI value for either chlorophyll-a or Secchi depth based on at least nine samples is 65 to 69, then the lake should be assessed as "partially supporting" designated uses, and the lake should be considered as a candidate for Section 303(d) listing. These lakes are likely to have moderate turbidity-related impacts of either algal or non-algal origin that interfere with support of aquatic life uses. TSI values from 65 to 69 are in the middle to upper range between eutrophic and hypereutrophic lakes. The

chlorophyll-a and Secchi depth threshold values for this use support category (65 to 69) are those used by the Minnesota Pollution Control Agency to identify Section 303(d)-impaired lakes in southern Minnesota (MPCA 2005). As such, this threshold is appropriate for identifying impairments in Iowa lakes.

Partially Supporting and “evaluated”: not candidates for Section 303(d) listing:

If the overall (2006-2008) lake-specific median summer TSI value for either chlorophyll-a or Secchi depth is 65 to 69, but the TSI value(s) is based on less than sufficient data (i.e., less than nine samples), then the lake should be assessed as “partially supporting” designated uses but should not be considered a candidate for Section 303(d) listing. These lakes may have turbidity-related impacts, of either algal or non-algal origin, that may interfere with support of designated uses for aquatic life. Thus, while the TSI values for Iowa lakes in this category *may* be impaired for Class B(LW) uses, insufficient data are available for developing Section 305(b) assessments having the high degree of confidence needed to justify Section 303(d) listing. These shallow lakes will be placed into Integrated Report categories 2b or 3b and will thus be added to Iowa's list of waters in need of further investigation.

Fully Supporting / Threatened and “monitored”: candidates for Section 303(d) listing:

EPA (2005) recommends that states consider as “threatened” those waters that are currently attaining water quality standards but which are expected to not meet water quality standards by the next listing cycle (every two years). For example, a water should be listed if an analysis demonstrates a declining trend in a specific water quality criterion, and the projected trend will result in a failure to meet a criterion by the date of the next list (i.e., 2012 for purposes of the 2010 assessment cycle); or, segments should be listed if there are proposed activities that will result in violations of water quality standards.

Shallow lakes with overall (2006-2008) summer-season median TSI values for chlorophyll-a and Secchi depth based on at least nine sample of less than 65, but that demonstrate adverse trends in either of these parameters such that impairment is likely for the next (2012) reporting/listing cycle, will be considered “fully supported/threatened (impaired)” and considered candidates for addition to IR Category 5 (Section 303(d) list). Because, however, sufficient data do not currently exist to determine the existence of water quality trends at Iowa's shallow lakes, identification of adverse trends will likely not be possible for the 2010 assessment/listing cycle.

Fully Supporting (not threatened); “monitored”: not candidates for Section 303(d) listing:

If the overall (2006-2008) lake-specific summer-season median TSI values for both chlorophyll-a or Secchi depth are less than 65 in the absence of any adverse water quality trend, and the overall median is based on based on at least nine samples, then the lake should be assessed as “fully supporting” its designated aquatic life uses. The assessment type should be considered “monitored” (i.e., of higher confidence), and the water should be placed into Category 1a of the Integrated Report. The TSI threshold values for both chlorophyll-a and Secchi depth in this category range from the middle range between eutrophic and hyper-eutrophic lakes to the upper range of mesotrophic lakes.

Fully Supporting (not threatened); “evaluated”: not candidates for Section 303(d) listing:

If the overall (2006-2008) lake-specific summer-season median TSI values for both chlorophyll-a or Secchi depth are less than 65 in the absence of any adverse water quality trend, and the overall median is based on based on fewer than nine samples, then the lake should be assessed as “fully supporting” its designated aquatic life uses. The assessment type, however, should be indicated as “evaluated” (i.e., of lower confidence).

MANAGEMENT AND ACCESSIBILITY OF ASSESSMENTS:

The Section 305(b) assessments of the degree of support of the Class B(LW) uses for the shallow lakes sampled as part of the IDNR survey are entered into Iowa DNR's Section 305(b) assessment database (ADBNet; <http://programs.iowadnr.gov/adbnet/index.aspx>).

References for Attachment 4:

- Carlson, R.E. 1977. A trophic state index for lakes. *Limnol Oceanogr.* 22:361-369.
- Carlson, R.E. and J. Simpson. 1996. *A Coordinator's Guide to Volunteer Lake Monitoring Methods*. North American Lake Management Society. 96 pp.
- Heiskary, Steven A. and C. Bruce Wilson. 2005. *Minnesota Lake Water Quality Assessment Report: Developing Nutrient Criteria*. 3rd edition.
- IAC. 2008. Chapter 567-61: water quality standards. Iowa Administrative Code [effective date 06/11/2008].
- IDNR. 2008. Shallow lake management initiative. Pgs. 27-30, In: *Lake restoration: 2008 report and 2009 plan*. Iowa Department of Natural Resources, Des Moines. 48 p. Available from <http://www.iowadnr.gov/water/lakerestoration/files/08report.pdf>.
- MPCA. 2005. *Guidance manual for assessing the quality of Minnesota surface waters for determinations of impairment: 305(b) report and 303(d) list*. Environmental Outcomes Division, Minnesota Pollution Control Agency. 106 pp. plus appendices.
- Oblesby, R.T., J.H. Leach, and J. Forney. 1987. Potential *Stizostedion* yield as a function of chlorophyll concentration with special reference to Lake Erie. *Can. J. Fish. Aquat. Sci.* 44(Suppl.)2:166-170.
- UMRCC. 2003. *Proposed water quality criteria necessary to sustain submersed aquatic vegetation in the Upper Mississippi River*. Water Quality Technical Section, Upper Mississippi River Conservation Committee. 6 p. Unpublished.
- U.S. EPA. 1997. *Guidelines for the preparation of the comprehensive state water quality assessments (305(b) reports) and electronic updates*. Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency, Washington, D.C.
- U.S. EPA. 2000. *Nutrient criteria technical guidance manual: lakes and reservoirs*. Report No. EPA-822-B00-001, Office of Water, U.S. Environmental Protection Agency, Washington D.C.
- U.S. EPA. 2003. *Guidance for 2004 assessment, listing, and reporting requirements pursuant to Sections 303(d) and 305(b) of the Clean Water Act, July 21, 2003*. Watershed Branch, Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency. 32 p.
- U.S. EPA. 2005. *Guidance for 2006 assessment, listing, and reporting requirements pursuant to Sections 303(d), 305(b), and 314 of the Clean Water Act, July 29, 2005*. Watershed Branch, Assessment and Watershed Protection Division, Office of Wetlands, Oceans, and Watersheds, Office of Water, U.S. Environmental Protection Agency. 70 p. plus appendices. (<http://www.epa.gov/owow/tmdl/2006IRG/>)

Table 4-1. Shallow natural (glacial) lakes monitored by Iowa DNR from 2006 through 2008.

Name	WB ID	Location	County	Designated Uses	Size (acres)	Year(s) Monitored
Barringer Slough	IA 06-LSR-02350-L_0	S14, T96N, R35W	Clay	B(LW), HH	778	2008
Big Wall Lake	IA 02-IOW-00860-L_0	S14, T90N, R24W	Wright	B(LW), HH	935	2008
Cheever Lake	IA 04-UDM-0505-L_0	S20, T99N, R34W	Emmet	[Not designated]	112	2007-08
Dan Green Slough	IA 06-LSR-02420-L_0	S20, T97N, R35W	Clay	B(LW), HH	311	2006-08
Diamond Lake	IA 06-LSR-3205-L_0	S15, T100N, R37W	Dickinson	B(LW), HH	166	2006
Elm Lake	IA 02-IOW-00870-L_0	S21, T92N, R24W	Wright	A1, B(WW2), HH, HQR	463	2008
Fourmile Lake	IA 04-UDM-0510-L_0	S19, T88N, R34W	Emmet	B(LW), HH	209	2006-07
Lizard Lake	IA-04-UDM-03110-L_0	S22, T91N, R34W	Pocahontas	B(LW), HH	268	2006-08
Pickereel Lake	IA 04-RAC-01690-L_0	S1, T93N, R35W	Buena Vista	A1, B(LW), HH	35	2006-08
South Twin Lake	IA 04-RAC-01395-L_0	S1, T88N, R33W	Calhoun	B(LW),HH	600	2006-08
Virgin Lake	IA 06-LSR-02330-L_0	S30, T96N, R34W	Palo Alto	B(LW), HH, HQR	200	2006-08

Table 4-2. Changes in temperate lake attributes according to trophic state (modified from U.S. EPA 2000, Carlson and Simpson 1996, and Oglesby et al. 1987).

TSI Value	Attributes	Primary Contact Recreation	Aquatic Life (Fisheries)
50-60	eutrophy: anoxic hypolimnia; macrophyte problems possible	[none]	warmwater fisheries only; percid fishery; bass may be dominant
60-70	bluegreen algae dominate; algal scums and macrophyte problems occur	weeds, algal scums, and low transparency discourage swimming and boating	Centrarchid fishery
70-80	hyper-eutrophy (light limited). Dense algae and macrophytes	weeds, algal scums, and low transparency discourage swimming and boating	Cyprinid fishery (e.g., common carp and other rough fish)
>80	algal scums; few macrophytes	algal scums, and low transparency discourage swimming and boating	rough fish dominate; summer fish kills possible

Table 4-3. Assessment and impairment thresholds for aquatic life uses of shallow lakes in Iowa based on trophic state index (TSI) values. TSI values are calculated using an overall three-year summer-season median value for chlorophyll-a and Secchi depth.

TSI value	Chlorophyll-a (median during growing season)	Secchi disk depth (median during growing season)	Rationale for threshold selection:	Assessment Decision:
60- < 65	20 to 33 ppb	1.0 m (3.3 ft) to 0.7 m (2.3 ft.)	Water quality is sufficient to support growth of aquatic macrophytes (UMRCC 2003).	<u>Full support:</u> TSI values less than 65 indicate full support of aquatic life uses and Clean Water Act goals.
60- < 65	20 to 33 ppb	1.0 m (3.3 ft) to 0.7 m (2.3 ft.)	Water quality degrading over time. As TSI values approach 65, the frequency of nuisance algal blooms increases and water clarity declines.	<u>Fully Supported/Threatened / Impaired:</u> Any adverse trends in apparent in data for chlorophyll-a or Secchi depth, however, suggest that full support is "threatened" such that impairment is likely by the time of the next 303(d) listing cycle.
65- ≤ 70	33 to 55 ppb	0.7 m (2.3 ft) to 0.5 ft (1.6 ft)	A TSI value of 65 is used by state of Minnesota as an impairment threshold for chlorophyll-a and Secchi depth in shallow lakes in the southern part of the state (Heiskary and Wilson 2005). TSI values 65 or greater indicate generally poor water transparency such that growth of aquatic macrophytes is suppressed or eliminated.	<u>Partially Supported / Impaired:</u> Water clarity is sufficiently poor that aquatic life uses can be considered moderately impaired.
≥ 70	55 ppb	0.5 m (1.6 ft.)	TSI values above 70 indicate heavy algal blooms in summer; light-limitation; hypereutrophic.	<u>Not Supported / Impaired:</u> Very poor water transparency suggest that aquatic life uses are severely impaired.

Attachment 5

Iowa DNR interpretations of Section 305(b)/303(d) causes of impairment.

Information is also included on the historical use of the individual cause categories for water quality assessments in Iowa and on the existence of numeric criteria in the *Iowa Water Quality Standards*. NA = "not applicable. Information is taken from several published and on-line sources (see "References, Attachment 5") as well as from IDNR staff experience from identifying these causes of impairment for Iowa waters.

Cause Category	Historically Used?	Numeric Criteria?	Description
ammonia (un-ionized)	yes	yes	Ammonia refers to the concentration of ionized (NH_4^+) and un-ionized ammonia (NH_3) in water. Ammonia is formed during bacterial decomposition of organic matter and is delivered to streams and rivers from wastewater discharges and from nonpoint sources. The primary source of ammonia dissolved in water comes from bacterial mineralization of dead plants and animals (Cole 1979). (Mineralization is the conversion of an element from an organic to an inorganic form as a result of microbial decomposition.) Impairments related to measured concentrations of ammonia in Iowa waters are rare. Most ammonia impairments are tied to fish kills caused by delivery of animal waste to streams; these impairments are based on the presumed presence of high levels of ammonia the high-strength animal waste generated by animal feeding operations to which fish kills are often attributed.
Arsenic	yes	yes	Arsenic is a naturally occurring element widely distributed in the earth's crust. In the environment, arsenic is combined with oxygen, chlorine, and sulfur to form inorganic arsenic compounds. Inorganic arsenic compounds are mainly used to preserve wood. Copper chromated arsenate (CCA) is used to make "pressure-treated" lumber. CCA is no longer used in the U.S. for residential uses; it is still used in industrial applications. Organic arsenic compounds are used as pesticides, primarily on cotton fields and orchards. Inorganic arsenic is known human carcinogen (source: ATSDR (http://www.atsdr.cdc.gov/toxfaqs/TF.asp?id=19&tid=3)). Arsenic impairments in Iowa waters are due to violations of Iowa's human health criterion designed to protect against adverse health impacts from consuming arsenic in water and fish. This criterion (0.18 parts per billion (ppb)) is well below what is believed to be the natural background concentration of arsenic in Iowa surface waters and groundwaters (~1 to 2 ppb) and is far below the U.S. EPA's maximum contaminant level of no more than 10 parts per billion in drinking water.
atrazine	yes	yes	A common pesticide (corn herbicide) that is in the triazine family of herbicides. The only criterion for atrazine in the <i>Iowa Water Quality Standards</i> is the maximum contaminant level of 3 ppb to protect drinking water (Class C) uses.
cause unknown	yes	NA	Causes of impairment are identified as "unknown" where results of water quality monitoring suggest an impact, but no cause of the impact is apparent. Most often, this cause category is used when results of biological monitoring identify an impact to biotic integrity but do not suggest a specific cause of the impact. In such cases, follow-up monitoring is often needed to determine the specific cause or causes of the impairment.
chloride	no	yes	Chloride (Cl^-) is a naturally-occurring negatively-charged dissolved constituent of water and is one of several similar ions that combine to constitute "total dissolved solids." Chloride is a major ion commonly found in streams and wastewater. Chloride may get into surface water from several

Cause Category	Historically Used?	Numeric Criteria?	Description
			<p>sources, including wastewater from certain industries, wastewater from communities that soften water, road salting, agricultural runoff, and produced water from oil and gas wells. Levels of chloride in Iowa surface waters are relatively low with a median concentration of 22 mg/l in the approximately 8,500 samples collected from 2000 through 2009 as part of Iowa DNR's ambient stream/river water quality monitoring network (IDNR 2010). Only 10% of these samples have had chloride levels greater than 39 mg/l; the maximum concentration in these samples was 170 mg/l. The only existing criterion for chloride in the Iowa Water Quality Standards is 250 mg/l to protect surface waters used as a source of a municipal water supply (i.e., Class C waters). Results of water quality monitoring to date have not shown levels of chloride in surface waters that suggest impairment of Class C uses. Lacking any approved water quality criteria to protect aquatic life from chloride, there are no historical or current aquatic life impairments for chloride in Iowa surface waters. IDNR has recently proposed water quality criteria to protect aquatic life from high levels of chloride; the criteria depend on both hardness and the sulfate concentrations (see http://www.iowadnr.gov/water/standards/files/ws_fact.pdf). At typical water hardness and sulfate concentrations, the chronic criterion to protect aquatic life from chloride would be approximately 380 mg/l. Chloride levels in Iowa waters are sufficiently low that implementation of IDNR's proposed criteria is unlikely to result in aquatic life impairments due to chloride.</p>
chlorine	yes	yes	<p>Chlorine and chloramines are widely used in treatment of potable water supplies and wastewater treatment plant effluents and are used in a variety of industrial applications, including power generating facilities and paper mills. Although the <i>Iowa Water Quality Standards</i> contain numeric criteria to protect aquatic life uses from adverse impacts of total residual chlorine, analytical difficulties have precluded analysis for total residual chlorine as part of ambient surface water monitoring since 1999. Currently, the only scenario that would lead to identification of chlorine as the cause of an impairment is the accidental release of chlorine to surface waters such that a fish kill occurs (e.g., as would potentially occur following a water main break).</p>
cyanide	no	yes	<p>Cyanide enters air, water, and soil from both natural processes and industrial activities. Cyanide is usually found joined with other chemicals to form compounds. Examples include hydrogen cyanide, sodium cyanide and potassium cyanide. Certain bacteria, fungi, and algae can produce cyanide. Cyanide and hydrogen cyanide are used in electroplating, metallurgy, organic chemicals production, photographic developing, manufacture of plastics, fumigation of ships, and some mining processes. Most cyanide in surface water will form hydrogen cyanide and evaporate. Cyanide in water does not build up in the bodies of fish (source: http://www.atsdr.cdc.gov/tfacts8.pdf). Detectable levels of cyanide are extremely rare in Iowa waters; there are no water quality impairments, historical or current, attributed to cyanide.</p>
dioxins	no	yes	<p>Dioxins and dioxin-like compounds are by-products of various industrial processes, and are commonly regarded as highly toxic compounds that are environmental pollutants and persistent organic pollutants. Dioxins are not intentionally produced and have no known use. They are the by-products of various industrial processes (i.e., bleaching paper pulp, and chemical and pesticide manufacture) and combustion activities (i.e., burning household trash, forest fires, and waste incineration). The defoliant Agent Orange, used during the Vietnam War, contained dioxins. Dioxins are found at low levels throughout the world in air, soil, water, sediment, and in foods such as meats, dairy, fish, and shellfish. The highest levels of dioxins are usually found in soil, sediment, and in the</p>

Cause Category	Historically Used?	Numeric Criteria?	Description
			fatty tissues of animals. Much lower levels are found in air and water. Sources: Wikipedia (http://en.wikipedia.org/wiki/Dioxins_and_dioxin-like_compounds) and ATSDR (http://www.atsdr.cdc.gov/substances/dioxin/policy/). In Iowa, doxins have been detected in samples of fish tissue but occur at extremely low levels (in the low parts per trillion range) and pose no known risk to human health or the aquatic environment.
excessive algal growth / chlorophyll-a	yes	no	Chlorophyll is the pigment in plants that is essential for photosynthesis whereby carbon dioxide and water are converted to carbohydrates and oxygen; chlorophyll-a is a form of chlorophyll that is common to all types of freshwater algae (e.g., green algae, cyanobacteria, and diatoms). For purposes of water quality assessment, chlorophyll-a is used as a surrogate measure of growth of algae in the water column. "Excessive algal growth" refers to an unusually large concentration of algal organisms (planktonic or benthic) that can adversely affect either the aesthetic quality of the surface water for water-based recreation or the ability of the waterbody to support the expected types and numbers of aquatic biota (see explanation for "turbidity" below). Scenarios that can lead to impairments due to "excessive algal growth" include the following: (1) large populations of common carp that increase water column nutrient levels through feeding and spawning activities such that algal blooms occur, (2) populations of grass carp that, through removal of littoral zone vegetation and feeding activities, lead to increased water column nutrient levels such that algal blooms occur, and (3) excessive growth of attached algae (periphyton) or attached filamentous algae on coarse substrates in stream riffle areas.
exotic species	yes	no	For purposes of Section 305(b) water quality assessments in Iowa, "exotic species" refers to a form "introduced into an area or ecosystem outside its historic or native geographic range; this includes both foreign (i.e., exotic) and transplanted species, and is used synonymously with "alien," "nonnative," and "introduced." Examples of exotic species in Iowa include <u>common carp</u> , <u>grass carp</u> , and the plant <u>purple loosestrife</u> . Scenarios that can lead to impairments due to "exotic species" include the following: (1) re-suspension of sediment and nutrients in a shallow lake by a large population of common carp such that turbidity and/or algal populations are increased to nuisance levels; (2) elimination of aquatic macrophytes from the littoral zone of a lake by grass carp such that the lake shifts from a clear-water to a turbid, phytoplankton-dominated (green) lake; and (3) the replacement of native wetland vegetation (e.g., grasses, sedges, cattails) with the exotic invasive purple loosestrife, thus degrading the habitat quality of the wetland for waterfowl and nutritional value of the wetland for wildlife.
flow alterations	yes	no	"Flow alterations" refer to human-related deviations from natural seasonal flow regimes that can adversely affect native biota. Flow alterations can result from several activities including water withdrawal for irrigation or water supplies, regulation of stream flow at dams, and drainage projects that lead to localized lowering of water tables such that lake/wetland water levels are adversely affected.
habitat alterations (other than flow alterations)	yes	no	"Habitat alterations" refer to manmade changes in the physical habitats of surface waters such that native aquatic biota may be adversely affected. When assessing impairments to Iowa surface waters for Section 305(b) reporting, "habitat alterations" refers primarily to impacts from (1) stream channelization (i.e., channel straightening), (2) removal of riparian vegetation, (3) pasturing of the riparian zone, and/or (4) streambank destabilization. All of these alterations tend to decrease the value of streams and rivers as high quality habitats for use by aquatic life through removal of

Cause Category	Historically Used?	Numeric Criteria?	Description
			important naturally-occurring habitat types (e.g., pools, riffles, sand bars, and snags). In addition, the alteration of aquatic habitat tends to increase the severity of impacts from other sources of pollution on aquatic life, especially the effects of siltation during low-flow periods.
metals	yes	yes	A general category that includes the following toxic metals: aluminum, antimony, arsenic, asbestos, beryllium, cadmium, chromium, copper, cyanide, lead, mercury, nickel, selenium, silver, thallium, zinc. All but aluminum are identified as "priority pollutants" under Section 307a of the Clean Water Act. Levels of toxic metals in Iowa waters are low. Impairments of Iowa waters for metals occur infrequently and tend to occur in rivers. Impairments are related to violations of chronic criteria to protect aquatic life uses from toxic metals. The occurrence of acutely toxic levels of toxic metals in Iowa surface waters is extremely rare.
nitrate	yes	yes	High levels of nitrate in drinking water can lead to infant methemoglobinemia (blue baby syndrome). This condition occurs as a result of ingestion of high levels of nitrate followed by the metabolism of nitrate to ammonia in the infant's digestive system. The conversion of nitrate to ammonia produces nitrite which can oxidize the iron atom in hemoglobin such that it can not carry oxygen. The lack of oxygen can give blood and oxygen-deficient tissues a bluish color. To protect against this condition, the U.S. EPA recommends that nitrate levels in water delivered by a public water supply to consumers should not exceed a maximum contaminant level (MCL) of 10 mg/l as nitrogen. The <i>Iowa Water Quality Standards</i> identify this 10 mg/l MCL as the water quality criterion to protect surface waters used as a source of a municipal water supply. At concentrations seen in surface waters, nitrate is not believed to be toxic to aquatic life; thus, there are no water quality criteria in the <i>Iowa Water Quality Standards</i> that apply to aquatic life uses.
nitrogen	yes	no	Nitrogen is an essential nutrient, is very abundant in the earth's atmosphere, and—like phosphorus—is implicated in eutrophication of surface waters such than excessive production of plant biomass occurs. Being considerably more abundant than phosphorus, nitrogen is much less often identified as a limiting (critical) nutrient in the eutrophication process. In water, nitrogen occurs in several forms (oxidation states) including nitrate, nitrite, and ammonia. Total nitrogen is defined as the sum of ammonia, nitrate, nitrite, and total Kjeldahl nitrogen (a measure of organic forms of nitrogen; e.g., in proteins). Total nitrogen is the measure most often proposed as an indicator of nutrient enrichment in surface waters and is the form proposed for inclusion into state water quality standards as a nutrient criterion. In Iowa waters, nitrate usually accounts for the majority of total nitrogen. Levels of total nitrogen in Iowa waters and in waters of other Corn Belt states are high relative to those in other states and are high relative to nutrient benchmark values for total nitrogen that have developed by nutrient criteria workgroups over the last decade (approximately 1 part per million for both rivers and lakes). Assuming that nitrate+nitrite concentrations approximate levels of total nitrogen in Iowa surface waters, the median level of nitrate+nitrite in the approximately 9,500 samples collected from 2000 through 2009 as part of Iowa DNR's ambient stream/river water quality monitoring network is 5.8 parts per million (ppm). Seventy-five percent of the samples had nitrate levels greater than 3.0 ppm (IDNR 2010).
noxious aquatic plants**	yes	no	"Noxious aquatic plants" refers to excessive growths of aquatic macrophytes or algae (e.g., bluegreen algae) that are known to interfere with recreational uses and be potentially harmful to human health as well as to the health of aquatic biota. Scenarios that can lead to impairments due to "noxious aquatic plants" include the following: dominance of a lakes' phytoplankton community by

Cause Category	Historically Used?	Numeric Criteria?	Description
			bluegreen algae.
nutrients	yes	no	<p>High levels of plant nutrients (primarily, nitrogen and phosphorus) indicate the potential for water quality problems in surface waters that result from excessive production of plant biomass. In lakes, high levels of nutrients can lead to excessive growth of aquatic plants, especially algae, which can interfere with recreational uses of a lake (e.g., boating, swimming, and fishing). Excessive plant growth can also lead to oxygen depletion of lake water through respiration related to bacterial decomposition of plant material and other organic matter that accumulates on the lake bottom. Severe cases of oxygen depletion can lead to fish kills. High levels of plant nutrients are generally attributed to agricultural nonpoint source pollution, to background levels in soil, and to naturally-occurring conditions, especially the internal nutrient recycling that occurs in the shallow glacial lakes of northern Iowa. Urban point sources and urban runoff, however, also contribute excessive amounts of nutrients to Iowa lakes with urban watersheds. Both the origin of high levels of plant nutrients and the nutrient concentrations that can impair aquatic life uses of Iowa's surface waters are poorly understood. Due to the natural fertility of Iowa's soils, levels of plant nutrients were likely relatively high prior to settlement in the mid-19th century (Menzel 1983). Application of fertilizers, however, especially for row crop agriculture, has increased nutrient levels in the state's surface waters over that during pre-settlement times. The threshold levels at which plant nutrients cause problems in Iowa's surface waters have not yet been identified. Thus, the <i>Iowa Water Quality Standards</i> does not contain water quality criteria for either levels of phosphorus or nitrogen related to protection for primary contact recreation (Class A) or for aquatic life (Class B) beneficial uses. Since 2004, IDNR has used a trophic state index to identify nutrient-related water quality problems in lakes due to poor water clarity caused by large populations of algae that are aesthetically objectionable and that thus suggest impairment of recreational uses. Algal impairment based on the trophic state index is the most commonly identified impairment at Iowa lakes. DNR, however, is currently (2010) proposing nutrient criteria to protect recreational uses in Iowa lakes and is in the process of developing numeric criteria for the nutrients nitrogen and phosphorus to protect aquatic life uses in Iowa's streams and rivers.</p>
oil and grease	no	no	<p>"Oil and grease" refers to adverse impacts to public water supplies or aquatic biota due to the presence of oils of petroleum or non-petroleum origin. Scenarios that can lead to impairments due to "oil and grease" include the following: (1) a fish kill caused by a spill of fuel oil and (2) adverse impacts to aquatic life resulting from contact of surface waters with coal tar waste.</p>
organic enrichment / low dissolved oxygen	yes	yes	<p>Impairments due to organic enrichment occur when the amount of organic material delivered to the waterbody exceeds the capacity of the stream to mineralize and assimilate this organic material with the result that levels of dissolved oxygen can fall below water quality criteria designed to protect aquatic life uses. In the absence of excessive inputs of oxygen-demanding organic material—as commonly measured through biochemical oxygen demand or "BOD"—streams, rivers, and lakes can process organic material without serious consequences to either chemical water quality or aquatic life. When inputs of organic materials exceed the stream or river's assimilative capacity, however, degradation of water quality will occur. The high rates of bacterial respiration resulting from the excessive amounts of organic material can lower the level of dissolved oxygen below that needed to support aquatic life. Most of the lakes with impacts due to organic enrichment are the relatively shallow natural lakes in north-central and northwest Iowa. Wind action at shallow lakes in summer</p>

Cause Category	Historically Used?	Numeric Criteria?	Description
			<p>tends to circulate lake water at all depths, thus resuspending sediments and nutrients that have settled to the bottom of the lake back into the water column. The increased levels of nutrients in the water column can increase plant production, usually in the form of algae. Continued resuspension of sediment and nutrients can lead to poor water transparency due to high levels of planktonic algae or due to high concentrations of suspended sediment. The rehigh levels of biological productivity in these lakes can lead to depletion of dissolved oxygen, and fish kills can occur. In temperate climates such as Iowa's, deeper lakes tend to thermally stratify during summer: a relatively cold and stagnant bottom layer of the lake (hypolimnion) becomes isolated from the relatively warm and wind-circulated surface layer (epilimnion) by a middle layer with a temperature gradient (metalimnion or thermocline). As summer progresses, bottom layers of stratified eutrophic lakes tend to become increasingly nutrient-rich and oxygen-poor. The isolation of this bottom layer, however, prevents movement of the poor-quality water to the surface layer of the lake. This isolation tends to improve the water quality of the surface layer of a lake that is used by aquatic life and is used for water-based recreation (e.g., swimming and water skiing). Water quality studies on Iowa lakes have shown that lakes with average depths greater than 13 feet tend to establish and maintain thermal stratification in summer and thus have better water quality than do shallower lakes (Bachmann et al. 1994).</p>
other inorganics	no	yes	<p>"Other inorganics" is a general cause category for inorganic substances that are not already included in a cause category.</p>
pathogens (pathogen indicators)	yes	yes	<p>"Pathogens," in the context of Section 305(b) reporting, actually refers to concentrations of typically non-pathogenic indicator bacteria (e.g., fecal coliforms or <i>E. coli</i>) in surface water samples. Iowa surface waters that support swimming, water skiing, and other primary body contact recreation that involves considerable risk of ingesting surface water are designated for one of several types of Class A (swimmable) use in the <i>Iowa Water Quality Standards</i>. Levels of fecal coliform bacteria and <i>E. coli</i> are monitored by DNR in rivers and lakes designated for Class A uses to <i>indicate</i> the health risks to persons using these waters for water-based recreation. Although typically not pathogenic, pathogen indicators such as fecal coliforms and <i>E. coli</i> are present in the intestines of warm-blooded animals and are commonly monitored by state environmental agencies to indicate the degree to which surface waters may contain waterborne pathogens (e.g., <i>Salmonella</i> and <i>Shigella</i>) that can cause disease in humans. "Pathogen indicators" (bacteria) is the most frequently identified impairment of Iowa streams and rivers. Despite the relatively high levels of indicator bacteria in Iowa streams and rivers, and despite the high numbers of impairments, reports of waterborne disease are extremely rare.</p>
PCBs	yes	yes	<p>Polychlorinated biphenyls are mixtures of up to 209 individual chlorinated organic compounds (congeners). There are no known natural sources of PCBs. PCBs are either oily liquids or solids that are colorless to light yellow. Some PCBs can exist as a vapor in air. PCBs have no known smell or taste. Many commercial PCB mixtures are known in the U.S. by the trade name Aroclor. PCBs have been used as coolants and lubricants in transformers, capacitors, and other electrical equipment because they don't burn easily and are good insulators. The manufacture of PCBs was stopped in the U.S. in 1977 because of evidence they build up in the environment and can cause harmful health effects. Products made before 1977 that may contain PCBs include old fluorescent lighting fixtures and electrical devices containing PCB capacitors, and old microscope and hydraulic oils (excerpted from ATSCR ToxFAQ: http://www.atsdr.cdc.gov/tfacts17.pdf). Levels of PCBs in</p>

Cause Category	Historically Used?	Numeric Criteria?	Description
			Iowa surface waters are too low to be detected in samples collected as part of ambient water quality monitoring. PCBs, however, like many chlorinated organic compounds, do accumulate (bioconcentrate) in animal tissue. In Iowa waters, the only Section 303(d) impairment caused by PCBs is their accumulation in fish tissue to levels that indicate the need to issue a fish consumption advisory (see http://www.iowadnr.gov/fish/news/consump.html). Levels of PCBs in Iowa fish and in fish nationwide, however, have declined greatly (by a factor of 100) since the banning PCB production in the United States in 1977.
pesticides	yes	yes	"Pesticides" refers to any substance, either currently or historically, used to kill plants, insects, algae, fungi, and other organisms; includes herbicides, insecticides, algacides, fungicides, and other substances. For purposes of 305(b)/303(d) reporting in Iowa, this category includes priority pesticides* (as defined in Section 307a of the Clean Water Act) as well as non-priority pesticides (e.g., cyanazine, and metolachlor).
pH	yes	yes	"pH" indicates the hydrogen ion concentration a water sample and indicates the intensity of an acid. The pH of natural waters is a measure of acid-base equilibrium achieved by the various dissolved compounds, salts, and gases. A pH of 7 is considered neutral (neither acidic nor basic). As the pH of waters decreases below 7, the waters become increasingly acidic. For example, the pH of tomatoes is 4.5, that of vinegar is approximately 2 and of battery acid is roughly 1 pH unit. As the pH increases above 7, the waters become increasingly basic. For example, the pH of baking soda is 8.3, that of ammonia is 11, and lye has a pH of 13. The pH scale varies logarithmically such that water with a pH of 5 is ten times more acidic (i.e., has ten times the hydrogen ion concentration) than water with a pH of 6. The ability of surface waters to resist changes in pH is called buffering capacity and is measured by alkalinity. The alkalinity of a surface water reflects the nature of the rocks within a drainage basin and is measured as milligrams of calcium carbonate (CaCO ₃) per liter (mg/l). Surface waters with high alkalinities resist lowering of pH values due, for example, to the addition of low-pH rainfall (acid precipitation). pH can have direct and indirect effects on aquatic life. Within a range of about pH 6.5 to 9, direct impact to aquatic life are minimal; outside of this range, adverse physiological impacts can occur and will increase as the pH deviates from this range. pH can also have indirect impacts on aquatic life as the toxicity of certain metals to aquatic life increases at lower pH and the toxicity of ammonia increases as pH levels increase. pH levels outside of the range of 6.5 to 9.0 can also impact swimmers by causing irritation to eyes (FWPCA 1968). Thus, because of the potential impacts to both aquatic life and primary contact recreation uses, the <i>Iowa Water Quality Standards</i> specify a range of pH values of 6.5 to 9.0 as protective of both aquatic life and primary contact recreation uses. Levels of pH in Iowa surface waters tends toward the basic side of neutral with lake pH values being somewhat higher than those found in rivers and streams. The median pH of over 9,000 stream/river samples collected from 2000-2009 was 8.2 units with over 90 percent of the samples greater than a pH of 7.8 units and with only 10 percent of the samples having a pH of greater than 8.6 (IDNR 2010). The median pH of almost 3,000 summer-season water samples collected from Iowa lakes from 2000-2007 was 8.6 units with over 90 percent of the samples having a pH of greater than 8.0 units; 17 percent of the samples had a pH greater than 9.0 units and thus are in violation of the Iowa water quality criterion. The tendency for lake pH values to be higher than rivers likely reflects the larger populations of algae in lakes versus rivers: the removal of carbon dioxide from the water column during algal photosynthesis results in an increase in pH levels.

Cause Category	Historically Used?	Numeric Criteria?	Description
phosphorus	yes	no	<p>Phosphorus is an essential nutrient for all living cells and functions in the storage and transfer of energy in living organisms and in their genetic systems. Igneous rock was the original source of phosphorus on earth; biotic sources of phosphorus (e.g., guano from sea birds) also exist. Phosphorus is highly reactive and is not found as a free element in Nature. In water, phosphorus can occur in several forms including dissolved and particulate. In addition, phosphorus concentrations in water can be reported in a number of ways depending on the type of sample analyzed (i.e., filtered versus unfiltered) and the type analytical methods used. (Sources: Wikipedia (http://en.wikipedia.org/wiki/Phosphorus) and Cole (1979)). IDNR's ambient stream/river and lake monitoring networks measure and report phosphorus as "total phosphorus as P." Although an essential nutrient and although not toxic at levels found in the aquatic environment, high levels of phosphorus in water can stimulate excessive production of plant biomass (for example, algae) such that adverse water quality impacts can occur. These impacts range from reduced water clarity due to algae suspended in the water column, excessive oxygen demand from bacterial mineralization of decomposing plant material, and production of large populations of cyanobacteria (blue-green algae) that can be aesthetically objectionable as well as potentially harmful to human health. Levels of total phosphorus in Iowa surface waters tend to be high relative to levels considered to be of concern. The median level of total phosphorus in the approximately 9,500 samples collected from 2000 through 2009 as part of Iowa DNR's ambient stream/river water quality monitoring network is 200 parts per billion (ppb) (IDNR 2010). Twenty-five percent of the samples had phosphorus levels greater than 340 ppb. Of the 131 Iowa lakes monitored from 2001 through 2009, 99 lakes (75%) had median phosphorus levels greater than 50 ppb. The summary statistics suggest that the majority of Iowa's rivers, streams, and lakes have levels of phosphorus above the nutrient benchmark values for total phosphorus that have developed by nutrient criteria workgroups over the last decade (approximately 50 ppb for lakes and 100 ppb for rivers). The <i>Iowa Water Quality Standards</i> does not contain water quality criteria for either levels of phosphorus or nitrogen related to protection for primary contact recreation (Class A) or for aquatic life (Class B) beneficial uses. Thus, despite the quite high levels of phosphorus in Iowa waters, very few impairments of Iowa waters have been specifically attributed to "nutrients," "phosphorus," or "nitrogen." Given the lack of numeric nutrient criteria, IDNR has used a trophic state index to identify nutrient-related water quality impacts in lakes (e.g., poor water clarity due to large populations of algae) that are aesthetically objectionable and that thus suggest impairment of recreational uses. Algal impairment based on the trophic state index is the most commonly identified impairment at Iowa lakes. DNR, however, is currently (2010) proposing nutrient criteria to protect recreational uses in Iowa lakes and is in the process of developing numeric criteria for the nutrients nitrogen and phosphorus to protect aquatic life uses in Iowa's streams and rivers.</p>
priority organics	yes	yes	<p>"Priority organics" are toxic organic pollutants listed in Section 307a of the federal Clean Water Act: "Priority organics" includes the following pollutant groups: chlorinated benzenes, chlorinated ethanes, chlorinated phenols, other chlorinated organics, haloethers, halomethanes, nitrosamines, non-chlorinated phenols, phthalate esters, polynuclear aromatic hydrocarbons (PAHs), pesticides and metabolites*, DDT and metabolites, polychlorinated biphenyls (PCBs), and other organics. For purposes of 305(b)/303(d) reporting in Iowa, this cause category does not include the following groups of priority organics: pesticides and metabolites, DDT and metabolites, or polychlorinated</p>

Cause Category	Historically Used?	Numeric Criteria?	Description
radiation (radium)	no	yes	<p>biphenyls (PCBs).</p> <p>Radiation is the energy emitted spontaneously in the process of decay of unstable atoms of radioisotopes. Sources of radiation include (1) the natural decay of primordial radioisotopes and their decay products and (2) manmade radioisotopes released into the environment beginning with testing and use of the atomic bomb in World War II. Radiation absorbed by plant and animal tissue may cause cellular and molecular damage that can adversely affect aquatic biota. Although routinely monitored for in Iowa groundwater monitoring networks, monitoring for radiation (radium) is not part of surface water monitoring networks in Iowa.</p>
siltation	yes	no	<p>Silt delivered to streams and rivers through nonpoint source runoff and/or through streambank erosion can degrade aquatic habitat through covering of coarse substrates, through deposition in pools, and through increasing the turbidity of the water. Siltation impacts in lakes refer to the erosion of soil particles by precipitation and movement of soil particles in runoff to lake basins where accumulation of silt occurs. The amount of silt delivered to Iowa's lakes relative to lake volume is an important factor in determining the quality of a lake for fishing, swimming and for use as a source of drinking water. Sedimentation is especially a problem for man-made lakes formed by dams placed across stream channels. Water quality impacts related to high rates of siltation/sedimentation include the delivery of excessive levels of plant nutrients (primarily phosphorus) to lakes, loss of lake volume, loss of surface area, a shortened useful life of the lake, interference with reproduction and growth of certain fish species, and impairments to recreational uses such as boating and fishing. While the delivery and accumulation of sediment is often the most serious problem in man-made lakes, it is generally less of a problem in the natural lakes of north-central and northwest Iowa. Natural lakes generally have much smaller watersheds relative to lake surface area, and their watersheds have less topographic relief and lower erosion rates than do lake watersheds in other regions of the state. Man-made lakes with low sedimentation rates tend to have clearer water and more productive fisheries than do lakes receiving large amounts of sediment. The man-made lakes in Iowa with the best water quality have relatively steep sides, small watersheds, and have well-controlled watersheds with a high percentage either in approved soil conservation practices or in non-crop land uses (e.g., pasture or forest) (see Hill 1981). Ideally, a man-made lake in Iowa would have a watershed-to-surface area ratio of from 20:1 up to 40:1. As watershed size increases relative to lake area, the more likely that the lake basin will be impacted (overloaded) with sediment delivered to the lake in rainfall runoff.</p>
sulfates	no	no	<p>Sulfate (SO_4^{2-}) is a naturally-occurring negatively-charged dissolved constituent of water and is one of several similar ions that combine to constitute "total dissolved solids." Sulfate may form salts with sodium, potassium, magnesium and other positively-charged ions. Sulfate is widely distributed in nature and may be present in natural waters at concentrations ranging from a few to several hundred milligrams per liter. At high levels (e.g., greater than 600 mg/l), sulfate in drinking water can have laxative effects on consumers. Levels of sulfate in Iowa surface waters are relatively low with a median concentration of 36 mg/l in the approximately 8,000 samples collected from 2000 through 2009 as part of Iowa DNR's ambient stream/river water quality monitoring network (IDNR 2010). Only 10% of these samples have had sulfate levels greater than 96 mg/l; the maximum concentration in these samples was 400 mg/l. IDNR has recently proposed water quality criteria to protect aquatic life from high levels of sulfate; the criteria depend on both hardness and the chloride concentrations</p>

Cause Category	Historically Used?	Numeric Criteria?	Description
			(see http://www.iowadnr.gov/water/standards/files/ws_fact.pdf). Lacking any water quality criteria for sulfate, there are no historical or current impairments for sulfate in Iowa surface waters. Sulfate levels in Iowa waters are sufficiently low that implementation of IDNR's proposed criteria is unlikely to result in aquatic life impairments due to sulfate.
suspended solids	yes	no	"Suspended solids" refers to the organic and inorganic particulate matter suspended in the water column. High levels of suspended solids in Iowa surface waters reduce water clarity and give a turbid or cloudy appearance to the water. Such material can originate from detritus carried by streams and rivers, atmospheric fallout, biological activity, chemical reactions, and re-suspension from bottom sediments as a result of current, wind/wave action, or movements of bottom-dwelling fish. The <i>Iowa Water Quality Standards</i> does not contain numeric aquatic life criteria for suspended solids
taste and odor	no	no	"Taste and odor" refer to the acceptability of drinking water to the user. Most taste and odor problems are related to the presence of phenolic compounds or to the presence of odor-producing organic substances produced by microorganisms or by human and industrial wastes.
thermal modifications	yes	yes	"Thermal modification" refers to a manmade deviation from natural seasonal water temperatures such that aquatic biota may be adversely affected. This deviation can include (1) addition of heat above physiological optimum levels of resident aquatic life, (2) the addition of heat such that state water quality standards are violated, or (3) the abrupt cessation of heated effluents during cooler seasons such that aquatic life cannot acclimate to the sudden change in ambient water temperature. Scenarios that can lead to impairments due to "thermal modifications" include the following: (1) discharge of heated effluent from power generating facilities such that ambient water temperatures violate water quality standards and (2) a fish kill caused by summer storm runoff with elevated water temperatures due to flow over super-heated impervious surfaces (streets, parking lots, etc) in urban areas. Criteria for water temperature are summarized in Table 7 of this document and can also be found in the <i>Iowa Water Quality Standards</i> (http://www.legis.state.ia.us/asp/ACODOCS/DOCS/567.61.pdf).
total dissolved solids / salinity / chlorides / sulfates	no	no	"Total dissolved solids" (TDS) refers to the concentration of inorganic salts, small amounts of organic material, and other dissolved materials in the water column. The principal inorganic anions dissolved in water are carbonates, chlorides, sulfates, and nitrates; the principal cations are calcium, magnesium, sodium, and potassium. Previous version of the <i>Iowa Water Quality Standards</i> contained a numeric criterion for TDS of 750 mg/l as part of "general water quality criteria." Recent changes in the Standards, however, have included replacement of the TDS criterion with separate criteria for chloride and sulfate with the goal of improved protection of aquatic life (see http://www.iowadnr.gov/water/standards/files/ws_fact.pdf).
total toxics	no	no	"Total toxics" refers to the cumulative adverse impact of toxic parameters from multiple groupings on water quality and aquatic biota.
turbidity	yes	no	For purposes of Section 305(b) assessments and Section 303(d) listings, "turbidity" refers to non-algal materials suspended in the water column, especially soil particles (silt or clay), that give the water a brown, cloudy appearance. Turbidity-related impairments due to planktonic algae (i.e., "green" water) are considered to be caused by "excessive algal growth/chlorophyll-a." Regardless of the cause, high levels of turbidity may suggest a water quality impairment. High levels of turbidity in surface waters, whether due to suspended algae or non-algal materials, can interfere with the growth

Cause Category	Historically Used?	Numeric Criteria?	Description
			<p>and reproduction of sight-feeding game fish (e.g., bluegill (<i>Lepomis macrochirus</i>), largemouth bass (<i>Micropterus salmoides</i>), and walleye (<i>Stizostedion vitreum</i>)), and excessive turbidity reduces the aesthetic appeal of surface waters for primary contact recreation such as swimming and water skiing. The primary sources of high turbidity in Iowa surface waters are (1) the resuspension of bottom sediments in shallow lakes through wind/wave action, (2) delivery of high amounts of silt and clay particles to the surface waters during precipitation runoff from agricultural areas, (3) contributions of silt and clay particles from erosion of stream banks or lake shorelines, or (4) bottom feeding fish (e.g., common carp (<i>Cyprinus carpio</i>) and bullheads (<i>Ameiurus</i> spp.) that increase turbidity through resuspension of sediment and nutrients during feeding and spawning activities. Surface waters that drain watersheds with certain types of clay-dominated soils may have chronic problems with turbidity regardless of the level of agricultural activity in the watershed. Historical evidence suggests that streams and rivers in the Missouri River drainage of southern and western Iowa had high levels of turbidity even during pre-settlement times. The presence of a turbidity tolerant fish fauna in these streams and rivers supports this assertion. Iowa surface waters with water quality problems due to high levels of turbidity are generally of three types: (1) man-made lakes in southern Iowa with relatively large watersheds having high rates of soil erosion (e.g., Bob White, Rock Creek, and Manteno lakes) and (2) shallow natural lakes of northern Iowa with high turbidities related to resuspension of silt and nutrients by bottom-feeding fish and/or wind/wave action (e.g., Ingham, Lower Gar, and North Twin lakes) and (3) streams and rivers with chronically high turbidities that may contribute to reduced aquatic diversity.</p>
unknown toxicity	yes	NA	<p>“Unknown toxicity” is identified as a cause of impairment when results of monitoring suggest some type of toxic impact but the identities of the substances causing toxicity are unknown. For example, results of a biological assessment that shows a complete lack of aquatic life in a stream strongly suggest the presence of toxic substances; the cause of impairment in such a case would be identified as “unknown toxicity.”</p>

* aldrin, dieldrin, chlordane, alpha-endosulfan, beta-endosulfan, endosulfan sulfate, endrin, endrin aldehyde, heptachlor, heptachlor epoxide, alpha BHC, beta BHC, gamma-BHC (lindane), delta-BHC, and toxaphene.

** Bluegreen algae (cyanobacteria) is considered a “noxious aquatic plant” by IDNR

References, Attachment 5:

- Bachmann, R.W., T.A. Hoyman, L.K. Hatch, and B.P. Hutchins. 1994. A classification of Iowa's lakes for restoration. Department of Animal Ecology, Iowa State University, Ames, Iowa. 517 p.
- Cole, G.A. 1979. Textbook of limnology. The C.V. Mosby Company. 426 p.
- FWPCA. 1968. Water quality criteria: report of the technical advisory committee to the Secretary of the Interior. Federal Water Pollution Control Administration, Washington D.C. 234 p.
(<http://www.eric.ed.gov/PDFS/ED046708.pdf>).
- Hill, K. 1981. Classification of Iowa lakes and their fish standing stocks. Iowa Conservation Commission.
- IAC. 2008. Chapter 567-61: water quality standards. Iowa Administrative Code
(<http://www.legis.state.ia.us/asp/ACODOCS/DOCS/567.61.pdf>).
- IDNR. 2010. Ambient monitoring program: water quality summary 2000-2009. Fact Sheet 2010-11. Geological and Water Survey Bureau, Iowa Department of Natural Resources
(<http://www.igsb.uiowa.edu/webapps/gsbpubs/pdf/WFS-2010-11.pdf>).
- Menzel, B.W. 1983. Agricultural management practices and the integrity of instream biological habitat. Pgs. 305-329 in: Schaller, F.W. and G.W. Bailey (eds): Agricultural management and water quality. Iowa State University Press, Ames. 472 p. <http://www.legis.state.ia.us/asp/ACODOCS/DOCS/567.61.pdf>