

Iowa Influenza Surveillance Network (IISN)

Influenza-like Illness (ILI) and Other Respiratory Viruses

Weekly Activity Report

For the week ending May 26, 2018 - Week 21

All data presented in this report are provisional and may change as additional reports are received



| Iowa Influenza Geographic Spread | | | | | | |
|----------------------------------|----------|-------|----------|------------|--|--|
| No Activity | Sporadic | Local | Regional | Widespread | | |

Note: See CDC activity estimates for definition www.cdc.gov/flu/weekly/overview.htm

| Quick Stats | |
|---|-----------------------|
| Percent of influenza rapid test positive | 3% (10/287) |
| Percent of RSV rapid tests positive | 4% (1/28) |
| Influenza-associated hospitalizations | 1/1886 inpatients |
| Percent of outpatient visits for ILI | 0.18% (baseline 1.9%) |
| Percent school absence due to illness | 1.33% |
| Number of long-term care influenza outbreaks | 0 |
| Number of schools with ≥10% absence due to illness | 0 |
| Influenza-associated mortality -all ages (Cumulative) | 271 |
| Influenza-associated pediatric mortality (Cumulative) | 0 |

Note: Deaths are considered influenza-associated when influenza is listed on the death certificate. This is an underestimate of influenza-related deaths. Cumulative mortality totals are from 10/1/2017 to the current week.

Iowa statewide activity summary:

This will be the last weekly report for the 2017-2018 surveillance season. Surveillance reports will be published monthly for the summer until weekly reports in the fall.


Influenza activity continues to decrease on most measures in Iowa, and the geographic spread is sporadic. For this reporting week, the State Hygienic Laboratory did not confirm any influenza viruses from submitted samples. There was one influenza-related hospitalization reported from sentinel hospitals during this reporting week. The proportion of outpatient visits due to influenza-like illness (ILI) was 0.18 percent, which is below the regional baseline of 1.9 percent. No long-term care influenza outbreaks were reported with first illness onset in this reporting week. No schools reported 10 percent or more illness this reporting week. There have been 271 influenza-related deaths reported this season. Surveillance sites reported detecting the following non-influenza respiratory illnesses with the most frequency: 77 rhinovirus/enterovirus, 48 parainfluenza virus type 3, and 15 adenovirus.

International activity summary - (WHO):

Influenza activity returned to inter-seasonal levels in most of the countries in the temperate zone of the northern hemisphere. Activity increased in some countries in tropical America. In the temperate zone of the southern hemisphere, influenza activity increased but remained below the seasonal thresholds in most countries. Worldwide, seasonal influenza subtypes A and B accounted for approximately the same proportion of influenza detections.

Visit <u>www.who.int/influenza/surveillance monitoring/updates/latest update GIP surveillance/en/</u> for more information. It was last updated 5/28/2018.

National activity summary - (CDC)-Last Updated in Week 20:



Synopsis: During week 20 (May 13-19, 2018), influenza activity continued to decrease in the United States.

Viral Surveillance: Overall, influenza A(H3) viruses have predominated this season. Since early March, influenza B viruses have been more frequently reported than influenza A viruses. The percentage of respiratory specimens testing positive for influenza in clinical laboratories decreased.

Pneumonia and Influenza Mortality: The proportion of deaths attributed to pneumonia and influenza (P&I) was below the system-specific epidemic threshold in the National Center for Health Statistics (NCHS) Mortality Surveillance System.

Influenza-associated Pediatric Deaths: One influenza-associated pediatric death was reported.

Influenza-associated Hospitalizations: A cumulative rate of 106.6 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported.

Outpatient Illness Surveillance: The proportion of outpatient visits for influenza-like illness (ILI) was 1.2%, which is below the national baseline of 2.2%. All 10 regions reported ILI below region-specific baseline levels. New York City, the District of Columbia, Puerto Rico and all 50 states experienced minimal ILI activity.

Geographic Spread of Influenza: The geographic spread of influenza in one state was reported as widespread; Guam, Puerto Rico and one state reported regional activity; seven states reported local activity; the District of Columbia, the U.S. Virgin Islands and 37 states reported sporadic activity; and four states reported no influenza activity.

Detailed information can be found online at www.cdc.gov/flu/weekly/.

Laboratory surveillance program:

The State Hygienic Laboratory (SHL) is the primary lab for influenza testing and reporting in Iowa. SHL reports the number of tests performed and the type and sub-type of positive tests to the influenza surveillance network daily.

Table 1: Influenza A viruses detected by SHL by age group


| | CURRENT WEEK | | | CURRENT WEEK CUMULATIVE (10/1/17- CURRENT WEEK) | | | | EK) | |
|--------------|------------------|-----------|-----------------------------|---|------------------|---------------|--------------|-----------------------------|-----------|
| Age Group | A(H1N1) pdm09 | A (H3) | Pending/ Not subtyped | Total | A(H1N1) pdm09 | A variants | A (H3) | Pending/ Not subtyped | Total |
| 0-4 | 0 | 0 | 0 | 0 (0%) | 10 | 0 | 73 | 1 | 84 (8%) |
| 5-17 | 0 | 0 | 0 | 0 (0%) | 8 | 0 | 82 | 1 | 91 (9%) |
| 18-24 | 0 | 0 | 0 | 0 (0%) | 11 | 1 | 128 | 0 | 140 (14%) |
| 25-49 | 0 | 0 | 0 | 0 (0%) | 11 | 1 | 77 | 1 | 90 (9%) |
| 50-64 | 0 | 0 | 0 | 0 (0%) | 10 | 0 | 113 | 1 | 124 (12%) |
| >64 | 0 | 0 | 0 | 0 (0%) | 7 | 0 | 489 | 3 | 499 (49%) |
| Total | 0 (0%) | 0 (0%) | 0 (0%) | 0 | 57 (6%) | 2 (0%) | 962 (94%) | 7 (1%) | 1028 |

Note: Cell counts of three or less are sometimes suppressed to protect confidentiality. Totals by age may not add up to totals by subtype/lineage due to missing age information. Only cases of Iowa residents are included. Specimens listed as "not subtyped" were not able to be subtyped due to weak positive lab results. This can be due to poor collection, timing of collection or stage of infection. Influenza A variants detected in 2017-18 include A(H1N1)v and A(H3N2)v.

Table 2: Influenza B viruses detected by SHL by age group

| | | | | -7 -0 - 0 | • | | | |
|-------|--------------|----------|---------|-----------|----------------|------------|---------|-----------|
| | CURRENT WEEK | | | CUMU | LATIVE (10/1/1 | L7- CURREN | T WEEK) | |
| Age | Victoria | Yamagata | Lineage | Total | Victoria | Yamagata | Lineage | Total |
| Group | Lineage | Lineage | Pending | Total | Lineage | Lineage | Pending | Total |
| 0-4 | 0 | 0 | 0 | 0 (0%) | 0 | 22 | 1 | 23 (5%) |
| 5-17 | 0 | 0 | 0 | 0 (0%) | 0 | 83 | 1 | 84 (17%) |
| 18-24 | 0 | 0 | 0 | 0 (0%) | 1 | 54 | 0 | 55 (11%) |
| 25-49 | 0 | 0 | 0 | 0 (0%) | 1 | 60 | 0 | 61 (13%) |
| 50-64 | 0 | 0 | 0 | 0 (0%) | 0 | 73 | 1 | 74 (15%) |
| >64 | 0 | 0 | 0 | 0 (0%) | 0 | 185 | 0 | 185 (38%) |
| Total | 0 | 0 | 0 | 0 | 2 | 477 | 3 | 402 |
| Total | (0%) | (0%) | (0%) | 0 | (0%) | (99%) | (1%) | 482 |

Note: Cell counts of three or less are sometimes suppressed to protect confidentiality. Totals by age may not add up to totals by subtype/lineage due to missing age information. Only cases of Iowa residents are included.




Rapid influenza and RSV test surveillance:


The State Hygienic Laboratory (SHL) runs a weekly web-based survey program where laboratorians report the number of influenza and respiratory syncytial virus (RSV) rapid tests performed and the number of tests positive. This table includes only the number of patients tested for influenza or RSV at laboratory surveillance sites throughout the state. This table does not provide case counts.

Table 3: Percent of influenza rapid tests positive and number of tests performed by region for the present week

| REGION* | | RAPID IN | FLUENZA TES | TS | F | RAPID RSV TES | TS |
|--------------------|--------|----------|-------------|------------|--------|---------------|------------|
| REGION | Tested | Flu A | Flu B | % Positive | Tested | Positive | % Positive |
| Region 1 (Central) | 84 | 0 | 2 | 2 | 2 | 0 | 0 |
| Region 2 (NE) | 9 | 0 | 0 | 0 | 5 | 0 | 0 |
| Region 3 (NW) | 10 | 0 | 0 | 0 | 6 | 0 | 0 |
| Region 4 (SW) | 8 | 1 | 0 | 13 | 2 | 0 | 0 |
| Region 5 (SE) | 26 | 5 | 0 | 19 | 5 | 0 | 0 |
| Region 6 (Eastern) | 150 | 1 | 1 | 1 | 8 | 1 | 13 |
| Total | 287 | 7 | 3 | 3 | 28 | 1 | 4 |

Note: See map in the school section for a display of the counties in each region.







Influenza-associated hospitalizations:

Sentinel hospitals that participate in IISN voluntarily track and report the number of influenza-associated hospitalizations and the total number of inpatients each week. Iowa hospitals interested in joining this surveillance program should contact Andy Weigel at 515-322-1937 or andy.weigel@idph.iowa.gov or more information.

Table 4: Number of influenza-associated hospitalization reported by age group

| | • | 1 1001 |
|-----------|--------------|-------------------------------------|
| AGE | CURRENT WEEK | CUMULATIVE (10/1/17 – CURRENT WEEK) |
| Age 0-4 | 0 | 124 |
| Age 5-24 | 0 | 99 |
| Age 25-49 | 0 | 169 |
| Age 50-64 | 0 | 344 |
| Age >64 | 1 | 1154 |
| Total | 1 | 1890 |






Outpatient health care provider surveillance program (ILINet):

Outpatient health care providers who participate in the ILINet (a national influenza surveillance program) report the number of patients seen with influenza-like illness and the total number of patient visits each week. This system is a key part of Iowa's influenza surveillance. Iowa health care providers interested in joining this surveillance program should contact Andy Weigel at 515-322-1937 or andy.weigel@idph.iowa.gov for more information.

Table 5: Outpatient visits for influenza-like illness (ILI)


| Week | Age 0- 4 | Age 5- 24 | Age 25- 49 | Age 50- 64 | Age > 64 | Total ILI | ILI Percent |
|------------------------|-------------|--------------|---------------|---------------|----------|-----------|----------------|
| Week 19, ending May 12 | 2 | 1 | 0 | 0 | 2 | 5 | 0.26 |
| Week 20, ending May 19 | 1 | 1 | 1 | 0 | 3 | 6 | 0.36 |
| Week 21, ending May 26 | 1 | 0 | 1 | 0 | 1 | 3 | 0.18 |

Note: Influenza-like Illness is defined as a fever of ≥100° F as well as cough and/or sore throat.



Seasonal influenza vaccination:

Seasonal influenza vaccination data in Iowa is based on doses reported to the Iowa Immunization Registry Information System (IRIS). IRIS is a confidential, computerized, population-based system that tracks immunization for children, adolescents and adults who are seen in a variety of public and private healthcare provider sites throughout the state of Iowa. For more information on the immunization data, contact Kim Tichy, IRIS coordinator at 515-281-4288 or kimberly.tichy@idph.iowa.gov.




Note: The data for the 2017-2018 season is provisional due to the lag between the vaccine administration date and the date reported to IRIS.

Long-term care outbreaks:

Table 6: Number of long-term care outbreaks investigated


| REGION | CURRENT WEEK | CUMULATIVE (10/1/17 – CURRENT WEEK) |
|--------------------|--------------|--|
| Region 1 (Central) | 0 | 21 |
| Region 2 (NE) | 0 | 12 |
| Region 3 (NW) | 0 | 5 |
| Region 4 (SW) | 0 | 13 |
| Region 5 (SE) | 0 | 19 |
| Region 6 (Eastern) | 0 | 20 |
| Total | 0 | 90 |

Note: See map in the school section for a display of the counties in each region.



School surveillance program

IDPH monitors illnesses in schools from two different types of reporting: 10 percent school absence reports and weekly sentinel illness reporting. Iowa schools (K-12) track and report (including non-influenza illnesses) when the number of students absent with illness reaches or exceeds 10 percent of total student enrollment. Iowa sentinel schools that participate in IISN voluntarily track and report absence due to all illness and the total enrollment each week. This data provides excellent trends for influenza activity as well as age-specific information used to target vaccination efforts and messages.



^{*}School data not reported for week 52 due to holiday closings

^{**}School data may underestimate influenza illness for week 11 when many schools closed for spring break


Table 7: Number of schools reporting >10% absenteeism due to any illness

| REGION | CURRENT WEEK | CUMULATIVE (10/1/17 – CURRENT WEEK)* |
|--------------------|---------------------|--------------------------------------|
| Region 1 (Central) | 0 | 78 |
| Region 2 (NE) | 1 | 35 |
| Region 3 (NW) | 1 | 26 |
| Region 4 (SW) | 0 | 25 |
| Region 5 (SE) | 0 | 36 |
| Region 6 (Eastern) | 0 | 106 |
| Total | 2 | 306 |

^{*}School data may underestimate influenza illness for week 11 when many schools closed for spring break

Note: A regional map with the total of schools by county that report at least 10 percent of students absent due to illness for this current reporting week is displayed below (region numbers and abbreviations are displayed in large black font near the middle of each region). Current week totals only include the first report for a school in the week, but cumulative totals may include multiple reports for the same school from different weeks.

Number of schools with at least 10 percent of students absent, by county and region




Non-influenza respiratory viruses:

The State Hygienic Laboratory (SHL) runs a weekly web-based survey program where laboratorians report the number of positive tests for non-influenza respiratory viruses. This table also includes the positive non-influenza virus tests reported from the Dunes Medical Laboratories at Mercy Medical Center in Sioux City. The table includes only the number of positive tests at laboratory surveillance sites throughout the state. The table does not provide case counts.

Table 8: Number of positive results for non-influenza respiratory virus collected by SHL, Mercy Dunes in Sioux City and Iowa Methodist Medical Center

| Viruses | CURRENT WEEK | CUMULATIVE (10/1/17 – CURRENT WEEK) |
|-----------------------------------|--------------|--|
| Adenovirus | 15 | 747 |
| Parainfluenza Virus Type 1 | 0 | 488 |
| Parainfluenza Virus Type 2 | 1 | 12 |
| Parainfluenza Virus Type 3 | 48 | 381 |
| Parainfluenza Virus Type 4 | 0 | 71 |
| Rhinovirus/Enterovirus | 77 | 3448 |
| Respiratory syncytial virus (RSV) | 4 | 2362 |
| Human metapneumovirus (hMPV) | 10 | 1349 |
| Coronavirus | 1 | 1599 |
| Total | 156 | 10457 |



Other resources:

Vaccine:

Influenza vaccine recommendation: idph.iowa.gov/immtb/immunization/vaccine

CDC vaccine information: www.cdc.gov/flu/faq/flu-vaccine-types.htm

Vaccine finder: http://vaccinefinder.org/

Neighboring states' influenza information:

Illinois: www.dph.illinois.gov/topics-services/diseases-and-conditions/influenza/surveillance

Minnesota: www.health.state.mn.us/divs/idepc/diseases/flu/stats/index.html

Missouri: health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php

South Dakota: doh.sd.gov/diseases/infectious/flu/

Wisconsin: flu.wisconsin.gov/