

Iowa Department of Public Heath | Center for Acute Disease Epidemiology | <u>West Nile Virus Website</u> All data presented in this report are provisional and may change as additional reports are received Date Issued: March 2, 2020

# West Nile Virus (WNV)

WNV is endemic in Iowa and activity usually peaks in late summer and early fall. IDPH works in collaboration with Local Public Health (LPH) and other appropriate partners to investigate all reported cases.

In 2018, Iowa experienced an increase in WNV activity and 104 human cases were identified. This is the highest number since 2003. Thus far in 2019, five human cases of WNV and one presumptive viremic blood donor have been identified. Three horses and eight mosquito samples have tested positive for WNV [Table 1].


				Mosquitoes	
County	Human	Blood Donor	Horse	Culex pipiens group	Culex restuans
Audubon	1	0	0	0	0
Davis	0	0	1	0	0
Harrison	1	0	0	0	0
Humboldt	0	1	0	0	0
Mitchell	0	0	1	0	0
Polk	0	0	0	7	1
Sioux	1	0	0	0	0
Union	0	0	1	0	0
Woodbury	2	0	0	0	0
Total	5	1	3	7	1

## Figure 1. 2019 West Nile virus case count and incidence rate by county of residence.





Figure 2. WNV disease cases reported to IDPH, by week of onset-Iowa, 2019



## National WNV Activity:

As of January 7<sup>th</sup>, 618 counties from 47 states and the District of Columbia have reported WNV activity to ArboNET for 2019, including 43 states and the District of Columbia with reported WNV human infections (i.e., disease cases or viremic blood donors) and four additional states with reported WNV activity in non-human species only (i.e., veterinary cases, mosquito pools, dead birds, or sentinel animals) [Figure 3].

To date, 917 human WNV disease cases have been reported from 261 counties in 43 states and the District of Columbia. Of the 917 reported cases, 607 (66%) were classified as neuroinvasive disease (e.g., meningitis or encephalitis) and 310 (34%) were classified as non-neuroinvasive disease [Figure 4]. Dates of illness onset for cases ranged from January-December [Figure 5].

Overall, 100 WNV PVD's has been reported from 23 states.





\*WNV human disease cases or presumptive viremic blood donors. Presumptive viremic blood donors have a positive screening test which has not necessarily been confirmed.

†WNV veterinary disease cases, or infections in mosquitoes, birds, or sentinel animals



#### Figure 4. WNV neuroinvasive disease incidence\* reported to ArboNET, by state - United States, 2019 (as of January 7, 2020)





Figure 5. WNV disease cases reported to ArboNET, by month of onset\*- United States, 2019 (as of January 7, 2020)



\*Cases missing onset date (n=13)



# **Mosquito Surveillance**


IDPH in collaboration with the State Hygienic Laboratory (SHL), Iowa State University (ISU), and local public environmental health partners conducts ecological surveillance in 16 counties across the state by monitoring mosquitoes and testing for WNV infected populations.

	# of Samples		
Species	Tested	WNV Negative	WNV Positive
Cx. pipiens	276	276	0
Cx. pipiens group	868	861	7
Cx. tarsalis	157	157	0
Cx. restuans	591	590	1
Cx. territans	39	39	0
Cx. erraticus	1	1	0
Cx. salinarius	11	11	0
Cx. species	2	2	0
Ae. japonicus	0	0	0
An. punctipennis	0	0	0
Ae. atropalpus	0	0	0
Ae. sticticus	0	0	0
Ae. triseriatus	2	2	0
Total	1947	1939	8

#### Table 2. 2019 mosquitoes tested for West Nile virus

In addition to viral testing for WNV, the population of mosquitoes in Iowa is monitored through trapping activities. All trapped mosquitoes are sorted by species. The figure [Figure 7] below shows where and when Aedes albopictus mosquitoes were detected 2017-2019.

#### Figure 7. Aedes albopictus identified in Iowa, 2017-2019





# **Dengue Fever**


Dengue is a disease caused by any one of four related viruses, which are passed by the bite of an infected *Aedes aegypti* or *Aedes albopictus* mosquito. Infection with one of the four viruses does not protect against the others and consecutive infections put people at greater risk of developing dengue hemorrhagic fever (DHF).

Dengue is not found in Iowa. Cases are in travelers and immigrants returning from parts of the world where dengue transmission occurs. Nine cases of dengue have been reported in Iowa, thus far in 2019. In 2018, five cases of dengue were reported to IDPH.

## National Dengue Activity:

As of January 8th, 45 states and three territories have reported dengue cases to ArboNET for 2019 [Figure 6].

#### Figure 6. Laboratory-positive travel-associated and locally-acquired dengue cases from the 50 states and five territories— United States, 2019 (as of January 8, 2020)



# Chikungunya

Chikungunya is a viral disease that is spread to people by the bite of an infected *Aedes aegypti* and *Aedes albopictus* mosquito. Mosquitoes become infected when they feed on a person already infected with this virus.

Two cases of chikungunya virus disease have been reported in Iowa. Cases are in travelers and immigrants returning from parts of the world where chikungunya transmission occurs. In 2018, no cases of chikungunya were reported in Iowa.

## Malaria

Malaria is a serious and sometimes fatal disease caused by a parasite that commonly infects *Anopheles* mosquitoes. Malaria is spread to humans by the bite of the infected female mosquito. Only *Anopheles* mosquitoes can transmit malaria and they must have been infected through a previous blood meal taken from an infected person.

Twenty-three cases of malaria have been reported in Iowa. Cases are in travelers and immigrants returning from parts of the world where malaria transmission occurs. In 2018, 22 cases of malaria were reported to IDPH.



# **Rocky Mountain spotted fever (RMSF)**

American dog ticks are carriers of *Rickettsia rickettsii*, the bacteria that causes RMSF. The American dog tick is the most common species of tick in Iowa and can be found in every county in the state. The tick is most active late March through August.

Eleven cases of RMSF have been reported in Iowa. In 2018, 22 cases of RMSF were reported to IDPH.

# Ehrlichiosis/Anaplasmosis

There are at least three species of bacteria responsible for ehrlichiosis/anaplasmosis in the United States: *Ehrlichia chaffeensis, Ehrlichia ewingii, and Anaplasma phagocytophilum*. Ehrlichiae are transmitted by the bite of an infected lone star tick (*Amblyomma americanum*) which is found in Iowa. *A. phagocytophilum* is transmitted by the bite of an infected blacklegged tick (or deer tick, *Ixodes scapularis*) in Iowa. The clinical signs and symptoms of these infections are similar.

Thirty-three cases of ehrlichiosis/anaplasmosis have been reported in Iowa. In 2018, 27 cases of ehrlichiosis/anaplasmosis were reported to IDPH.

## **Babesiosis**

Babesiosis is caused by microscopic parasites that infect red blood cells. Most human cases in the United States are caused by the parasite *Babesia microti*. *Babesia microti* is spread by the blacklegged tick (or deer tick, *Ixodes scapularis*). The parasite typically is spread by the young nymph stage of the tick. They are most common during the warm months of spring and summer in areas with woods, brush, or grass.

One case of babesiosis has been reported in Iowa. In 2018, no cases of babesiosis were reported to IDPH.

#### Lyme

Lyme disease is caused by *Borrelia burgdorferi* and in Iowa is transmitted to humans by the bite of an infected tick, the blacklegged tick (or deer tick, *Ixodes scapularis*). Ticks are most likely to spread the Lyme disease bacterium during their pre-adult stage (nymph). They are most common between May and July and found in tall grasses and brush of wooded areas.

As of March 2<sup>nd</sup>, 302 confirmed and probable cases of Lyme disease have been reported in Iowa [Figure 8]. In 2018, 284 cases of Lyme disease were reported to IDPH.

