## IOWA PRENATAL CARE BARRIERS PROJECT DATA FROM 2018 SURVEYS

2018 Respondents from All Hospitals in Iowa: 20,035

2018 Respondents with Babies Born in Iowa County: 75\*

\*Because of the small number of cases, data may not be representative of the entire population of new mothers whose babies were in this county.

| Demographic Indicators  | State | County |
|---|-------|--------|
| Mother's Age  |       |        |
| Less than 18 years  | 01%   | 01%    |
| 18-19 years   | 03%   | 04%    |
| 20-25 years   | 24%   | 07%    |
| 26-30 years   | 37%   | 41%    |
| 31-35 years   | 26%   | 35%    |
| More than 35 years  | 09%   | 12%    |
| Mother's Ethnicity  |       |        |
| Hispanic  | 08%   | 05%    |
| Non-Hispanic  | 92%   | 95%    |
| White   | 90%   | 99%    |
| Black   | 06%   | <01%   |
| Asian or Pacific Islander   | 03%   | <01%   |
| American Indian or Native Alaskan   | 01%   | 01%    |
| Socioeconomic Indicators  | State | County |
| Mother's Education  |       |        |
| Less than high school   | 07%   | 04%    |
| High school   | 21%   | 19%    |
| Some college  | 33%   | 31%    |
| College graduate (Bachelor's degree)  | 27%   | 28%    |
| Graduate school   | 12%   | 19%    |
| Household Income  |       |        |
| Less than \$10,000  | 11%   | 03%    |
| \$10,000 - \$19,999   | 08%   | 07%    |
| \$20,000 - \$29,999   | 09%   | 06%    |
| \$30,000 - \$39,999   | 08%   | 04%    |
| \$40,000 - \$49,999   | 08%   | 04%    |
| \$50,000 or more  | 56%   | 76%    |
| Information Received in 12 Months Prior to Becoming Pregnant  | State | County |
| Doctor, Nurse, or Other Health Care Provider Talked About:  |       |        |
| Taking folic acid or a multivitamin   | 46%   | 52%    |
| Smoking   | 35%   | 44%    |
| Drinking alcohol  | 32%   | 39%    |
| Your weight   | 29%   | 31%    |
| Regular exercise  | 33%   | 36%    |
| How romantic relationships affect your health   | 14%   | 19%    |
| Oral health   | 21%   | 29%    |
| Did not see a doctor, nurse, or other health care professional for a check-up in the 12 months prior to becoming pregnant | 33%   | 35%    |

| o Birth Control was Used at Time of Conception  nintended Pregnancy and Did NOT Use Birth Control  an to return to work outside of home following birth of this baby | 87%<br>70%<br>74%<br>26% | 93%<br>76% |
|--|--------------------------|------------|
| an to return to work outside of home following birth of this baby  | 74%                      | 76%        |
| s  |                          |            |
|  |                          |            |
|  | 26%                      | 80%        |
|  |                          | 20%        |
| pes worksite have designated room/location (not counting bathroom stalls) r mothers to breastfeed or pump/express their breast milk                                  |                          |            |
| S  | 55%                      | 55%        |
|  | 28%                      | 31%        |
| on't Know  | 17%                      | 15%        |
| pes worksite have written policy about breastfeeding?  |                          |            |
| S  | 22%                      | 20%        |
|  | 28%                      | 20%        |
| on't Know  | 50%                      | 60%        |
| noked within the past 2 years  |                          |            |
| S  | 21%                      | 20%        |
|  | 79%                      | 80%        |
| garettes Smoked Per Day During Last 3 Months of Pregnancy  |                          |            |
| ro cigarettes per day  | 53%                      | 47%        |
| 10 per day   | 41%                      | 33%        |
| -20 per day  | 06%                      | 20%        |
| ore than 20 per day  | 01%                      | <01%       |
| onsumed alcohol in the past 2 years  |                          |            |
| S  | 70%                      | 80%        |
|  | 29%                      | 20%        |
| coholic Drinks During Last 3 Months of Pregnancy   |                          |            |
| ro drinks per week   | 98%                      | 96%        |
| -2 drinks per week   | 02%                      | 04%        |
| 5 drinks per week  | <01%                     | <01%       |
| 10 drinks per week   | <01%                     | <01%       |
| ore than 10 drinks per week  | <01%                     | <01%       |
| enatal Care  | State                    | County     |
| other received prenatal care as early in the pregnancy as wanted   |                          |            |
| S  | 95%                      | 97%        |
|  | 05%                      | 03%        |
| here mother went for most of prenatal care   |                          |            |
| ivate doctor's office, clinic or OBGYN   | 78%                      | 85%        |
| idwife's office or home office from a midwife  | 07%                      | 04%        |
| ıblic clinic   | 15%                      | 10%        |
| her  | 01%                      | <01%       |

|  | State                           | County                          |
|--|---------------------------------|---------------------------------|
| HIV/AIDS Testing During Pregnancy  |                                 |                                 |
| Yes  | 45%                             | 49%                             |
| No   | 28%                             | 24%                             |
| No, Testing was offered, but did not want to be tested   | 02%                             | 03%                             |
| Don't know   | 26%                             | 24%                             |
| Received Flu Shot During Pregnancy   |                                 |                                 |
| Yes  | 65%                             | 78%                             |
| No, Received flu shot before became pregnant   | 06%                             | 07%                             |
| No, Did not want flu shot during pregnancy   | 12%                             | 07%                             |
| No, Doctor said to wait until after pregnancy  | 01%                             | 01%                             |
| No, Do not ever get flu shots  | 13%                             | 07%                             |
| No, Flu shot was not available   | 02%                             | <01%                            |
|  |                                 |                                 |
| Hospital Services  | State                           | County                          |
| Hospital Services  Hospital Staff Encouraged Breastfeeding During First 24 Hours After   | State                           | County                          |
| ·  | State                           | County                          |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After  | State<br>85%                    | County<br>89%                   |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby  |                                 | <u> </u>                        |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby Yes  | 85%                             | 89%                             |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby Yes No   | 85%<br>13%                      | 89%<br>09%                      |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby Yes No Don't remember  | 85%<br>13%                      | 89%<br>09%                      |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby Yes No Don't remember  During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other   | 85%<br>13%                      | 89%<br>09%                      |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby Yes No Don't remember  During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other Healthcare Professional Provided Information or Scheduled a Time to Discuss:  | 85%<br>13%<br>02%               | 89%<br>09%<br>01%               |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby Yes No Don't remember  During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other Healthcare Professional Provided Information or Scheduled a Time to Discuss: Breastfeeding or breastfeeding techniques  | 85%<br>13%<br>02%               | 89%<br>09%<br>01%               |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby Yes No Don't remember  During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other Healthcare Professional Provided Information or Scheduled a Time to Discuss: Breastfeeding or breastfeeding techniques Action steps to decrease frustration when infant cries | 85%<br>13%<br>02%<br>83%<br>43% | 89%<br>09%<br>01%<br>84%<br>45% |