IOWA PRENATAL CARE BARRIERS PROJECT DATA FROM 2014 SURVEYS

2014 Respondents from All Hospitals in Iowa: 24,696 2014 Respondents with Babies Born in **Hancock**: 69*

^{*}Because of the small number of cases, data may not be representative of the entire population of new mothers whose babies were in this county.

| Demographic Indicators | State | County |
|--|---|--|
| Mother's Age | | |
| Less than 18 years | 01% | 01% |
| 18-19 years | 04% | 03% |
| 20-25 years | 28% | 26% |
| 26-30 years | 35% | 45% |
| 31-35 years | 25% | 20% |
| More than 35 years | 07% | 04% |
| Mother's Ethnicity | | |
| Hispanic | 07% | 01% |
| Non-Hispanic | 93% | 99% |
| White | 91% | 99% |
| Black | 05% | 01% |
| Asian or Pacific Islander | 03% | 00% |
| American Indian or Native Alaskan | 01% | 00% |
| | | |
| Socioeconomic Indicators | Ctata | County |
| | State | County |
| Mother's Education | State | County |
| Mother's Education Less than high school | 08% | 06% |
| | | |
| Less than high school | 08% | 06% |
| Less than high school High school | 08% 19% | 06% 17% |
| Less than high school High school Some college | 08% 19% 36% | 06% 17% 52% |
| Less than high school High school Some college College graduate (Bachelor's degree) | 08% 19% 36% 27% | 06% 17% 52% 25% |
| Less than high school High school Some college College graduate (Bachelor's degree) Graduate school | 08% 19% 36% 27% | 06% 17% 52% 25% |
| Less than high school High school Some college College graduate (Bachelor's degree) Graduate school Household Income | 08% 19% 36% 27% 10% | 06% 17% 52% 25% 00% |
| Less than high school High school Some college College graduate (Bachelor's degree) Graduate school Household Income Less than \$10,000 | 08% 19% 36% 27% 10% | 06% 17% 52% 25% 00% |
| Less than high school High school Some college College graduate (Bachelor's degree) Graduate school Household Income Less than \$10,000 \$10,000 - \$19,999 | 08% 19% 36% 27% 10% | 06% 17% 52% 25% 00% |
| Less than high school High school Some college College graduate (Bachelor's degree) Graduate school Household Income Less than \$10,000 \$10,000 - \$19,999 \$20,000 - \$29,999 | 08% 19% 36% 27% 10% 14% 10% | 06% 17% 52% 25% 00% 09% 12% 06% |

| Information Received in 12 Months Prior to Becoming Pregnant | State | County |
|---|------------|--------|
| Doctor, Nurse, or Other Health Care Provider Talked About: | | |
| Taking folic acid or a multivitamin | 47% | 39% |
| Smoking | 37% | 28% |
| Drinking alcohol | 32% | 28% |
| Your weight | 30% | 25% |
| Regular exercise | 33% | 23% |
| How romantic relationships affect your health | 13% | 12% |
| Did not see a doctor, nurse, or other health care professional for a check-up | 34% | 39% |
| in the 12 months prior to becoming pregnant | | |
| | | |
| | . . | |
| Behavioral Indicators During Pregnancy | State | County |
| No Birth Control was Used at Time of Conception | 86% | 94% |
| Unintended Pregnancy and Did NOT Use Birth Control | 68% | 79% |
| Official Control | 0070 | 7570 |
| Plan to return to work outside of home following birth of this baby | | |
| Yes | 72% | 82% |
| No | 28% | 18% |
| | | |
| Does worksite have designated room/location (not counting bathroom | | |
| stalls) for mothers to breastfeed or pump/express their breast milk | 100/ | 400/ |
| Yes | 48% | 48% |
| No Don't Known | 32% | 30% |
| Don't Know | 19% | 22% |
| Does worksite have written policy about breastfeeding? | | |
| Yes | 18% | 13% |
| No | 30% | 33% |
| Don't Know | 52% | 54% |
| | | |
| Cigarettes Smoked During Last 3 Months of Pregnancy | | |
| Not smoked in past 2 years | 77% | 71% |
| Zero cigarettes per day | 03% | 19% |
| 1-10 per day | 13% | 09% |
| 11-20 per day | 06% | 00% |
| More than 20 per day | 01% | 01% |
| Alcoholic Drinks During Last 3 Months of Pregnancy | | |
| Not had any alcoholic drinks in past 2 years | 23% | 15% |
| Zero drinks per week | 26% | 84% |
| 1 drink per week | 32% | 00% |
| 2 or more drinks per week | 41% | 00% |
| 2 of more drinks per week | 71/0 | 01/0 |

| Prenatal Care | State | County |
|--|------------|------------|
| Mother received prenatal care as early in the pregnancy as wanted | | |
| Yes | 93% | 92% |
| No | 07% | 08% |
| Where mother went for most of prenatal care | | |
| Private doctor's office or clinic | 81% | 85% |
| Midwife's office or home office from a midwife | 05% | 00% |
| Public clinic | 13% | 12% |
| Other | 01% | 04% |
| HIV/AIDS Testing During Pregnancy | | |
| Yes | 50% | 46% |
| No | 25% | 24% |
| No, Testing was offered, but did not want to be tested | 04% | 01% |
| Don't know | 22% | 28% |
| Received Flu Shot During Pregnancy | | |
| Yes | 62% | 50% |
| No, Received flu shot before became pregnant | 07% | 03% |
| No, Did not want flu shot during pregnancy | 13% | 19% |
| No, Doctor said to wait until after pregnancy | 01% | 01% |
| No, Do not ever get flu shots | 14% | 21% |
| No, Flu shot was not available | 02% | 06% |
| Hospital Services | State | County |
| Hospital Staff Encouraged Breastfeeding During First 24 Hours After Birth of the Baby | | |
| Yes | 83% | 74% |
| No | 15% | 24% |
| Don't remember | 03% | 03% |
| During Pregnancy or Time at the Hospital, a Doctor, Nurse, or Other | | |
| Healthcare Professional Provided Information or Scheduled a Time to | | |
| Discuss: Breastfeeding or breastfeeding techniques | 700/ | 020/ |
| Action steps to decrease frustration when infant cries | 78% 40% | 83% 61% |
| • | 49% | 61% |
| How to lay the baby down to sleep | 56% | 67% |
| Dangers of shaking an infant Did not receive any information and have not scheduled a time to discuss | 48% | 61% |
| Did not receive any information and have not scheduled a time to discuss | 09% | 07% |

Technical Note: Because percentages were rounded, the sum of values may not equal 100%. *For more information:* Call the University of Northern Iowa Center for Social & Behavioral Research: (319) 273-2105.

Data source: Iowa Barriers to Prenatal Care Project, Iowa Department of Public Health, University of Northern Iowa Center for Behavioral Research: 2014 Data.