

Immunization Schedule with Combination Vaccines

EVERY FALL: FLU VACCINE⁶ for anyone 6 months and older

	2 MONTHS	4 MONTHS	6 MONTHS	12 MONTHS	15 MONTHS	18 MONTHS	4-6 YEARS		
PEDIARIX[®] PROQUAD[®] QUADRACEL[™] or KINRIX[®]	PEDIARIX[®] DTaP, IPV, HepB + PCV Rotavirus Hib	PEDIARIX[®] DTaP, IPV, HepB ¹ + PCV Rotavirus Hib	PEDIARIX[®] DTaP, IPV, HepB + PCV Rotavirus ² Hib ³	HepA MMR ⁴ Varicella ⁴ PCV ⁵ Hib ⁵	DTaP	HepA	QUADRACEL[™] or KINRIX^{®7} DTaP, IPV + PROQUAD[®] MMRV		
	PENTACEL^{®8} PROQUAD[®] QUADRACEL[™] or KINRIX[®]	PENTACEL[®] DTaP, IPV, Hib + PCV Rotavirus HepB	PENTACEL[®] DTaP, IPV, Hib + PCV Rotavirus HepB ¹	PENTACEL[®] DTaP, IPV, Hib + PCV Rotavirus ² HepB	HepA MMR ⁴ Varicella ⁴ PCV ⁵	PENTACEL[®] DTaP, IPV, Hib	HepA	QUADRACEL[™] or KINRIX^{®7} DTaP, IPV + PROQUAD[®] MMRV	
		VAXELIS^{™8} PROQUAD[®] QUADRACEL[™] or KINRIX[®]	VAXELIS[™] DTaP, IPV, Hib, HepB + PCV Rotavirus	VAXELIS[™] DTaP, IPV, Hib, HepB ¹ + PCV Rotavirus	VAXELIS[™] DTaP, IPV, Hib ³ , HepB + PCV Rotavirus ²	HepA MMR ⁴ Varicella ⁴ PCV ⁵ Hib ⁵	DTaP	HepA	QUADRACEL[™] or KINRIX^{®7} DTaP, IPV + PROQUAD[®] MMRV

This is a suggested schedule using combination vaccines. For alternatives, details and catch-up, refer to the most current “Recommended Immunization Schedules” from CDC.

- 1 A dose of HepB vaccine is not necessary at 4 months if doses are given at birth and 2 months, but may be included as part of combination vaccine.
- 2 The 6 month dose is not needed if Rotarix is used exclusively for both dose 1 and 2 in rotavirus vaccine series.
- 3 The 6 month Hib dose is not needed if PedvaxHIB is used exclusively for the 2nd and 4th month doses.
- 4 CDC recommends administration of separate MMR and Varicella vaccines for the 12-15 month dose; however MMRV is acceptable based on clinical discretion.
- 5 ACIP recommends whenever feasible that the same manufacturer’s DTaP be used for each dose in the series, however vaccination should not be delayed based on the type of DTaP previously administered.
- 6 The child may need 1 or 2 doses of flu vaccine based on previous vaccination history. Refer to most current influenza recommendations.
- 7 Licensed for children 4-6 years. ACIP recommends using the same manufacturer’s DTaP vaccine for each dose in the series, but vaccination should not be deferred because the previous dose of DTaP is unavailable or unknown.
- 8 Licensed for children 6 weeks-4 years of age (prior to the 5th birthday).